

GENERAL INDEX—VOLUME 35

(A1 and A2 = Abstracts Supplements 1 and 2; E = Editorial; L = Letter to the Editor)

- AARONS L *see* Teh LS
- ABAD MA, GONZÁLEZ CM, LÓPEZ-LONGO FJ, ARIZA A, FERNÁNDEZ-ESPARTERO MC, GODOY H, RODRÍGUEZ-MAHOU M, CARREÑO L. Clinical significance of non-precipitating anti-nRNP antibodies in patients with systemic lupus erythematosus, (A1) 18
- ABAD MA *see also* López-Longo FJ; Monteagudo I
- ABDON P *see* McNicholas MJ
- ABDUL-WAHAB R *see* Lim KL
- ABE T. Letter from Japan, 579
- ABINUN M. Failure of first-line therapy with intravenous immunoglobulin in a child with scleromyositis, (L) 1029
- ABRAHAM DJ *see* Shiwen X
- ABU-SHAKRA M *see* Koh ET
- ACASUSO M *see* Cissal A
- ACEBES JC, HASSAN LA, OLIVÉ A, BAETHGE BA, ORIOL A, TENA X, LISSE JR. Systemic lupus erythematosus, comparison of disease activity of ethnic groups belong to 2 university hospitals, (A1) 34
- ACKROYD C *see* Carr AJ
- Acromegalic patients, articular complaints: a preliminary report on clinical and radiological features, (A1) 94
- Acromegaly, magnetic resonance imaging, (A1) 95
- Activity, 24hr in rheumatoid arthritis: comparison with HAQ, (A1) 233
- Acute rheumatology referral service, clinical need, 389
- ADAMS G *see* Chernajovsky Y
- ADAMS MA, BHATIA LS, DOLAN P. Sustained loading reduces the stress-equalising property of cartilage, (A1) 239
- ADAMS MA, CLINTON-WATKINS A, KERIN AJ, WISNOM MR. Sustained loading makes articular cartilage stronger, (A1) 240
- ADEBAJO AO. Dengue arthritis, (L) 909
- Adhesion mechanisms, role in the pathogenesis of chronic synovitis, (A1) 125, 1198
- Adhesion molecule expression
does lipocortin mediate the inhibition induced by glucocorticoids?, (A1) 64
glucocorticoid induced inhibition is reversed by IL-2, but not by TNF- α , IL-1 β /IL-4 or IL-10, (A1) 7
- Adhesion molecules, soluble, relationship with anti-endothelial cell antibodies in vasculitis, (A1) 52
- Adhesion of rheumatoid lymphocytes to mucosal endothelium: the gut revisited, 218
- Adolescents with rheumatic diseases and their transition into adulthood, (E) 80
- Adrenal function in AA amyloidosis, (A1) 76
- β_2 -Adrenergic receptors on synovial fluid lymphocytes and peripheral blood lymphocyte subsets in patients with rheumatoid arthritis, (A1) 65
- Adult Still's disease
cyclosporin A in the treatment, (A1) 77
DRB1 associations, (A1) 78
- Adulthood, transition into of adolescents with rheumatic diseases, (E) 80
- Adults, young with arthritis: meeting their transitional needs, 84
- AERSSENS J *see* Dequeker J
- Agalactosyl IgG
in aggregates from the rheumatoid joint, 335
[Gal(O)]—an analysis of its clinical utility in the long-term follow-up of patients with rheumatoid arthritis, 1063
is a marker of severe spondylitis, (A1) 153
- Age effects on the efficacy and tolerance of methotrexate in rheumatoid arthritis, 453
- Aggrecan
age-related changes in the C-terminal from human cartilage, (A1) 236
cartilage domains, production as recombinant fusion proteins to analyse recognition by autoreactive T lymphocytes, (A1) 184
and link protein mRNA quantitation in human articular cartilage of different ages by competitive RT-PCR, (A1) 237
- AGNEW JE *see* MacGregor AJ
- AHMED S *see* Salih AM
- AHMED Z, ARCHER JR, BROWN RA. Expression of the *tch* gene in mouse embryos, (A2) 8
- AHO K *see* Heliövaara M
- AHO S *see* Maillefert JF
- AIDS, successful treatment of Reiter's syndrome in a patient with methotrexate and corticosteroids, (L) 295
- AIMS2, evaluation of a Dutch version for patients with rheumatoid arthritis, 755
- AKAIL M *see* Evans MD
- AKBAR BC *see* Gordon C
- AKIL M, AMOS RS, STEWART P. Infertility may sometimes be associated with NSAID consumption, 76
- AKIL M *see also* Yeap SS
- AKIYA K *see* Nishikai M
- AKOĞLU T *see* Şahin (Özgün) Ş
- AL DALAAN A *see* El Ramahi KM; Karrar A
- AL JISHI F *see* Karrar A
- AL SHAIKH A *see* El Ramahi KM; Karrar A
- AL-AWADI A *see* Malaviya AN
- AL-GAURER S *see* Malaviya AN
- AL-JARALLAH K *see* Malaviya AN
- AL-MAZIDI M *see* Malaviya AN
- AL-MUGHALES J *see* McInnes IB
- AL-SAIED K *see* Malaviya AN
- AL-SALEM I *see* Malaviya AN
- ALAVI A *see* Homaei H; Pool AJ
- ALBORNOZ J, QUIJADA J, POVEDANO J, PÉREZ-VILCHEZ D, DE LA IGLESIA JL, ALBORNOZ J, GARCÍA-LÓPEZ A, FERNÁNDEZ-RODRÍGUEZ A. Comparison of tenoxicam and bromazepam in the treatment of fibromyalgia. A randomized, double-blind, placebo-controlled trial, (A1) 138
- ALBORNOZ J *see also* Albornoz J; De La Iglesia JL
- ALCANIZ C *see* Román JA
- ALCOCK J *see* Tunn AJ
- ALDABERTH T *see* McNicholas MJ
- ALEGRE C *see* Escóla A
- ALEXANDER G *see* Naughton M
- ALGRA D *see* Odling E
- ALI NN, KANZLER JM, SAHINOGLU T, BLAKE DR, STEVENS CR. rhTNF α -induced resorption of mouse calvariae *in vitro* is suppressed by xanthine oxidase inhibitors, (A1) 241
- ALISON MR *see* Ravirajan CT
- Alkaline phosphatase half-life in the comparison of potency between bisphosphonate compounds in the treatment of Paget's disease, (A1) 92
- ALLAIN J *see* Sarau A
- ALLARD SA, BARKLEY DEH, WOODROW DF, MAINI RN. Clinical, arthroscopic and histological features of synovitis in SLE, (A1) 36
- ALLEN IV *see* Kane D
- ALLEN PJ *see* Hogarth MB
- ALLEN RL, GILLESPIE GMA, EDMUNDS S, HALL MA, BOWMAN SJ, WORDSWORTH BP, LEIRISALO-REPO M, MCMICHAEL AJ, BOWNESS P. T lymphocyte expansions in blood and synovial fluid in reactive arthritis, (A1) 183
- ALLEN RL *see also* Bowness P
- ALMOGUERA MI *see* Carreño L
- ALONSO J *see* López-Longo FJ; Monteagudo I
- ALONSO RUIZ A *see* Perez-Ruiz F
- ALONSO-RUIZ A *see* Calabozo M
- Alternative/complementary medicine, a survey of the attitudes of rheumatologists and private medical insurers, (A1) 128
- ALVES J *see* Ames PRJ; Ruiz-Irastorza G
- AMAT M, DIAZ C, VILA L. Study of the cooperation between chondrocytes and polymorphonuclear leucocytes in leukotriene biosynthesis, (A1) 62
- Ambulation, factors affecting in rheumatoid arthritis, (A2) 20
- Ambulatory patients' attitudes towards a proposed telephone follow-up service, (A1) 130
- AMENGUAL O, ATSUMI T, KHAMASHTA MA, HUGHES GRV. Beta 2-glycoprotein I antibody specificity in patients with the antiphospholipid syndrome, (A1) 24
- AMENGUAL O, ATSUMI T, KHAMASHTA MA, KOIKE T, HUGHES GRV. Specificity of ELISA for antibody to β_2 -glycoprotein I in patients with antiphospholipid syndrome, 1239
- American College of Rheumatology
core set and other validated indices of disease activity in RA, comparison of Overall Status in Rheumatoid Arthritis, (A1) 196
core set and SF-36 in rheumatoid arthritis—physical and social functioning measures of the SF-36 are strongly associated with the Health Assessment Questionnaire, (A1) 195
- AMÉRIGO MJ *see* Brito M
- AMES P *see* Méndez MJ
- AMES PRJ, BRILLANTE M, IANNACCONE L, CIMINO R, LUPOLI S, BRANCACCIO V. Fibrinolysis abnormalities in primary antiphospholipid syndrome: sex dependency and clustering with low free protein S, (A1) 38
- AMES PRJ, PYKE S, ALVES J, RAMOS P, KHAMASHTA MA, HUGHES GRV. Fibrinogen in systemic lupus erythematosus: an atherothrombotic link?, (A1) 122

- AMES PRJ, ROES L, LUPOLI S, PICKERING M, BRANCACCIO V, KHAMASHTA MA, HUGHES GRV. Thrombosis in Churg-Strauss syndrome. Beyond vasculitis, 1181
- AMES PRJ *see also* Lútrano E; Steuer A
- Amino acid concentrations, decreased plasma in fibromyalgia patients, (A1) 137
- Aminohexane bisphosphonate suppresses bone turnover in postmenopausal women more rapidly than oestrogen-gestagen therapy, 636
- Amitriptyline, placebo-controlled study in ankylosing spondylitis, (A1) 155
- AMOR B *see* Cisin A; Ravaud P
- AMOS N *see* Hall FC; Richards PJ
- AMOS RS *see* Akil M; Till SH
- AMOUROUX J *see* Guillevin L
- Amyloid, inappropriately high ESR results—a feature complicating rheumatic disorders?, (A1) 77
- Amyloidosis
- AA, adrenal function in, (A1) 76
 - AA in RA and JCA, (A1) 76
 - AL, intensive chemotherapy, (A1) 12
 - and giant cell arteritis, lack of association between, (A1) 57
 - new roles for organ transplantation, (A1) 12
 - secondary in patients with rheumatoid arthritis: diagnostic and prognostic value of gastroduodenal biopsy, 44
- AN TN, KHOA ND. Rheumatic diseases and the development of rheumatology in Vietnam, 680
- Anaemia, chronic of rheumatoid arthritis, evidence for abnormal erythroblast iron metabolism, (A1) 191
- Anatomic index, video image analysis of hands: development as a potential outcome measure in rheumatoid arthritis, 1274
- ANCA in RA patients, (L) 1328
- ANDERSON ME, HOLLIS S, MOORE T, JAYSON MIV, HERRICK AL. Non-invasive assessment of vascular reactivity in forearm skin of patients with primary Raynaud's phenomenon and systemic sclerosis, 1281
- ANDERSON ME, MOORE TL, ROSSER S, JAYSON MIV, HERRICK AL. Digital neurovascular mechanisms in primary Raynaud's phenomenon and systemic sclerosis as assessed by iontophoresis and laser Doppler techniques, (A1) 48
- ANDREU JL, LABIANO I, PEÑA F, ORTIZ-BERROCAL J, PÉREZ-PÁRAMO M, LARREA A, MULERO J. Isonitrite-Tc99 bone scan vs radiologic evaluation in the assessment of the malignant nature of a tumour-like bone lesion, (A1) 241
- ANDREU JL *see also* Díez E; Mulero J
- ANDREWS D *see* Devlin J
- Angiogenesis: a critical process in the pathogenesis of RA—a role for VEGF, (E) 917
- Angiotensin converting enzyme, synovial in human arthritis, (A1) 66
- ANILKUMAR TV *see* Ravirajan CT
- Animal models of rheumatoid arthritis, (E) 1041
- Ankylosing spondylitis
- acute physiological effect of hydrotherapy, (A1) 156
 - antibodies to *Klebsiella*, *Proteus* and the peptides HLA-B27 and HLA-DR4 (DW15) in Japanese patients, (A2) 12
 - assessment of disease activity measures: their relationship with acute phase reactants, (A1) 153
 - autoantibodies: molecular analysis of the antigen, (A1) 152
 - Bath patient global score (BAS-G), 66
 - biochemical markers of bone turnover, (A1) 151
 - cervical spondylodiscitis, (L) 1034
 - compliance monitoring of NSAID drug therapy, experiences with an electronic monitoring device, 60
 - disease activity: selection of a core set of variables and a first step in the development of a disease activity score, 867
 - do childhood infections affect the development, (A2) 23
 - does genetic anticipation occur in?, (A2) 24
 - effect of 30 min cycle ergometry, 167
 - epidemiology of spondylodiscitis in—a controlled study, 660
 - ethnic differences, (A1) 149
 - evolution of the pulmonary function, (A1) 151
 - immunohistologic analysis of peripheral joint disease, (A1) 154
 - intracellular calcium release in T-lymphocytes, (A1) 11
 - jejunal lesions are due to non-steroidal anti-inflammatory drug therapy, (A1) 157
 - methotrexate in severe: clinical and serological response compared with rheumatoid arthritis and psoriatic arthritis, (A1) 155
 - Paget's disease in a patient with—a diagnostic dilemma, 1011
 - placebo-controlled study of low dose amitriptyline, (A1) 155
 - and psoriasis, atypical antineutrophil cytoplasmic antibodies (ANCA), (A1) 161
 - radiological outcome: use of the Stoke Ankylosing Spondylitis Spine Score (SASSS), 373
 - radiology in: comparison of two scoring systems, (A2) 23
 - risk factors for worse disease activity and function, (A2) 24
 - serum prolactin and salivary testosterone in young and middle-aged men, (A1) 152
 - similarly increased serum IgA1 and IgA2 subclass antibody levels against *Klebsiella pneumoniae* bacteria with/without extra-articular features, 125
 - spinal pseudoarthrosis complicating: a report of two patients, 1319
 - treatment of chronic painful focal lesions with low-dose radiotherapy, (A1) 156
- Anterior pituitary function in patients with newly diagnosed rheumatoid arthritis, 350
- Anti- β 2-glycoprotein I, specificity of ELISA in patients with antiphospholipid syndrome, 1239
- Anti-CD4 and anti-IL-6 monoclonal antibodies combination therapy in a case of severe spondylarthropathy, (L) 1331
- Anti-CD8 treatment reduces the severity of inflammatory arthritis, but not vasculitis, in mercuric chloride-induced autoimmunity, (A1) 10
- Anti-CD44 antibody inhibits human lymphocyte adhesion to endothelial cells and synoviocytes by inducing proteolytic cleavage of CD44, (A1) 60
- Anti-endothelial cell antibodies
- characterization in systemic lupus erythematosus by controlled immunoblotting, 1231
 - longitudinal study of activity in systemic sclerosis, (A1) 46
 - relationship with soluble adhesion molecules in vasculitis, (A1) 52
- Anti-ganglioside antibodies in patients with rheumatoid arthritis complicated by peripheral neuropathy, 725
- Anti-IL-6 and anti-CD4 monoclonal antibodies combination therapy in a case of severe spondylarthropathy, (L) 1331
- Anti-keratin antibodies
- anti-RA33 antibody, antiperinuclear factor and antinuclear antibody, diagnostic value in early rheumatoid arthritis: comparison with rheumatoid factor, 620
 - disease activity and radiological damage in rheumatoid arthritis, (A2) 14
- Anti-nRNP antibodies, non-precipitating, clinical significance in patients with systemic lupus erythematosus, (A1) 18
- Anti-p57, sero-epidemiology and correlations with HLA, (A1) 24
- Anti-RA33 antibody, antikeratin antibody, antiperinuclear factor and antinuclear antibody, diagnostic value in early rheumatoid arthritis: comparison with rheumatoid factor, 620
- Anti-RNP antibody, high titre in association with carcinoma in a child who did not have a rheumatic disease, (A2) 37
- Anti-Sm antibodies detected by ELISA, diagnostic relevance in systemic lupus erythematosus, (A1) 19
- Anti-TCRBV monoclonal antibodies in the detection and monitoring of large granular lymphocyte leukaemia in Felty's syndrome, (A1) 182
- Anti-tumour necrosis factor alpha antibody, engineered human, reduces serum interleukin-6 and urine bone markers but has no effect on soluble CD4 and soluble interleukin-2 receptor in rheumatoid arthritis, (A1) 214
- Antibodies
- defective production in patients with rheumatoid arthritis and bronchiectasis, (A1) 201
 - levels to *Clostridium perfringens* and response to sulphasalazine, (L) 1327
 - to endothelial cells and to β 2-glycoprotein I in the antiphospholipid syndrome: prevalence and isotype distribution, 523
 - to four Gram-negative bacteria in rheumatoid arthritis which share sequences with the rheumatoid arthritis susceptibility motif, 592
 - trispecific stimulation, rheumatoid T cell functional deficiency can be reversed with, (A1) 13
- Anticardiolipin antibodies
- human monoclonal and β 2-glycoprotein I effects on the protein S / C4B-binding protein system, (A1) 19
 - human monoclonal IgG, identification of idiotypes on and in the sera of patients with autoimmune rheumatic diseases, (A1) 22
 - and renal disease in patients with systemic lupus erythematosus, (A') 20
- Antigen-presenting cells but not lymphocytes in the joint may indicate the cause of reactive arthritis, 1082
- Antileukoproteinase serine proteinase inhibitor specifically accumulates in normal but not in arthritic cartilage, (A1) 68
- Antineutrophil cytoplasmic antibodies
- abnormal angiograms and pathological findings in polyarteritis nodosa and Churg-Strauss syndrome, 958
 - are they associated with spondylarthropathies?, (L) 1032
 - atypical in patients with ankylosing spondylitis and psoriasis, (A1) 161
 - in rheumatoid arthritis patients, 38
 - in systemic lupus erythematosus, 625
- Antineutrophil cytoplasmic antibody-positive polyarthritis associated with minocycline therapy, (L) 1327
- Antineutrophil cytoplasmic autoantibodies, atypical Cogan's syndrome associated with, (L) 601
- Antinuclear antibody, anti-RA33 antibody, antikeratin antibody and antiperinuclear factor, diagnostic value in early rheumatoid arthritis: comparison with rheumatoid factor, 620
- Antinuclear and antinuclear antibodies in patients with scleroderma-polymyositis overlap syndrome, (L) 1326

- Antiperinuclear factor
anti-RA33 antibody, antikeratin antibody and antinuclear antibody, diagnostic value in early rheumatoid arthritis: comparison with rheumatoid factor, 620
in spondylarthropathies, (L) 804, (L) 805
- Antiphospholipid antibodies
from patients infected with human B19 parvovirus have similar specificity and co-factor dependence to those found in SLE patients, (A1) 22
screen, (A1) 23
- Antiphospholipid syndrome
antibodies to beta-2-glycoprotein I recognise a conformational epitope and are highly specific, (A1) 21
antibodies to endothelial cells and to β 2-glycoprotein I in the antiphospholipid syndrome: prevalence and isotype distribution, 523
beta 2-glycoprotein I antibody specificity, (A1) 24
fibrinolysis abnormalities: sex dependency and clustering with low free protein S, (A1) 38
specificity of ELISA for antibody to β 2-glycoprotein I, 1239
and systemic lupus erythematosus, obstetric outcome: the St Thomas' Hospital Lupus in Pregnancy clinic experience, (A1) 123
- Antiprotease antibody, phosphatidyl serine-dependent is exclusive to patients with lupus anticoagulant, 589
- α 1-Antitrypsin and IgA, normal urine provides an environment that augments complex formation between, (A2) 13
- ANTOLIN J see Cantabrana A
ANTÓN R see Camacho M
ANTTILA P see Kauppi M
AOYAGI K see Kobayashi H
- Apolipoproteins, changes in levels following low-calorie dieting in hyperuricemic patients, (A1) 145
- Apoptosis analysis in lymphoid organs and lupus disease in murine systemic lupus erythematosus, (A1) 40
- APPERLEY JF see Persey MR
ARA 1987 criteria do not perform well in early RA, (A2) 18
ARAGON A. Familial *Salmonella*-triggered reactive arthritis, (L) 908
ARCHER JR. Mapping of *tch*, a mouse dwarfing gene, (A2) 9
ARCHER JR see also Ahmed Z
ARDEN N see Etherington J
ARDEN NK, GRIFFITHS G, HART DJ, DOYLE DV, SPECTOR TD. Association between radiographic osteoarthritis and osteoporotic fractures: the Chingford study, (A1) 116
ARDEN NK, GRIFFITHS GO, HART DJ, DOYLE DV, SPECTOR TD. Association between osteoarthritis and osteoporotic fracture: the Chingford study, 1299
ARDEN NK see also Dolan AL
ARETXABALA I see Calabozo M; Perez-Ruiz F
ARFI S see Le Thi Huong D
ARIZA A see Abad MA; González M
ARIZA J see Fitter J
ARMADANS L see Lience E
ARMONA J see González-Gay MA
ARMSTRONG C see Luqmani RA
ARMSTRONG GP, WHALLEY GA, DOUGHTY RN, GAMBLE GD, FLETT SM, TAN PLJ, SHARPE DN. Left ventricular function in scleroderma, 983
ARNALICH F see De Miguel E
ARNOLD MM see Green M; Mason JC
ARO H see Roivainen A
AROMAA A see Heliövaara M
Arteritis temporalis, reduced CD8+ T-cell concentration in peripheral blood of patients with carotid artery stenosis, 649
Arthritic joint, electron paramagnetic resonance study, (A1) 204
Arthritis
acute, tenidap reduces cell infiltration and MCP-1 mRNA expression in a rabbit model, (A1) 73
adjuvant, synergistic interaction between morphine and indomethacin attenuates pain sensation and tissue destruction, (A1) 75
adjuvant-induced, role of endogenous serotonin in the rat, 112
associated with monoclonal gammopathy: clinical characteristics, 241
chronic
autoimmune induction in DBA/1 mice by oral administration of type II collagen and *Escherichia coli* lipopolysaccharide, 828
single-blind randomized controlled trial of an educational booklet for patients, 775
tenidap decreases synovial growth (hyperplasia and increased extra-cellular matrix production) and cell infiltration in rabbits, (A1) 72
of coeliac disease: prevalence and pattern in 200 adult patients, 1314
collagen-induced
amelioration following treatment with rolipram, (A1) 75
modification of murine with antibodies to interleukin 12, (A1) 108
pattern of cytokine secretion during development, (A1) 5
prevention by a non-depleting anti-CD4 antibody, (A1) 108
pro-inflammatory cytokine expression, (A1) 6
community based self-management courses for older people, (A2) 31
dengue, (L) 909
elevated levels of corticotrophin-releasing factor binding protein in the blood of patients and the presence of novel ligands in synovial fluid, 120
experimental models of acute and chronic, antiinflammatory effects of two new platelet activating factor receptor antagonists, (A1) 13
inflammatory
does it affect sexuality?, (A2) 19
evaluation of disease and drug monitoring, (A1) 230
handicap in, 891
lymphatic function, (L) 396
reversible ovulatory failure associated with development of luteinized unruptured follicles in women taking non-steroidal anti-inflammatory drugs, 458
use of multi-bacterial nested PCR to screen synovial fluid and membrane samples, (A1) 113
low incidence in persistent human parvovirus B19 infection in immunocompromised HIV-infected patients, (A1) 167
Lyme, Lyme borreliosis in rheumatological practice: identification and diagnostic aspects in a Swedish county with high endemicity, 853
mono-articular, administration of local and systemic soluble complement receptor one (SCR1) has a beneficial effect on development, (A1) 74
murine collagen-induced, inhibition by intranasal administration of collagen type II peptides, (A1) 73
oncoprotein expression in human synovial tissue: an immunohistochemical study of different types of arthritis, 933
rat antigen-induced, single intra-articular injection of liposomally conjugated methotrexate suppresses joint inflammation, 719
reactive
antigen-presenting cells but not lymphocytes in the joint may indicate the cause, 1082
 β -haemolytic *Streptococcus* and *Staphylococcus aureus*, (L) 912
familial *Salmonella*-triggered, (L) 908
HLA-B27-restricted peptide presentation to cytotoxic T cells, (A1) 114
nature of the synovial infiltrate in the synovial membrane, (A1) 163
T lymphocyte expansions in blood and synovial fluid, (A1) 183
temporomandibular joint in the rabbit, mapping abnormal synovial vascular permeability using MRI, (S3) 24
type II collagen-induced transfer from DBA/1 to SCID mice can be prevented by blockade of MAC-1, (A1) 74
- Arthropathy
inflammatory, intra-articular pressure dynamics, (A1) 71
ochronotic: a non-rheumatological series of 16 patients, (A1) 95
Arthroplasty, lower limb, use of steroids in patients undergoing, (A1) 209
Arthroscopic synovectomy in rheumatoid and psoriatic knee joint synovitis: long-term outcome, 463
Arthrotec 75 and Diclofenac 75, comparison of effect on patient activity using the Numact activity monitor, (A2) 33
- Articular cartilage
in vitro effects of methotrexate, 342
and synovial tissue metabolism, serum markers in patients treated with prednisolone for early rheumatoid arthritis, (A1) 238
- Articular complaints in acromegalic patients: a preliminary report on clinical and radiological features, (A1) 94
- Articular destruction mechanisms in the feet and hands may differ, (A1) 192
- ARTLETT CM, BLACK CM, BRIGGS DC, STEPHENS C, WELSH KI. DNA allelic alterations within VNTR loci of scleroderma families, 1216
- ARTLETT CM, BLACK CM, BRIGGS DC, STEVENS CO, WELSH KI. Telomere reduction in scleroderma patients: a possible cause for chromosomal instability, 732
- Ascorbic acid absorption in patients with systemic sclerosis, (A1) 43
- ASENSI E see Escóla A; Obach J
ASPE B see Graña J
Aspirin-induced stomach erosions, effects of an amino-versus and alkyl-bisphosphonate on healing in healthy subjects, (A2) 7
- ASTORGA R see De La Iglesia JL
ATERO F see Rodriguez R
ATHANASSIOU P, NORSWORTHY P, WALPORT MJ, DAVIES KA. Antibodies to C1q(CLR) are markers for severe disease in systemic lupus erythematosus, (A1) 21
ATIEA G, KAY L, HANFY M, WALKER DJ. Disease expression and HLA association of RA: comparison of Egypt and the UK, (A1) 185
- ATKINS M see Fox RA
ATKINS RM see Perry MJ
ATKINSON P see Mishra MB
Atlantoaxial subluxation, rheumatoid, stiff collar for treatment, 771
ATRA E see Ciconelli RM

- ATSUMI T, KHAMASHTA MA, ICHIKAWA K, KOIKE T, HUGHES GRV. Effect of β 2-glycoprotein I and human monoclonal anticardiolipin antibodies on the protein S / C4B-binding protein system, (A1) 19
 ATSUMI T *see also* Amengual O
- Audit
 of domiciliary visits for locomotor problems, (A2) 30
 of guidelines for treatment and prevention of osteoporosis, (A1) 81
 of muscle biopsies, (A1) 128
 of needle arthroscopy service, (A1) 127
 of shared care between hospital and general practice in the management of rheumatoid arthritis, (L) 398
- AULELEY G-R *see* Ravaut P
- AUSTIN CA *see* Peel NFA
- Australia, the 1994-5 Michael Mason Travelling Fellowship: goodbye the Commonwealth, hello Australia, 1324
- Autoantibodies in ankylosing spondylitis: molecular analysis of the antigen, (A1) 152
- Autopsy, cause of death in 22 patients with rheumatoid arthritis, (A2) 22
- AVERNS HL, HALL J, WEBLEY M. Keratoconjunctivitis sicca in rheumatoid arthritis: prevalence in an outpatient population and role of Opticare eye drop dispenser, (A2) 20
- AVERNS HL, OXTORY J, TAYLOR HG, JONES PW, DZIEDZIC K, DAWES P. Radiological outcome in ankylosing spondylitis: use of the Stoke Ankylosing Spondylitis Spine Score (SASSS), 373
- AWADALLAH F, DALZIEL M, AXFORD JS. Expression of the Lewis X trisaccharide on serum IgG in rheumatoid arthritis, (A1) 175
- AXFORD JS *see* Awadallah F; Dalziel M; Homaei H; Pool AJ
- AYED K *see* Hamzaoui A
- AYRAL X, MAYOUX-BENHAMOU A, DOUGADOS M. Proposed scoring system for assessing synovial membrane abnormalities at arthroscopy in knee osteoarthritis, (S3) 15
- Azathioprine and sulphasalazine combination therapy, (L) 493
- B19 parvovirus, antiphospholipid antibodies from patients have similar specificity and co-factor dependence to those found in SLE patients, (A1) 22
- B lymphocytes, facilitated expression by synovial fibroblasts of factors involved in maturation and survival, (A1) 110
- BABA H *see* Wada M
- BACIC G *see* Poleksic L
- Bacillary angiomatosis, bone, in an HIV-infected patient, 901
- Back extensor muscles
 muscle fibre size and distribution: relationship with muscle fatigability, (A1) 144
 normal muscle fibre size and type distribution, (A1) 144
- Back pain
 in children, 1022
 future, employment and work activities as predictors, (A1) 141
 graded status, pain coping strategies as predictors for: results of a population based follow-up study, (A1) 142
 in Great Britain and Germany, (A1) 142
 relationship with vertebral osteophytosis, (A1) 143
 value of SPECT scans in identifying pain likely to benefit from facet joint injection, 1269
- Backache, school Christmas shows... 'look out the backache is behind you', (L) 1335
- BACKER V *see* Johansen JS
- BACON P *see* Exley A
- BACON PA *see* Carruthers DM; Gordon C; Luqmani RA; Pugh MT
- Bacterial urease increased activity in faeces in juvenile chronic arthritis: evidence of altered intestinal microflora?, 689
- BADLEY EM *see* Fitzpatrick F
- BAERWALD CGO, LAUFENBERG M, SPECHT T, WAHLE M, EHLENZ K, WICHERT P VON, KRAUSE A. β 2-adrenergic receptors on synovial fluid lymphocytes and peripheral blood lymphocyte subsets in patients with rheumatoid arthritis, (A1) 65
- BAERWALD CGO, PANAYI GS, LANCHBURY JS. Three novel polymorphisms in the regulatory region of the corticotropin releasing hormone gene in patients with rheumatoid arthritis and controls, (A1) 107
- BAERWALD CGO, WAHLE M, ULRICH S, JONAS D, WICHERT P V, KRAUSE A. Reduced influence of catecholamines on proliferation of lymphocytes from patients with rheumatoid arthritis *in vitro*, (A1) 65
- BAETHGE BA *see* Acebes JC
- BAIGENT S *see* Woods RJ
- BAILDAM E *see* Holt PJL
- BAILDAM EM, KELSEY A, HOLT PJL. Polyarteritis nodosa in two brothers presenting at the same age, (A2) 38
- BAILDAM EM *see also* Stathopoulou E
- BAKER E *see* Teh LS
- BAKER J *see* Spector TD
- BALAKRISHNAN C, JONES SM, EVISON G, MCHUGH NJ. Hand radiology in psoriatic arthritis: a five year follow-up study, (A1) 158
- BALAKRISHNAN C *see also* Flora RS; Jones SM; Taylor AL
- BALDOVIN M *see* Fiocco U
- BALDWIN D *see* Preedy VR
- BALLO R *see* Beighton PH
- BALSA A, DIXEY J, SANSOM DM, MADDISON PJ, HALL ND. Differential expression of the costimulatory molecules B7.1 (CD80) and B7.2 (CD86) in rheumatoid synovial tissue, 33
- BALSA A *see also* Uson J
- BAMJI AN, ERHARDT CC, PRICE TR, WILLIAMS PL. Painful shoulder: can consultants agree?, 1172
- BANBURY P *see* Exley A
- BANKHEAD CR *see* Harrison BJ; Symmons DPM
- BANNISTER G *see* Carr AJ
- BANSAL S *see* Tiwana H
- BARCELÓ P *see* Wojtulewski JA
- BAREN A *see* Ramos P
- BARKLEY DEH *see* Allard SA
- BARLOW JH, PENNINGTON DC. Randomised controlled study of information leaflets for people with rheumatoid arthritis, (A2) 32
- BARLOW JH, WILLIAMS B, WRIGHT C. Community based arthritis self-management courses for older people, (A2) 31
- BARNES J *see* Mackay K
- BARON D *see* Sarau A
- BARRATT JA, Scurry, osteoporosis and megaloblastic anaemia due to alleged food intolerance, (L) 701
- BARRETT EM, SYMMONS DPM, SCOTT DGI. Employment attrition in a community based inception cohort of rheumatoid arthritis patients, (A1) 235
- BARRETT EM *see also* Harrison BJ; Symmons DPM
- BARRINGTON NA *see* Peel NFA
- BARTOSIK I. Intermittent iloprost infusion therapy of pulmonary hypertension in scleroderma—a pilot study, (L) 1187
- BAS S, CUNNINGHAM T, KVIEK TK, GLENNÄS A, MELBY K, VISCHER TL. Synovial fluid and serum antibodies against *Chlamydia* in different forms of arthritis: intra-articular IgA production in *Chlamydia* sexually acquired reactive arthritis, 548
- BAS S, CUNNINGHAM T, KVIEK TK, GLENNÄS A, MELBY K, VISCHER TL. Value of isotype determination of serum antibodies against *Chlamydia* for the diagnosis of *Chlamydia* reactive arthritis, 542
- BAŞARAN G *see* Yazici H
- BASSIOUNI M *see* Troughton PR
- Bath ankylosing spondylitis patient global score (BAS-G), 66
- BAVINGTON C, WRIGHT MO, JOBANPUTRA P, LIN H, MORRISON PT, BRENNAN FR, SALTER D, NUKI G. Accelerated proteoglycan synthesis in human chondrocytes following cyclical pressurisation is mediated by β 1 integrin and stretch activated ion channels, (A1) 110
- BAVINGTON C *see also* Brennan FR; Jobanputra P
- BAYLISS M *see* Howat S; Venn G
- BAYLISS MT *see* Bolton MC; Duddhia J; McAlinden A; McKenna LA
- bcl-2 expression and peripheral lymphocyte apoptosis in systemic lupus erythematosus: correlation with disease activity, (A1) 39
- bcl-2 expression in rheumatoid arthritis, 611
- bcl-2 proto-oncogene overexpression in rheumatoid synovium, (L) 803
- BEGLEY J *see* Dunne CA
- BEGUM S *see* Koutedakis Y
- Behçet's disease
 coagulation factor V gene mutation increases the risk of venous thrombosis, 1178
 endothelin-1 expression in serum and bronchoalveolar lavage, 357
 magnetic resonance imaging shows brain lesions in patients without central nervous system symptoms, (A1) 52
 monocyte activity, 424
- Behçet's syndrome
 neurological involvement in the UK, (A2) 28
 ten-year mortality, 139
- BEIGHTON PH, BALLO R, VILJOEN D, RAMESAR R. Mseleni joint disease, (A1) 106
- BEKKELUND SI, MELLGREN SI, PRØVEN A, HUSBY G. Quantified neurological examination with emphasis on motor and sensory functions in patients with rheumatoid arthritis and controls, 1116
- BELL A *see* Knight J
- BELL AL, MCNALLY J, MAGILL K, IRVINE AE. Progressive loss of reactive oxygen species generation in neutrophils undergoing apoptosis: an antiinflammatory effect in synovitis, (A1) 61
- BELL AL *see also* Smyth AE
- BELL F *see* Hamilton J
- BELL J *see* Hardwick LJ; Mishra MB
- BELL L *see* Dasgupta B
- BELLAICHE L *see* Lioté F
- BELMONT HM *see* Markov G
- BELMONTE MA, BELTRAN J, FURIÓ MP. Comparative study of the short vs Stanford versions of the Spanish HAQ. Correlation between Likert and VAS scales for pain and health status, (A1) 232
- BELTRAN J *see* Belmonte MA

- BELTRANDI E *see* Boiardi L
- BELZUNEGUI J, GONZÁLEZ C, FIGUEROA M, DELVAL M, ERAUSQUIN C, RODRIGUEZ-VALVERDE V. Infectious spondylodiscitis: a revision of 98 cases. (A1) 166
- BENDIX G, BIELLE A. Adding low-dose cyclosporin A to parenteral gold therapy in rheumatoid arthritis: a double-blind placebo-controlled study. 1142
- BENNETT RJM *see* Teh LS
- Benzbromarone and allopurinol, comparative study of efficacy for the therapy of hyperuricemia in chronic primary gout. (A1) 145
- BERENBAUM F. Successful treatment of Reiter's syndrome in a patient with AIDS with methotrexate and corticosteroids. (L) 295
- BERESFORD JN *see* Minaur NJ
- BERGIN C *see* Ormond P
- BERGLUND J, BLOMBERG I, HANSEN BU. Lyme borreliosis in rheumatological practice: identification of Lyme arthritis and diagnostic aspects in a Swedish county with high endemicity. 853
- BERNARD J *see* Coombes G
- BERNHARDT M *see* Ghoslan PR
- BERRY H *see* Houssein DA; Houssien DA; Smith EC
- BERTHOLET J-M *see* Maugars Y
- BERTI A *see* Leak AM
- BEVIS PJR, BIRD HA, LAPHAM G
Open study to assess safety and tolerability of meloxicam 15 mg in subjects with rheumatic disease and mild renal impairment. (S1) 56
Study to assess the tolerability and safety of meloxicam in patients with rheumatoid arthritis who have mild renal impairment. (A1) 223
- BEVIS PJR *see also* Wojtulewski JA
- BEXENDINE S *see* Shearing-Lynne C
- BEYELER C, JORDI B, GERBER NJ, IM HOF V. Pulmonary function in rheumatoid arthritis treated with low-dose methotrexate: a longitudinal study. 446
- BEYELER C *see also* Lehmann T
- BEYNON HLC *see* Davies KA
- BEYNON K *see* Ocal L
- BHALLA AK *see* McFarlane XA; Minaur NJ
- BHATIA LS *see* Adams MA
- BHATT B *see* Sahinoglu T
- BHAYANI H *see* Sibanda EN
- BHRF1, rheumatic disease patients, prone to Sjögren's syndrome and/or lymphoma, mount an antibody response to BHRF1, the Epstein-Barr viral homologue of BCL-2. 1075
- BIAGINI P, PIPITONE N, GOULDING NJ, LEE A, GERLI R, PANAYI GS, PITZALIS C. Does lipocortin mediate the inhibition of adhesion molecule expression induced by glucocorticoids? (A1) 64
- BIAGINI P *see also* Pipitone N
- BIDDLE P *see* DYCE C
- BIJLSMA JWI *see* van der Veen MH; Van Valburg AA; Verhoef CM
- BILAG disease activity index, further validity in patients with systemic lupus erythematosus. (A1) 29
- BILLINGHAM MEJ, MEIJERS T, MAHWINNEY B, MALCOLM A. Spontaneous osteoarthritis in guinea pigs: cartilage degeneration is preceded by loss of subchondral trabecular bone. (A1) 104
- BILLINGHAM MEJ *see also* Fernihough JK
- BINDER A *see* Tiwana H
- Binding proteins, production and role of the insulin-like growth factor I binding protein 3 in human articular cartilage explants. 515
- BINNS RM *see* Chapman PT
- Biochemical markers and disease activity, relationships between clinical measures in rheumatoid arthritis. (A1) 230
- BIRD C *see* Buckland-Wright JC
- BIRD H *see* Troughton PR
- BIRD HA *see* Bevis PJR
- BIRKELUND S *see* Campbell F
- BIRKENSHAW G *see* MacDonald AG
- BIRKHOLDER J *see* Sakkas LI
- BIRRELL FN, HASSELL A, DAWES PT
American College of Rheumatology core set and SF-36 in rheumatoid arthritis—physical and social functioning measures of the SF-36 are strongly associated with the Health Assessment Questionnaire. (A1) 195
Why not use OSRA? Comparison of Overall Status in Rheumatoid Arthritis with American College of Rheumatology core set and other validated indices of disease activity in RA. (A1) 196
- Bisphosphonate
amino-versus and alkyl-bisphosphonate effects on the healing of aspirin-induced stomach erosions in healthy subjects. (A2) 7
comparison on TNF α production by PBMCs. (A1) 4
- BJARNASON I *see* Mahmud T
- BUELLE A *see* Bendix G
- BLAS St *see* Specker Ch
- BLACK AJ, CHARLWOOD C, DURHAM B, FRASER WD. Timed therapy with calcium and phosphate supplements restores phosphate and parathyroid hormone rhythms in patients with osteoporosis, with an anabolic effect on spinal bone mineral density. (A1) 88
- BLACK C *see* Bryan C; McHugh NJ; Stratton R
- BLACK CM *see* Artlett CM; Bryan C; Denton CP; MacGregor AJ; Shiwen X
- BLADES S *see* Blake DR; Gaffney K
- BLAKE D. Training in rheumatology. (L) 1190
- BLAKE DR, BLADES S, COUMBE A, MAPP PI. Nitration of tyrosine in the inflamed human synovium: evidence for the generation of peroxynitrite *in vivo*. (A1) 70
- BLAKE DR. Measuring outcome in rheumatoid arthritis—which measures are suitable for routine clinical use. (L) 301
- BLAKE DR *see also* Ali NN; Edmonds SE; Gaffney K; Jawed S; Naughton DP; Nazhat NB; Sahinoglu T; Walsh DA; Zhang Z
- BLANCO R, MARTÍNEZ-TABOADA V, GARCÍA-FUENTES M, GONZÁLEZ-GAY MA, RODRIGUEZ-VALVERDE V. Henoch-Schönlein purpura in adulthood: a more severe syndrome with a good outcome. (A1) 55
- BLANCO R, MARTÍNEZ-TABOADA V, GONZÁLEZ-GAY MA, SÁNCHEZ-ANDRADE A, RODRIGUEZ-VALVERDE V. Successful therapy with danazol in refractory thrombocytopenia associated with autoimmune diseases. (A1) 55
- BLANCO R *see also* González-Gay MA; Rodríguez-Valverde V
- BLAND M *see* Casey ATH
- BLANN AB *see* Chaloner C
- BLANN AD, SANDERS P, HERRICK A, JAYSON MIV. Reduced levels of soluble L-selectin in systemic sclerosis and vasculitis. (A1) 42
- Blindness and low-dose corticosteroids in giant cell arteritis. (L) 702
- BLOMBERG J *see* Berglund J
- Blood flow responses, digital and microvascular in patients with systemic sclerosis and primary Raynaud's phenomenon. (A1) 42
- BLUHMKI E *see* Distel M; Ghoslan PR; Hosie J; Huskisson EC; Linden B; Register JY; Wojtulewski JA
- BLYTH T *see* Gordon MM
- BLYTH TH *see* Hunter JA
- BODMAN-SMITH K, SUMAR N, SINCLAIR H, ROITT I, ISENBERG D, YOUNG A. Agalactosyl IgG [Gal(o)]—an analysis of its clinical utility in the long-term follow-up of patients with rheumatoid arthritis. 1063
- BODMAN-SMITH K *see also* Kjeldsen-Kragh J
- Body composition and weight loss in rheumatoid arthritis. (A1) 200
- BOERBOOMS A *see* Stolk J
- BOERBOOMS AMT *see* van den Hoogen FHJ
- BOHEMEN EK. Flatfoot in pregnancy. (L) 396
- BOIARDI L, SALVARANI C, MACCHIONI P, CASADEI MALDINI M, MANCINI R, BELTRANDI E, PORTIOLO I. CD8 lymphocyte subsets in active polymyalgia rheumatica: comparison with elderly-onset and adult rheumatoid arthritis and influence of prednisone therapy. 642
- BOIVIN C *see* Devlin J
- BOLGER T, COYLE M, POWER C, FITZGERALD O, MCCARTHY D, BRESNIHAN B. Antiphospholipid antibody screen. (A1) 23
- BOLLOW M *see* Braun J
- BOLOGNA C, VIU P, JORGENSEN C, SANY J. Effect of age on the efficacy and tolerance of methotrexate in rheumatoid arthritis. 453
- BOLOGNA C *see also* Jorgensen C
- BOLTON M *see* Howat S
- BOLTON MC, DUDHIA J, BAYLISS MT. Quantitation of aggrecan and link protein mRNA in human articular cartilage of different ages by competitive RT-PCR. (A1) 237
- BOLTON MC *see also* McKenna LA
- BOMBARDIERI S *see* Ferri C
- BOND A *see* Dalziel M
- Bone densitometry, a funded service—audit of referrals over a 2 year period. (A2) 5
- Bone density
low in patients falling outside Compston guidelines for DEXA scanning. (A1) 83
lumbar spine measurements: the effect of posterior facet joint osteoarthritis in a middle age female population. (A1) 98
- Bone disease
duration effects in rheumatoid arthritis on appendicular bone mineral measurements. (A1) 86
metastatic, iliopsoas bursitis as unusual presentation. 285
- Bone erosion, prognostic value of contrast enhanced Gd-DTPA MRI for development in rheumatoid arthritis. (S3) 27
- Bone formation and degradation marker relationship in serum and synovial fluid in rheumatoid patients. (A1) 91
- Bone lesion, tumour-like, isonitrite-Tc99 bone scan vs radiologic evaluation in the assessment of the malignant nature. (A1) 241
- Bone loss
effect of cyclical didronel in early postmenopausal women. (A1) 87
trabecular and cortical bone loss in systemic lupus erythematosus. 244
- Bone markers, effect of corticosteroids on in rheumatoid arthritis. (A1) 120
- Bone mass
clinical associations of dual-energy X-ray absorptiometry measurement in rheumatoid arthritis. 1256

- effects of corticosteroids over 2 years when used in management of early rheumatoid arthritis, (A1) 210
- and polymorphism at the oestrogen receptor gene in pre- and post-menopausal women, (A1) 89
- Bone mass measurement**
- and bone metabolism in rheumatoid arthritis: a review, 309
- peripheral—is there any clinical value in rheumatoid arthritis?, (E) 109
- Bone metabolism**
- and bone mass measurement in rheumatoid arthritis: a review, 309
- marker changes in early rheumatoid arthritis, (A2) 3
- Bone mineral density**
- and bone turnover in treated coeliac disease, (A1) 85
- fluoride pharmacokinetics and changes in lumbar spine and femoral neck, (A1) 84
- fore-arm in non-steroid treated rheumatoid arthritis is reduced in patients on long-term methotrexate compared with other disease-modifying drugs, (A1) 209
- hand, cross-sectional area of forearm musculature and grip strength in RA, (A1) 205
- how much does position of the spine affect values?, (A1) 82
- Bone mineral measurements, appendicular, effects of bone disease duration in rheumatoid arthritis, (A1) 86**
- Bone mineral metabolism, effect of intramuscular corticosteroid on markers in rheumatoid patients, (A1) 81**
- Bone morphogenetic proteins, and osteoblastic bone nodules are induced bone growth stimulating pulsed electromagnetic fields, (A1) 115**
- Bone resorption, serum cross-linked telopeptides of type I collagen are a useful marker in the rat, (A1) 91**
- Bone turnover**
- biochemical markers in ankylosing spondylitis, (A1) 151
- and polymorphisms of the vitamin D receptor in female twins, (A1) 88
- suppression by aminohexane bisphosphonate in postmenopausal women, more rapid than oestrogen-gestagen therapy, 636
- BONK C see Dreses-Werlingloer U**
- BOONEN A, SCHREY G, VAN DER LINDEN Sj. Clubbing in human immunodeficiency virus infection, 292**
- BOONEN S see Dequeker J**
- BOOTH J see Snowden N**
- BOOTSMA H see Spronk PE**
- BORNHANN M see Mau W**
- BORYSIEWICZ LK see Lawson TM**
- BOTTO M see Davies KA; Naughton M**
- BOUMPAS DT. Novel action of glucocorticoids—NF- κ B inhibition, (E) 709**
- BOUVIER M see Maillefert JF**
- BOWMAN S see Hameed K**
- BOWMAN SJ, GEDDES GC, CORRIGALL V, PANAYI GS, LANCHBURY JS. Large granular lymphocyte expansions in Felty's syndrome have an unusual phenotype of activated CD45RA⁺ cells, 1252**
- BOWMAN SJ see also Allen RL**
- BOWNESS P, ALLEN RL, McMICHAEL AJ. HLA-B27-restricted peptide presentation to cytotoxic T cells in reactive arthritis, (A1) 114**
- BOWNESS P see also Allen RL**
- BPI-ANCA system, immunodiagnostic importance, (A1) 50**
- BRADLEY SM. Antibody levels to *Clostridium perfringens* and response to sulphasalazine, (L) 1327**
- BRADLOW A see Turner HE**
- BRAGANZA JM see Chaloner C**
- BRANCACCIO V see Ames PRJ**
- BRANCOLINI V see Hardwick L**
- BRASCH RC see Demars F**
- BRAUN J, TUSZEWSKI M, EHLERS S, HÄBERLE J, BOLLOW M, EGGENS U, SIEFER J. Use of multi-bacterial nested PCR to screen synovial fluid and membrane samples of patients with inflammatory arthritis and sacroiliitis, (A1) 113**
- BRAUN J see also Wilkinson NZ**
- BREEDVELD F see Paleolog E**
- BREEDVELD FC see de Vries E; Dothain RJEM; Ronday HK; Simon CH; Vliet Vlieland TPM**
- BREGEON C see Combe B**
- BRENNAN F see Jobanputra P**
- BRENNAN FM, WILLIAMS L, GIBBONS DL, MAINI RN, FELDMANN M. Paradoxical effects of a synthetic metalloproteinase inhibitor which blocks both p55 and p75 TNF receptor shedding and TNF α processing in RA synovial membrane cell cultures, (A1) 181**
- BRENNAN FM see also Malfait A-M**
- BRENNAN FR, MIKECZ K, PINDER S, JOBANPUTRA P, BAVINGTON C, LIN H, MORRISON P, GLANT TT, NUKI G. Anti-CD44 antibody inhibits human lymphocyte adhesion to endothelial cells and synoviocytes by inducing proteolytic cleavage of CD44, (A1) 60**
- BRENNAN FR see also Bavington C**
- BRENNAN P, SILMAN A. Are hormonal associations with rheumatoid arthritis due to HLA DR4, (A1) 1189**
- BRENNAN P, SILMAN A, HARRISON B, OLLIER B, JAWAHEER D, SYMMONS D. Role of the HLA-DRB1 genotype in the susceptibility to and severity of inflammatory polyarthritis, (A1) 186**
- BRENNAN P see also Harrison BJ; Symmons DPM**
- BRENNAN PJ see Bryan C**
- BRESNIHAN B see Bolger T; Costello P; Cunnane G; Mulherin D**
- BRICENO F see Sanchez J**
- BRIGGS D see Davies KA; Öcal L**
- BRIGGS DC see Artlett CM**
- BRILLANTE M see Ames PRJ**
- British Paediatric Rheumatology Group news, 686**
- BRITO M, CANTABRANA A, RAMOS P, SEGOVIA C, AMÉRIGO MJ, ROCES A. Multiple dysplasia in ochronosis: a non-rheumatological series of 16 patients, (A1) 95**
- BRITO M see also Cantabrana A**
- BROAD I see Etherington J**
- BROCK JH see Houston T**
- Bromazepam and tenoxicam comparison in the treatment of fibromyalgia. A randomized, double-blind, placebo-controlled trial, (A1) 138**
- Bronchiectasis and rheumatoid arthritis, defective antibody production, (A1) 201**
- Bronchoalveolar lavage in mixed cryoglobulinaemia associated with hepatitis C virus, 978**
- BROOKES SM see Price EJ**
- BROOKS PM. Undergraduate education in rheumatology, (E) 203**
- BROOKS PM see also Jones G**
- BROSTOFF J see Kjeldsen-Kragh J**
- BROUSSE S see Lioté F**
- BROWN MA, JEPSON A, YOUNG A, GREENWOOD B, WORDSWORTH BP. Spondylarthritis in the Gambia—evidence for a non-HLA B27 protective effect, (A1) 113**
- BROWN RA see Ahmed Z**
- BRUCE IN see Smyth AE**
- BRUCKNER FE see Laversuch CJ**
- BRÜGGEMAN M see Harmer IJ**
- BRUNZEEEL CAFM see Verhoef CM**
- BRUS HLM, VAN DE LAAR MAFJ, TAAL E, RASKER JJ, WIEGMAN O. Influence of formal education on compliance and disease activity in early RA patients starting sulphasalazine, (A1) 216**
- BRIJN GAW, GRIEF EN. Eruption of vasculitis during treatment with leflunomide for rheumatoid arthritis, (A1) 226**
- BRYAN C, BRENNAN PJ, HOWARD Y, SILMAN AJ, BLACK CM. Predictors of mortality in a cohort of patients with newly diagnosed scleroderma, (A1) 122**
- BRYAN C, HOWARD Y, BRENNAN P, BLACK C, SILMAN A. Survival following the onset of scleroderma: results from a retrospective inception cohort study of the UK patient population, 1122**
- BRYDSON G see Hamilton J**
- BUCHANAN N see Etherington J**
- BUCHANAN NMM see Khamashta MA; Ruiz- Irastorza G**
- BUCHNER E see Burkhardt H**
- BUCKLAND-WRIGHT C see Sharma S**
- BUCKLAND-WRIGHT JC, LYNCH JA, BIRD C. Microfocal techniques in quantitative radiography: measurement of cancellous bone organization, (S3) 19**
- BUCKNALL RC see MacDonald AG; Over KE**
- BUKHARI M, HERRICK A, MANNING J, MOORE T, JAYSON MIV. Digital and microvascular blood flow responses in patients with systemic sclerosis and primary Raynaud's phenomenon, (A1) 42**
- BUKHARI M, HERRICK AL, MOORE T, MANNING J, JAYSON MIV. Increased nailfold capillary dimensions in primary Raynaud's phenomenon and systemic sclerosis, 1127**
- BULPITT KJ see Isaacs JD**
- BUNN C see Stratton R**
- BURD GP, MAINI RN, MAGEED RA. Different patterns of cytokine and CD40-ligand expression in spleen germinal centres of NZB/W F1 mice compared with normal CBA mice, (A1) 5**
- BURGE PD see Chitnavi: J**
- BURKHARDT H, MEYER P, BUCHNER E, PALOMBO-KINNE E, KALDEN RW. Serine proteinase inhibitor antileukoproteinase specifically accumulates in normal but not in arthritic cartilage, (A1) 68**
- BURMEISTER G see Wedekind F**
- BURMEISTER GR see Riemekasten G**
- BURNS A see Stratton R**
- BURT J see Macfarlane GJ**
- BUSCH U see Türk D**
- BUSCOMBE J see MacGregor AJ**
- BUSKILA D see Ehrenfeld M**
- BUSTABAD S see Hdez-Berian JA**
- BUTCHER S see Hardwick L; Hardwick LJ**
- BUTLER DM see Malfait A-M**
- BUTLER RC see Martin JC**
- BÜTZOW C see Kohlman T**

- BYNG P *see* Coggon D
- BYRNE PAC, EVANS WD, RAJAN KT. How much does position of the spine affect bone mineral density values? (A1) 82
- BYRNE PAC, REES JIS, WILLIAMS BD. Iliopsoas bursitis, an unusual presentation of metastatic bone disease, 285
- BYRNE PAC. Treatment of gout following cardiac transplantation, (L) 1329
- CIQ deficiency, complete, molecular basis in three patients, (A1) 27
- Clq(CLR) antibodies are markers for severe disease in systemic lupus erythematosus, (A1) 21
- C-reactive protein
is elevated in osteoarthritis, (A1) 99
low level increases are present in early osteoarthritis of the knee and predict progressive disease, (A1) 99
- CABEZAS JA *see* Uson J
- CABO X *see* Valverde J
- CABRAL DA *see* Foster HE
- CABRÉ E *see* Olivé A
- Cachexia causes in rheumatoid arthritis, (A1) 200
- CAEIRO F *see* Hardwick L
- CALABOZO M, ALONSO-RUIZ A, PEREZ-RUIZ F, SAEZ F, RUIZ-LUCEA E, ARETXABALA I, FRAIX E. Evaluation of the efficacy of radiosynovectomy of the knee joint with gadolinium-enhanced magnetic nuclear resonance, (A1) 224
- CALABOZO RALUY M *see* Perez-Ruiz F
- Calcific periarthritis involving multiple sites. A familial condition, (A1) 133
- Calcitropic hormones and osteocalcin levels in lung transplant recipients, (A1) 86
- Calcium and phosphate supplements timed therapy restores phosphate and parathyroid hormone rhythms in patients with osteoporosis, with an anabolic effect on spinal bone mineral density, (A1) 88
- Calcium pyrophosphate dihydrate (CPPD) crystal arthropathy and gout, diagnosis, (E) 306
- Calcium release, intracellular in T-lymphocytes from patients with rheumatoid arthritis and ankylosing spondylitis, (A1) 11
- CALIN A *see* Gomez K; Haigh RC; Jones SD; Kabasakal Y; Kennedy LG; Koh WH; Mack CS; Roussou E; Taylor AL
- CALLARD P *see* Guillemin L
- Calman Report, consequences on academic rheumatology, (E) 608
- CALVER D *see* Mishra MB
- CALVO-ALÉN J, CORRALES A, FERNÁNDEZ-ECHEVARRÍA MA, SANCHEZ-ANDRADA S, PÉRA JL, RODRÍGUEZ-VALVERDE V. Reliability of simple parameters in the assessment of clinical activity in RA, (A1) 232
- CALVO-ALÉN J *see also* Sánchez-Andrada S
- CAMACHO M, ANTÓN R, GARCÍA M, VILA L. Linoleic acid is a better substrate for PGH-synthase-2 than for PGH-synthase-1 tested in whole endothelial cells in culture, (A1) 219
- CAMBRIDGE G *see* Leigh RD
- CAMILLERI J *see* Hall FC
- CAMILLERI JP *see* Williams AS
- CAMPATH-1H in rheumatoid arthritis—an intravenous dose-ranging study, 231
- CAMPBELL F, BIRKELUND S, WARD ME, PANAYI GS, KINGSLEY GH. Synovial fluid derived T cell clones respond to *Chlamydia trachomatis* protein antigens, (A1) 114
- CAMPBELL L *see* Coggon D
- CAMPBELL R *see* Speed CA
- CAMPION G *see* Jevtic V
- CAMPOY E *see* Narváez J
- Cancellous bone, measurement of organization, microfocal techniques in quantitative radiography, (S3) 19
- Cancer, does exposure to immunosuppressive therapy increase the 10 year malignancy and mortality risks in rheumatoid arthritis? A matched cohort study, 738
- CANTABRANA A, BRITO M, SEGOVIA C, RAMOS P, ANTOLIN J, ROCES A. Ochronotic arthropathy: a non-rheumatological series of 16 patients, (A1) 95
- CANTABRANA A *see also* Brito M
- CAPELL H *see* England AJ
- CAPELL HA, MORRISON E, DUNCAN MR, McENTEGART A, PORTER DR, MADHOK R, THOMSON EA. Are we serving female patients less well than males? Baseline characteristics of rheumatoid arthritis patients enrolled into sulphasalazine studies, (A1) 213
- CAPELL HA, MUNRO R. Weight loss and body composition in rheumatoid arthritis, (A1) 200
- CAPELL HA *see also* Duncan MR; McEntegart A; Morris AJ; Morrison E; Munro R; Porter DR
- CARACUEL MA *see* González J
- CARBON RJ, MACEY MG, MCCARTHY DA, PEREIRA FP, PERRY JD, WADE AJ. Effect of 30 min cycle ergometry on ankylosing spondylitis, 167
- Cardiac abnormalities in a patient with localized scleroderma, (L) 703
- Cardiac function, echocardiographic assessment in Paget's disease of bone, (A1) 93
- Cardiac transplantation, treatment of gout following, (L) 1329
- Care path, developing for low back pain and sciatica: from primary through secondary to tertiary care, (A1) 143
- CARLI PM *see* Maillefert JF
- CARLSON MR *see* Lanza FL
- Carotid artery stenosis, CD8+ T-cell reduced concentration in peripheral blood: relation to arteritis temporalis, 649
- CAROTTI M *see* Manganelli P
- Carpal tunnel syndrome, idiopathic, management: survey of rheumatologists, (A2) 34
- CARP A *see* Hardwick L
- CARR AJ. Patient-centred approach to evaluation and treatment in rheumatoid arthritis: the development of a clinical tool to measure patient-perceived handicap, 921
- CARR AJ, THOMPSON PW, KIRWAN JR. Quality of life measures, 275
- CARR AJ, WOOLHEAD GM, WILLIAMS MH, BANNISTER G, ACKROYD C, THOMPSON PW. Patterns of symptom impact in acute and chronic knee pain, (A1) 133
- Psychological impact of knee pain, (A1) 135
- CARR AJ, WOOLHEAD GM, WILLIAMS MH, WILKINSON M. Impact of a common symptom: validation of the disease repercussion profile for knee pain, (A1) 133
- CARR AJ *see also* Chitnavis J; Harwood RH
- CARRERO L, LÓPEZ-LONGO FJ, GONZÁLEZ CM, LAMPREAVE JL, GONZÁLEZ M, VEGAZO O, MINGUEZ MD, FERNÁNDEZ-ESPARTERO MC, ALMOGUERA MI, MONTEAGUDO I. Influence of non-steroidal antiinflammatory drugs on the evolution of abdominal scintigraphy in patients with spondylarthropathies, (A1) 154
- CARRERO L *see also* Abad MA; González M; López-Longo FJ; Monteagudo I
- CARRUTHERS D *see* Exley A
- CARRUTHERS DM, WATTS RA, SYMMONS DPM, SCOTT DGI. Wegener's granulomatosis, increasing incidence or increased recognition?, 142
- CARRUTHERS DM, YOUNG SP, BACON PA. Intracellular calcium release in T-lymphocytes from patients with rheumatoid arthritis and ankylosing spondylitis, (A1) 11
- CARTER M *see* Naughton M
- Cartilage
actual repair by Ilizarov joint distraction tested in the Pond-Nuki dog model for osteoarthritis, (A1) 104
- aggregrin domains, production as recombinant fusion proteins to analyse recognition by autoreactive T lymphocytes, (A1) 184
- articular, sustained loading makes it stronger, (A1) 240
- collagen destruction, oncostatin M is raised in rheumatoid synovial fluids and interacts with IL-1, (A1) 109
- osteoarthritic, increased expression of the Ed-B-containing fibronectin (an embryonic isoform of fibronectin), 407
- and perichondrium implanted in the joint, (A1) 240
- sustained loading reduces the stress-equalising property, (A1) 239 and synovium
- MRI in the assessment, (S3) 3
- pathophysiology, (S3) 11
- CARTY H *see* Lees C
- CARVALHO MD *see* Denton CP
- CASADEI MALDINI M *see* Boiardi L
- CASADEMONT J *see* Font C
- CASAN P *see* Giral P
- CASASSUS P *see* Guillemin L
- CASATTA L *see* Ferraccioli GF
- CASEY ATH, YOUNG A, CROCKARD HA, BLAND M. Myelopathy functional disability index—a step towards objectivity, (A2) 19
- CASEY E *see* Guerin J
- CASEY EB *see* Donnelly S; McGonagle D; Montague S; Ormond P
- CASSISI GA *see* Fiocco U
- CASTELL M *see* Ros S
- Catalase, increased activity in synovial fluid of rheumatoid arthritis patients compared to osteoarthritis patients, (A1) 70
- Catecholamines, reduced influence on proliferation of lymphocytes from patients with rheumatoid arthritis *in vitro*, (A1) 65
- CATRAVAS J *see* Walsh DA
- CAUGHEY D *see* Rudge SR
- CAULI A *see* Iannone F; Yanni G
- Cause of death
at autopsy in 22 patients with rheumatoid arthritis, (A2) 22
- in two hundred patients with systemic lupus erythematosus, (A1) 37
- CAVANAGH G *see* Davidson BKS
- CRAWLEY MID *see* Smith RW; Wong WM
- CRAWSTON TE *see* Clark IM; Ellis AJ; Summers CA
- CAWTE SA *see* Yeap SS
- CAZABON JK *see* Loizou S
- CD1 expression in skin and synovial membrane of rheumatoid arthritis and psoriatic arthritis patients, (A1) 179
- CD8+ T cells
characterisation in psoriatic arthritis synovial effusions, (A1) 160

- reduced concentration in peripheral blood of patients with carotid artery stenosis: relation to arteritis temporalis, 649
and their T cell receptor repertoires in rheumatoid arthritis, (A1) 183
- CD8 lymphocyte subsets in active polymyalgia rheumatica: comparison with elderly-onset and adult rheumatoid arthritis and influence of prednisone therapy, 642
- CD40 primary impaired pathway, hyper-IgM syndrome associated with rheumatoid arthritis, 282
- CD40-ligand and cytokine expression, different patterns in spleen germinal centres of NZB/W F1 mice compared with normal CBA mice, (A1) 5
- CD44, proteolytic cleavage induction, anti-CD44 antibody inhibits human lymphocyte adhesion to endothelial cells and synoviocytes, (A1) 60
- CD45, large granular lymphocyte expansions in Felty's syndrome have an unusual phenotype of activated CD45RA⁺ cells, 1252
- CD69 on synovial T cells in rheumatoid arthritis correlates with disease activity, (L) 397
- CD95/FAS/APO-1, murine, histological analysis, (A1) 39
- CEDRASCHI C see Weber M
- Cellular immunity and endothelial responses, contemporaneous activation in limited cutaneous systemic sclerosis, (A2) 27
- Central nervous system side effects with low dose methotrexate and mycristin therapies in rheumatoid arthritis, (A1) 216
- CERAVOLO GP see Marchesoni A
- CEREJO MJ see González-Gay MA
- CERF I see Lioté F
- CERVERA R see García-Carrasco M; Navarro M
- Cervical myelopathy, unusual complications of cervical spine surgery in rheumatoid arthritis, 682
- Cervical narrow disc spaces, importance in patients with rheumatoid arthritis, (A1) 206
- Cervical spine surgery for cervical myelopathy, unusual complications in rheumatoid arthritis, 682
- Cervical spondylodiscitis in a patient with ankylosing spondylitis, (L) 1034
- CERVINI C see Manganello P
- CHABBOU A see Hamzaoui A
- CHACKFIELD M see Mills SY
- CHAKRAVARTY K. Phenyltoin-induced pseudo-Sjögren's syndrome, (L) 1033
- CHAKRAVARTY K see also Symmons DPM
- Chlamydia* reactive arthritis
sexually acquired, synovial fluid and serum antibodies against *Chlamydia* in different forms of arthritis: intra-articular IgA production, 548
value of isotype determination of serum antibodies against *Chlamydia* for diagnosis, 542
- CHALONER C, RIELEY F, BLANN AB, HERRICK A, BRANGANZA JM. Contemporaneous activation of endothelial responses and cellular immunity in limited cutaneous systemic sclerosis, (A2) 27
- CHAMBERLAIN MA, ROONEY CM. Young adults with arthritis: meeting their transitional needs, 84
- CHAMBERLAIN MA see also Tennant A
- CHAMBERS JB see Mishra MB
- CHAMBERS TJ see Gallagher AC; Tobias JH
- CHAN D see Mok CC
- CHAN E see Mok CC
- CHAN JK-M see Patel S
- CHAN SY see Mok CC
- CHANDLER BJ see Kent RM
- CHANDLER C see Chatterjee S; Kay LJ; Walker DJ
- CHAPMAN PT, JAMAR F, HARRISON AA, SCHOFIELD JB, PETERS AM, BINNS RM, HASKARD DO. Characterization of E-selectin expression, leucocyte traffic and clinical sequelae in urate crystal-induced inflammation: an insight into gout, 323
- CHAPMAN PT, JAMAR F, PETERS AM, HASKARD DO. Imaging inflammation in rheumatoid arthritis: comparisons between ¹¹¹In-anti-E-selectin MAB 1.2B6 and ^{99m}Tc- and ¹¹¹In-immunoglobulin, (A1) 203
- CHAPMAN PT, YARWOOD H, HARRISON AA, STOCKER CJ, JAMAR F, GUNDEL RH, PETERS AM, HASKARD DO. Endothelial activation in urate crystal-induced inflammation: *in vitro* and *in vivo* studies on the role of tumour necrosis factor, (A1) 146
- CHARLES P see Davis DM
- CHARLIER C see Maugars Y
- CHARLWOOD C see Black AJ
- CHASTANG C see Ravaud P
- CHATARD C see Maillefert JF
- CHATELIER G see Cordonnier C
- CHATTERJEE S, LINGARD E, PLUMMER CJ, CHANDLER C, HESLOP PS, WALKER DJ. How well do 10m walking speed and treadmill tests reflect 24 hour activity?, (A1) 194
- Chemotherapy, intensive for AL amyloidosis, (A1) 12
- CHERNAJOVSKY Y, ADAMS G, NEVE R, HERMAN C, TRIANTAPHYLLOPOULOS K. Use of lymphoid cells capable of transferring disease as carriers of immunomodulators—a rheumatoid arthritis model, (A1) 182
- CHERNAJOVSKY Y see also Gould DJ; Herman C; Kissnerghis M
- CHERREZ C see Hdez-Berriain JA
- Chest pain, atypical of musculoskeletal origin, (A1) 136
- CHEUNG NT, TAYLOR DJ, CLARKE S, FISHER J, JONES PW, MATTEY DL, DAWES PT. Effect of 6 months treatment on serum pro-stromelysin and pro-collagenase levels in rheumatoid arthritis patients, (A1) 120
- CHEUNG NT see also Close DR
- CHEVALIER X, GROULT N, EMOD I, PLANCHENAUT T. Proteoglycan-degrading activity associated with the 40 kDa collagen-binding fragment of fibronectin, 506
- CHEVALIER X, GROULT N, HORNEBECK W. Increased expression of the Ed-B-containing fibronectin (an embryonic isoform of fibronectin) in human osteoarthritic cartilage, 407
- CHEVALIER X, TYLER JA. Production of binding proteins and role of the insulin-like growth factor I binding protein 3 in human articular cartilage explants, 515
- CHIECO-BIANCHI F see Fiocco U
- CHIKANZA I see Jawed S
- CHIKANZA IC see Kassimos D
- CHILD A see Mishra MB
- Childhood infections, do they affect the development of ankylosing spondylitis, (A2) 23
- Children
back pain, 1022
pattern of joint involvement in Lyme arthritis, 1016
- CHITNAVIS J, CLIPSHAM K, BURGE PD, CARR AJ. Familial tendency to severe symptomatic osteoarthritis of the hip and knee, (A1) 100
- Chlamydia*, synovial fluid and serum antibodies against in different forms of arthritis: intra-articular IgA production in *Chlamydia* sexually acquired reactive arthritis, 548
- Chlamydia pneumoniae*
and *Chlamydia trachomatis* detection in the synovium of patients with a range of rheumatic diseases, (A1) 162
reactive arthritis and persistent infection, (L) 1189
- Chlamydia trachomatis*
effect of ciprofloxacin and interferon-gamma on infection in the monocytic cell line THP-1, (A1) 169
synovial fluid derived T cell clones respond to protein antigens, (A1) 114
- Cholesterol, serum and risk of rheumatoid arthritis in a cohort of 52 800 men and women, 255
- Chondrocalcinosis
familial, synovial chondromatosis prevalence, (A1) 147
genetic linkage to chromosome 5p15.1-15.2, (A1) 148
- Chondrocytes
human, accelerated proteoglycan synthesis following cyclical pressurisation is mediated by $\beta 1$ integrin and stretch activated ion channels, (A1) 110
and polymorphonuclear leucocytes, study of cooperation in leukotriene biosynthesis, (A1) 62
- Chondroitin and keratan sulphate markers in normal and arthritic synovial fluid, 951
- Chondroitin sulphate isomers and proteoglycan markers in synovial fluids from healthy volunteers and patients with arthritis, (A1) 239
- CHOW CC see Szeto CC
- CHOW JWM see Gallagher AC
- CHOWDHRY KM see Roussou E
- CHOWDREY HS see Harbuz MS
- CHOY EHS, RANKIN ECC, KASSIMOS D, VETTERLEIN O, SOPWITH M, EASTELL R, ISENBERG DA, KINGSLEY GH, PANAYI GS. Engineered human anti-tumour necrosis factor alpha antibody reduces serum interleukin-6 and urine bone markers but has no effect on soluble CD4 and soluble interleukin-2 receptor in rheumatoid arthritis, (A1) 214
- CHOY EHS see also Houssien DA; Kassimos D
- Chromosomal instability, telomere reduction in scleroderma patients, 732
- Chronic fatigue syndrome, seronegative Sjögren's syndrome manifested as a subset, 471
- CHU C-Q see Kang RY
- CHU CQ, LONDEI M
Inhibition of murine collagen-induced arthritis by intranasal administration of collagen type II peptides, (A1) 73
Prevention of collagen-induced arthritis by a non-depleting anti-CD4 antibody, (A1) 108
- CHU CQ see also Taylor PC
- Churg-Strauss syndrome
antineutrophil cytoplasmic antibodies, abnormal angiograms and pathological findings, 958
in a district general hospital, (A1) 53
thrombosis in. Beyond vasculitis, 1181
- CIACCI C see Lubrano E
- CICONELLI RM, FERRAZ MB, VISIONI RA, OLIVEIRA LM, ATRA E. Randomized double-blind controlled trial of sulphasalazine combined with pulses of methylprednisolone or placebo in the treatment of rheumatoid arthritis, 150
- CICUTTINI F see Cooper C; Spector TD

- CID MC *see* Font C
- CIMAZ R. Methotrexate treatment every other week in patients with juvenile chronic arthritis, (L) 1030
- CIMINO R *see* Ames PRJ
- Ciprofloxacin and interferon-gamma, effect on the infection of *Chlamydia trachomatis* in the monocytic cell line THP-1, (A1) 169
- CISNAL A, COLLANTES E, MUÑOZ E, ACASUSO M, Spanish Spond. Study Group. Assessment of performance of ESSG criteria according to diagnostic age, (A1) 150
- CISNAL A, MUÑOZ E, COLLANTES E, HAJJAL-HASSOUNI N, DOUGADOS M, AMOR B. Spondylarthropathies, their distribution by diagnostic criteria in three countries, (A1) 150
- CISNAL A *see also* Collantes E
- CLAASSEN E *see* Dolhain RJEM
- CLARK B *see* Foley-Nolan D; Griffiths B
- CLARK IM, ROWAN AD, EDWARDS DR, CAWSTON TE. Mapping the human TIMP-1 gene promoter, (A1) 68
- CLARK IM *see also* Summers CA
- CLARKE S *see* Cheung NT
- CLAVAGUERA MT *see* Valverde J
- CLAVAGUERA T *see* Narváez J
- CLEARY R *see* Walker DJ
- CLELAND LG. Animal models of rheumatoid arthritis, (E) 1041
- CLEMENTS PJ *see* Seibold JR
- Clinic attenders, comparison of disease status in rheumatoid arthritis patients attending and not attending a specialist clinic, 1169
- Clinical activity, reliability of simple parameters in the assessment in RA, (A1) 232
- Clinical audit. Isotope bone scans for foot problems, (A2) 8
- Clinical and radiological progression, differences in rheumatoid arthritis, (A1) 199
- CLINTON-WATKINS A *see* Adams MA
- CLIPSHAM K *see* Chitnavis J
- Clodronate, effects on disease activity and bone mineral density in rheumatoid arthritis, (A2) 21
- CLOSE DR, CHEUNG NT, EVANS EL, CLOSE JA, DAWES PT. Evidence that silica glass fragments from steroid containing ampoules augment the intra-articular inflammatory response, (A1) 121
- CLOSE JA *see* Close DR
- Clostridium perfringens* antibody levels and response to sulphasalazine, (L) 1327
- CLOTET B *see* Olivé A
- Clubbing in human immunodeficiency virus infection, 292
- COAKLEY G, LANCHBURY JS. MHC-encoded TAP genes in rheumatic diseases, (E) 501
- Coagulation factor V gene mutation increases the risk of venous thrombosis in Behçet's disease, 1178
- COCKRAM CS *see* Szeto CC
- Coeliac disease
- arthritis: prevalence and pattern in 200 adult patients, 1314
 - bone mineral density and bone turnover, (A1) 85
 - co-existent with scleroderma, (L) 807
- Cogan's syndrome, atypical, associated with antineutrophil cytoplasmic autoantibodies, (L) 601
- COGGON D, EGGER P, CAMPBELL L, BYNG P, CROFT P, McLAREN M, COOPER C. Postmenopausal oestrogen use and osteoarthritis of the hip, (A1) 101
- COGGON D, EGGER P, CAMPBELL L, CROFT P, McLAREN M, COOPER C. Occupational lifting and osteoarthritis of the hip, (A1) 100
- COGGON D *see also* Cooper C; Smedley J
- COHEN MR, ISENBERG DA. Ultraviolet irradiation in systemic lupus erythematosus: friend or foe?, 1002
- COHEN SBA *see* Price EJ
- COHEN TERVAERT JW *see* Spronk PE
- COLAO A *see* Scarpa R
- Colitis, ischaemic, polyarthritis nodosa presenting as, (L) 1334
- COLL-VINENT B *see* Font C
- Collagen
- interleukin-4 can prevent release from cartilage stimulated with proinflammatory cytokines *in vitro*, (A1) 8
 - type I mRNA expression, atypical synovial cells are involved in rheumatoid arthritis, (A1) 178
 - type I, serum cross-linked telopeptides are a useful marker of bone resorption in the rat, (A1) 91
 - type II, induction of chronic autoimmune arthritis in DBA/1 mice by oral administration, 828
 - type II peptides, inhibition of murine collagen-induced arthritis by intranasal administration, (A1) 73
 - type II-induced arthritis transfer from DBA/1 to SCID mice can be prevented by blockade of MAC-1, (A1) 74
- Collagenase
- responsible for the destruction of the collagen network in human cartilage, (E) 818
- and stromelysin, effect of low dose steroid treatment on levels in RA patients followed for 3 years, (A1) 210
- COLLANTES E, LÓPEZ A, CISNAL A, MONTILLA C, ROLDAN R, LÓPEZ E. Stereological analysis of the synovial membrane in rheumatic disorders, (A1) 177
- COLLANTES E, TINAHONES F, CISNAL A, PÉREZ G, SORIGUER F, SANCHEZ P. Changes in apolipoproteins levels following low-calorie dieting in hyperuricemic patients, (A1) 145
- COLLANTES E *see also* Cisnal A
- COLLINS I *see* Ng T
- COLLUM L *see* Mullaney P
- COLVIN E *see* Sharif M
- COLVIN EM *see* Sharif M
- COMBE B, GOUPILLE P, KUNTZ JL, TEBIB J, LIOTÉS F, BREGEON C. Sulphasalazine in psoriatic arthritis: a randomized, multicentre, placebo-controlled study, 664
- Combination therapy with sulphasalazine and azathioprine, (L) 493
- COMER M *see* Houssien DA
- Community clinic, comparison with a hospital out-patient clinic in rheumatology, 385
- Complement C3, extrahepatic secreted contributes to circulating C3 levels in humans, (A1) 66
- Complement C4 null alleles and HLA-DRB1 alleles in systemic lupus erythematosus in a Spanish population, (A1) 26
- Complement, immune complexes and systemic lupus erythematosus, 5
- Complement receptor one (SCR1), administration of local and systemic has a beneficial effect on the development of rat mono-articular arthritis, (A1) 74
- Compliance monitoring of NSAID drug therapy in ankylosing spondylitis, experiences with an electronic monitoring device, 60
- Compstone DEXA scanning guidelines
- low bone density in patients falling outside, (A1) 83
 - may underestimate community needs, (A1) 84
- Computerized tomography, high resolution and sequential DTPA clearance in monitoring interstitial lung disease in dermatomyositis, 164
- CONCA W *see* Machein U
- Connective tissue disease
- joint involvement and its associations with the third hypervariable region (HVR3), (A1) 30
 - prevalence of underlying in patients with suspected primary Raynaud's phenomenon, (A2) 26
- CONROZIER T, JOUSSEAUME C, MATHIEU P, PIPERNO M, VIGNON E. Factors affecting the outcome of hip osteoarthritis, (A1) 102
- CONSIDINE E *see* Hardwick L
- CONSIGLI G *see* Manganelli P
- CONTI L *see* Ferri C
- COOKE N *see* Walne AJ
- COOKE S *see* Rigby S
- COOKE SP, FORREST GR, MITCHELL TJ, VENABLES PJW. Human endogenous retrovirus-3 a highly conserved immunoregulatory sequence?, (A1) 17
- COOKSON J *see* Gaffney K
- COOMBS EJ *see* Sinclair D
- COOMBS G, MCCLOSKEY E, BERNARD J, REANEY L, KANIS J. Morphometric X-ray absorptiometry and radiographic vertebral morphometry in established postmenopausal osteoporosis, (A1) 83
- COOMBS GM, TEH LS, DENTON J, JOHNSON AS, JONES AKP. *Mycobacterium xenopi*—an unusual presentation as tenosynovitis of the wrist in an immunocompetent patient, 1008
- COOPER AM, SHARIF M, EASTELL R, KIRWAN JR. Effect of corticosteroids on bone markers in rheumatoid arthritis, (A1) 120
- COOPER C, EGGER P, COGGON D, HART D, MASUD T, CICUTTINI F, DOYLE D, SPECTOR T. Generalised osteoarthritis in women: pattern of joint involvement and approach to definition, (A1) 97
- COOPER C *see also* Coggon D; Nahit ES; O'Neill TW; Reid DM; Smedley J
- COOPER RG *see* Mannion AF
- COPE A *see* Davis DM
- Coping strategy influence on outcome in systemic lupus erythematosus, (A1) 33
- CORDONNIER C, MEYER O, PALAZZO E, DE BANDT M, ELIAS A, NICAISE P, HAIM T, KAHN MF, CHATELLIER G. Diagnostic value of anti-RA33 antibody, antikeratin antibody, antiperinuclear factor and antinuclear antibody in early rheumatoid arthritis: comparison with rheumatoid factor, 620
- CORRALES A *see* Calvo-Alén J; Sánchez-Andrada S
- Correlates of disablement in polyarticular juvenile chronic arthritis—a cross-sectional study, 91
- CORRIGALL V *see* Bowman SJ; Dolan AL; Iannone F
- Corticosteroid
- assessment of the efficacy of sacroiliac injections in spondylarthropathies: a double-blind study, 767
 - effects on bone markers in rheumatoid arthritis, (A1) 120
 - effects on bone mass over 2 years when used in management of early rheumatoid arthritis, (A1) 210

- intramuscular, effect on markers of bone mineral metabolism in rheumatoid patients, (A1) 81
- low-dose and blindness in giant cell arteritis, (L) 702
- in management of rheumatoid arthritis, (E) 2
- mediated change in lymphocyte adhesion to mucosal endothelium by mononuclear cells from rheumatoid arthritis patients, is it due to a quantitative change in lymphocyte subsets?, (A1) 14
- multiple joint injections in polyarticular juvenile chronic arthritis: a single-blind one year follow-up study, (A1) 121
- Corticotrophin-releasing factor binding protein, elevated levels in the blood of patients suffering from arthritis and septicemia and the presence of novel ligands in synovial fluid, 120
- Corticotrophin-releasing hormone gene, three novel polymorphisms in the regulatory region in patients with rheumatoid arthritis and controls, (A1) 107
- Cost
- of rheumatoid arthritis, 781
 - to society of treatment of rheumatoid arthritis, 791
- COSTELLO P, O'FARRELLY C, BRESNIHAN B, FITZGERALD O. Characterisation of CD8+ T cells in psoriatic arthritis synovial effusions, (A1) 160
- Costimulatory molecules B7.1 (CD80) and B7.2 (CD86), differential expression in rheumatoid synovial tissue, 33
- COUMBE A *see* Blake DR; Gaffney K
- COUPLAND CA *see* Patel S
- COX N *see* Prouse P; Talamo J
- COX NL, MISRA A, YOUNG KA. Simple index of range of movement of large joints as an outcome measure in RA, (A1) 194
- COX-2 inhibitors, and non-acidic pro-NSAIDs do not uncouple mitochondrial oxidative phosphorylation, (A1) 222
- COYLE M *see* Bolger T
- COYLE PV *see* Kane D
- COZZI L *see* Fiocco U
- CPPD deposition disease, familial, spondyloepiphyseal dysplasia associated with: study of a pedigree, (A1) 147
- CRAWFORD A *see* Foey A
- CREAMER P, KEEN M, ZANANIRI F, WATERTON J, MACIEWICZ R, DIEPPE P, WATT I. MRI can differentiate effects of intra-articular therapy on synovial proliferation and acute inflammatory changes in the knee joint, (A1) 205
- CREEMERS MCW, VANT HOF MA, FRANSSEN MJAM, VAN DE PUTTE LBA, GRIBNAU FWJ, VAN RIEL PLCM. Disease activity in ankylosing spondylitis: selection of a core set of variables and a first step in the development of a disease activity score, 867
- CREMER MA *see* Terato K
- CRILLY A, FIELD M, STURROCK RD, MADHOK R. Determination of cytokine profiles in RA synovial tissue using reverse transcriptase PCR, (A1) 6
- CROCKARD HA *see* Casey ATH
- CROFT P *see* Coggon D
- CROFT PR *see* Macfarlane GJ; Nahit ES; Pope DP; Pope DR
- CROWLEY B *see* Kane D
- CRUICKSHANK K *see* Öcal L
- CRUWYS SC, GARRETT NE, JOLLIFFE VA, KIDD BL. Synergistic interaction between morphine and indomethacin attenuates pain sensation and tissue destruction in adjuvant arthritis, (A1) 75
- CRUWYS SC *see also* Morris VH
- Cryofibrinogenemia, primary, familial presence, report of three cases, (L) 102
- Cryoglobulinaemia, mixed and hepatitis C virus genotypes, (A1) 50
- Cryptococcal brain abscesses in systemic lupus erythematosus, (L) 1334
- CERNOK E *see* Johnston TW; Stoffel MP
- CUERVAS-MONS V *see* Diez E
- CUNNANE G, BRESNIHAN B, FITZGERALD O. Immunohistologic analysis of peripheral joint disease in ankylosing spondylitis, (A1) 154
- CUNNINGHAM JL *see* Speed CA
- CUNNINGHAM P *see* Tiwana H
- CUNNINGHAM T *see* Bas S
- CURRAN MD *see* Kane D
- CURREY J. Scalp necrosis in giant cell arteritis, (A2) 36
- CURRY VA *see* Ellis AJ
- CURSON D. Patient education: treatment or nice extra. The British experience, (L) 805
- CUTOLO M *see* Ferraccioli GF
- CWYFAN-HUGHES S *see* Fernihough JK
- Cycle ergometry, 30 min, effect on ankylosing spondylitis, 167
- Cyclooxygenase, inducible, selective inhibition by meloxicam: summary of pharmacological data and clinical pharmacology, (A1) 221
- Cyclooxygenase-2
- expression in human and an animal model of rheumatoid arthritis, 711
 - inhibitors—and renal damage, (L) 1332
- Cyclophosphamide
- and prednisolone, intermittent pulse for the treatment of interstitial lung disease in systemic sclerosis, (A2) 27
 - pulse in the treatment of neuropsychiatric systemic lupus erythematosus, (A1) 32
- Cyclosporin
- combination therapy in rheumatoid arthritis, (S2) 19
 - is it a disease-controlling antirheumatic therapy? review of the two-year results of the GRISAR study, (A1) 215
 - potential role, new perspectives on controlling progression of severe rheumatoid arthritis, (S2) 1
- Cyclosporin A
- adding to parenteral gold therapy in rheumatoid arthritis: a double-blind placebo-controlled study, 1142
 - in the long-term treatment of psoriatic arthritis, (L) 1332
 - in the treatment of adult Still's disease, (A1) 77
 - in the treatment of systemic lupus erythematosus: results of an open clinical study, 669
- Cytokine
- and CD40-ligand expression, different patterns in spleen germinal centres of NZB/W F1 mice compared with normal CBA mice, (A1) 5
 - inflammatory, expression of MT-MMP1 by human fibroblast-like synoviocytes and its regulation, (A1) 67
 - inhibition of the Na/K-ATPase exerts differential effects on production by monocytes and fibroblasts, (A1) 67
 - inhibitor levels, changes in naturally occurring after TNF α blockade in rheumatoid arthritis, (A1) 181
 - pro-inflammatory expression in early collagen-induced arthritis, (A1) 6
 - profile determination in RA synovial tissue using reverse transcriptase PCR, (A1) 6
 - secretion pattern during the development of collagen induced arthritis, (A1) 5
 - Th1 and Th2 in rheumatoid arthritis, (A1) 179
- Cytotoxin, synthesis of active by the cross-linking of human TNF α and ricin A-chain, (A1) 3
- CZERNY C *see* Kainberger F
- DA SILVA JAP *see* Hall GM
- DACRE JE, GRIFFITH SM, JOLLY BC. Rheumatology and medical education in Great Britain, 269
- DAHINDEN C *see* Lehmann T
- DAIMARU Y *see* Kobayashi H
- DALLARD MC *see* Hellio le Graverand MP
- DALVIT G *see* Weber M
- DALZELL N *see* Tobias JH
- DALZIEL M, BOND A, REES DHE, HAY FC, AXFORD JS. Sialylation of serum immunoglobulin G in rheumatoid arthritis in relation to galactosylation and rheumatoid factor expression, (A1) 175
- DALZIEL M *see also* Awadallah F
- DAMATO R *see* Ferraccioli GF
- DAMEN CA *see* van der Veen MH
- Danazol, successful therapy in refractory thrombocytopenia associated with autoimmune diseases, (A1) 55
- DANOFF D *see* Newkirk MM
- DARBOIS Y *see* Le Thi Huong D
- DARKE C *see* Hall FC
- DASGUPTA B, DOLAN AL, PANAYI GS, FERNANDES L. Controlled trial of intramuscular methylprednisolone vs oral prednisolone in polymyalgia rheumatica, (A1) 59
- DASGUPTA B, WORRALL JG, WIND K, BELL L. Survey of NSAID usage in hospital and community and serious NSAID related gastrointestinal bleeding, (A1) 218
- DASGUPTA B *see also* Dolan AL; Ismail AA
- DASHWOOD MR *see* Shiwen X
- DAVID J *see* Turner HE; Woods RJ
- DAVIDSON BKS, McDONAGH JE, FOSTER HE, CAVANAGH G, GRIFFITHS ID, WALKER DJ. Prevalence of secondary Sjögren's syndrome in multicase RA families, (A2) 10
- DAVIDSON JE *see* Elliott M; Lees C
- DAVIE MWJ *see* Martin JC
- DAVIES EJ, PEPPER L, THE LS, ORDI J, OLLIER WER. HLA-DRB1 and complement C4 null alleles in systemic lupus erythematosus in a Spanish population, (A1) 26
- DAVIES EV *see* Lloyds D
- DAVIES KA. Complement, immune complexes and systemic lupus erythematosus, 5
- DAVIES KA, THEODORIDIS E, BOTTO M, THOMPSON EM, ISENBERG DA, BRIGGS D, BEYNON HLC, WALFORD MJ. IgG Fc γ IIA polymorphism in SLE: genotypic analysis in Caucasian, Chinese and Afro-Caribbean patients, (A1) 27
- DAVIES KA *see also* Athanassiou P; Hogarth MB; Slingsby JH
- DAVIES P *see* Prouse P; Talamo J
- DAVIES RP *see* Rischmueller M
- DAVIES EJ *see also* Hajeer AH; Stanworth SJ

- DAVIS DM, CHARLES P, ELLIOTT M, POTTER A, COPE A, FELDMAN M, MAINI RN. Changes in naturally-occurring cytokine inhibitor levels after TNF α blockade in rheumatoid arthritis, (A1) 181
- DAVIS DM, ELLIOTT MJ, CHARLES P, MAINI RN, FELDMAN M. Role of TNF α as an inhibitor of erythropoiesis in rheumatoid arthritis: insights from TNF α blockade *in vivo*, (A1) 217
- DAWES P *see* Hajer A; Hassell A; Stanworth SJ
- DAWES PT *see* Birrell FN; Cheung NT; Close DR; Plant MJ; Ryan SJ; Salih AM; Scott LJ
- DAWSON JK
Health economics as an aspect of health outcome: basic principles and application in rheumatoid arthritis, (L) 494
Thyroid autoimmunity in systemic lupus erythematosus: the clinical significance of a fluctuating course, (L) 195
- DAWSON JK, GRAHAM DR, KENNY J, DU BOIS RM, LYNCH MP. Clinical features of HRCIT diagnosed interstitial lung disease associated with rheumatoid arthritis, (A2) 16
- DAY A *see* Howat S
- D'CRUZ D *see* Ng T; Ross EL
- DE ABREU R *see* Stolk J
- DE BANDT M. ANCA in RA patients, (L) 1329
- DE BANDT M, MEYER O, HAIM T, KAHN M-F. Antineutrophil cytoplasmic antibodies in rheumatoid arthritis patients, 38
- DE BANDT M *see also* Cordonnier C
- DE BRASI D *see* Scarpa R
- DE CANDIA A *see* Fiocco U
- DE GRAAF R *see* Stolk J
- DE KLERK E, VAN DER LINDEN SJ. Compliance monitoring of NSAID drug therapy in ankylosing spondylitis, experiences with an electronic monitoring device, 60
- DE KONING D *see* Stolk J
- DE LA FUENTE R *see* Graña J
- DE LA IGLESIA JL, MORENO I, POVEDANO J, PÉREZ-VILCHEZ D, QUIJADA J, ALBORNOZ J, GARCÍA-LÓPEZ A. Association of the shared epitope with radiological severity of rheumatoid arthritis, (A1) 187
- DE LA IGLESIA JL, POVEDANO J, GARCÍA A, GARCÍA-PESQUERA F, PÉREZ-VILCHEZ D, SILVA H, ASTORGA R, LEAL-CERRO A. Somatotrophic axis in the fibromyalgia syndrome, (A1) 139
- DE LA IGLESIA JL, POVEDANO J, PÉREZ-VILCHEZ D, QUIJADA J, ALBORNOZ J, GARCÍA-LÓPEZ A, LEAL-CERRO A. Nocturnal growth hormone pulsatility and sleep study in patients with fibromyalgia, (A1) 138
- DE LA IGLESIA JL, POVEDANO J, QUIJADA J, GARCÍA A, PÉREZ-VILCHEZ D, ALBORNOZ J. Atypical chest pain of musculoskeletal origin, (A1) 136
- DE LA IGLESIA JL *see also* Albornoz J
- DE LA MATA J, FERNÁNDEZ-DAPICA P, POVEDA MJ, JIMENEZ M, VEIGA R, RESINES C. Synovial chondromatosis: prevalence in familial chondrocalcinosis, (A1) 147
- DE LA MATA J *see also* Fernández-Dapica P
- DE LEY M *see* Dolhain RJEM
- DE LORD DA. Case of chronic *Escherichia coli* osteomyelitis of the clavicle, (L) 192
- DE MIGUEL E, PLAZA MA, ARNALICH F, PANTOJA L, MARTÍN-MOLA E, HERNÁNZ A. Neuropeptides stimulate *in vitro* cytokine release in human leucocytes from rheumatoid arthritis in patients, (A1) 8
- DE MIGUEL E *see also* Rodríguez R
- DE VITA S *see* Ferraccioli GF
- DE WILDE EJ *see* van der Net J
- DECK R *see* Kohlman T
- Deflazacort and prednisolone effects on bone metabolism in polymyalgia rheumatica, (A1) 58
- DEGNER F *see* Distel M
- DEGNER FL *see* Frölich JC; Huskisson EC; Stichtenoth DO
- Degradation marker and bone formation relationship in serum and synovial fluid in rheumatoid patients, (A1) 91
- DEIGHTON C, DOHERTY M. Treatment advances—when and how to inform the public, 1175
- DEIGHTON CM. Towards a musculoskeletal care package, (A1) 126
- DEL VAL N *see* Rodríguez-Valverde V
- DELAMERE JP *see* Koutedakis Y
- DELLA CASA-ALABERIGHI O *see* Ferraccioli GF
- DELLESTABLE F. Heterotopic ossification complicating long-term sedation, (L) 700
- DELL'ORSO S *see* Patoia L
- DELVAL M *see* Belzunegui J
- DEMARS F, VAN DUKE CF, KIRK BA, KAPILA S, PETERFY CG, ROBERTS TPL, SHAMES DM, TOMAZIC S, MANN J, BRASCH RC. Mapping abnormal synovial vascular permeability in temporomandibular joint arthritis in the rabbit using MRI, (S3) 24
- DEMARS F *see also* Jevtic V
- Dengue arthritis, (L) 909
- DENMAN AM. Provision of rheumatological information, (E) 503
- DENMAN AM *see also* Steuer A
- DENTON CP, CARVALHO MD, STRATTON R, BLACK CM, PEARSON JD. Longitudinal study of anti-endothelial cell antibody activity in systemic sclerosis, (A1) 46
- DENTON CP, HOWELL K, STRATTON R, SHIWEN X, BLACK CM. Pilot study of low molecular weight heparin treatment in Raynaud's phenomenon and scleroderma, (A1) 43
- DENTON CP, SHIWEN X, PEARSON JD, BLACK CM. Endothelial cell derived interleukin 1 and basic fibroblast growth factor independently modulate scleroderma fibroblast properties, (A1) 41
- DENTON J *see* Coombes GM; Sanders PA
- DENTON CP *see also* Shiwen X
- DEODHAR AA, WOOLF AD. Bone mass measurement and bone metabolism in rheumatoid arthritis: a review, 309
- DEPRESSEUX G *see* Houssiau FA
- Depression
and quadriceps strength are important factors in knee pain and associated disability, (A1) 135
self-reported, prevalence in patients with rheumatoid arthritis, 879
- DEQUEKER J, BOONEN S, AERSENS J, WESTHOVEN R. Inverse relationship osteoarthritis-osteoporosis: what is the evidence? What are the consequences?, (E) 813
- Dermatology and rheumatology: sister specialties in Italy and Malta, 1312
- Dermatomyositis
T-lymphocyte immunophenotyping, 839
use of sequential DTPA clearance and high resolution computerized tomography in monitoring interstitial lung disease, 164
- DESOYA R *see* Martin JC
- DEVI B, NEILD G, ISENBERG DA. Analysis of renal disease and its involvement in mortality in patients with systemic lupus erythematosus, (A1) 37
- DEVLIN J, LILLEY J, GOUGH A, HUISSOON A, HOLDER R, REECE R, PERKINS P, EMERY P. Clinical associations of dual-energy X-ray absorptiometry measurement of hand bone mass in rheumatoid arthritis, 1256
- DEVLIN J, MOOTS R, ANDREWS D, BOVIN C, EMERY P. High intestinal permeability in patients with inflammatory arthritis is more strongly associated with NSAID therapy than with other factors, (A1) 221
- DEVLIN J *see also* Prouse P; Talamo J
- DEVOLLAER J-P *see* Houssiau FA
- DEVRIES E, BREEDVELD FC. Approach to therapeutic trials in rheumatology: the foundation for applied rheumatology research, 1000
- Diabetic cheiroarthropathy, ultrasonographic features, 676
- DIAMOND AG *see* Warnock MG
- DIAZ C *see* Amat M
- DIAZ C *see* Giral P
- DIAZ-FLORES L *see* Hdez-Beraián JA
- DICENTA-SOUSA M *see* Moragues-Pastor C
- DICKSON DJ. What do general practitioners expect from rheumatology clinics, (E) 920
- Diclofenac 75 and Arthrotec 75, comparison of effect on patient activity using the Numact activity monitor, (A2) 33
- Diclofenac sodium and meloxicam in osteoarthritis: a six-month, double-blind comparison, (S1) 39
- Diclofenac versus tenidap in rheumatoid arthritis, (L) 602
- Didronel, cyclical, effect on bone loss in early postmenopausal women, (A1) 87
- DIEPPE P *see* Creamer P
- DIEPPE PA *see* Sharif M
- DIEZ E, CUERVAS-MONS V, MULERO J, LABIANO I, LARREA A, ANDREU JL. Rheumatic manifestations in liver transplant recipients, (A1) 87
- DIEZ ILE *see* Isasi C
- DIEZ MANGLANO J *see* Pina Salvador JM
- Diffuse idiopathic skeletal hyperostosis (DISH), clinical manifestations, (E) 1193
- Digital neurovascular mechanisms in primary Raynaud's phenomenon and systemic sclerosis as assessed by iontophoresis and laser Doppler techniques, (A1) 48
- 1,25-Dihydroxyvitamin D₃ combined with cyclical etidronate on spine and femoral neck bone density in postmenopausal women, (A1) 115
- DUKMAN'S BAC *see* Simon CH
- DIMITRIYADIS I *see* Yazici H
- DIONISI MS *see* Patoia L
- DIRESKENELI H *see* Şahin (Özgün) Ş
- Disability
associated with rheumatic disease, survey in a rural population, (A1) 131
dissatisfaction with: repeated associations with mood and beliefs about function, (A1) 196
locomotor, association of abnormalities on physical examination of the hip and knee in the Rotterdam study, 884
overview, 184
- Disablement correlates in polyarticular juvenile chronic arthritis—a cross-sectional study, 91

- Disease activity**
 in ankylosing spondylitis: selection of a core set of variables and a first step in the development of a disease activity score, 867
 assessment of measures in ankylosing spondylitis: their relationship with acute phase reactants, (A1) 153
 and biochemical markers, relationships between clinical measures in rheumatoid arthritis, (A1) 230
 measurement of outcome in rheumatoid arthritis using the Nottingham Health Profile, (A1) 231
 remission in a prospective study of patients with rheumatoid arthritis. American Rheumatism Association preliminary remission criteria in relation to disease activity score, 1101
 in systemic lupus erythematosus, comparison of ethnic groups belong to 2 university hospitals, (A1) 34
 vs illness perception as predictors of health status and psychological well-being in patients with systemic lupus erythematosus: longitudinal analysis, (A1) 34
- Disease expression and HLA association of RA: comparison of Egypt and the UK, (A1) 185**
- Disease progression, is the control within our grasp? review of the GRISAR study, (S2) 8**
- Disease repercussion profile, impact of a common symptom: validation for knee pain, (A1) 133**
- Disease severity, is RA becoming a milder disease? or are we starting second-line therapy in patients with milder disease?, 1305**
- Disease status comparison in rheumatoid arthritis patients attending and not attending a specialist clinic, 1169**
- Disease susceptibility genes**
 dominant, effect on phenotype in BXSb lupus-prone mice, (A1) 112
 sensitivity of different analytical methods to detect in RA sibling pair families, (A1) 190
- Disease-modifying anti-rheumatic drug (DMARD) therapy, rational: is it possible?, (E) 707**
- Disease-modifying drug termination in psoriatic arthritis: study of 109 courses of treatment, 564**
- DISTEL M, MUELLER C, BLUMMKI E, FRIES J, DEGNER F. Safety of treatment with a new NSAID, meloxicam, which shows selectivity towards COX-2: a global analysis, (A1) 219**
- DISTEL M, MUELLER C, BLUMMKI E, FRIES J. Safety of meloxicam: global analysis of clinical trials, (S1) 68**
- DISTEL M see also Hosie J; Linden B; Reginster JY; Wojtulewski JA**
- DIXEY J see Balsa A; Haigh RC; McFarlane XA; Talamo J**
- DIXEY JJ see Martin JC**
- DMARDs**
 monitoring in daily rheumatological practice: are laboratory tests of any use?, (A1) 217
 response index for monitoring efficacy in rheumatoid arthritis, (A2) 21
- DNA allelic alterations within VNTR loci of scleroderma families, 1216**
- DOBRO J see Markov G**
- DODDS W see Halsey J**
- DOHERTY M see Deighton C; Lanyon P; Ledingham J; O'Reilly SC**
- DOHERTY SM, STEEL SA, PURDIE DW. Bone densitometry, a funded service—audit of referrals over a 2 year period, (A2) 5**
- DOLAN AL, DASGUPTA B, FERNANDES L, PANAYI GS, MONIZ C. ESR at presentation is a prognostic indicator for bone loss and duration of treatment in polymyalgia rheumatica, (A1) 58**
- DOLAN AL, MONIZ C, CORRIGALL V, DASGUPTA B, PANAYI GS. Systemic IL6 may be responsible for increased bone turnover in polymyalgia rheumatica, (A2) 4**
- DOLAN AL, RYAN PJ, ARDEN NK, STRATTON R, WEDLEY JR, HAMANN W, FOGELMAN I, GIBSON T. Value of SPECT scans in identifying back pain likely to benefit from facet joint injection, 1269**
- DOLAN AL see also Dasgupta B**
- DOLAN P see Adams MA**
- DOLHAIN RJEM, TER HAAR NT, HOEFAKKER S, TAK PP, DE LEY M, CLAASSEN E BREEDVELD FC, MILTENBURG AMM. Increased expression of interferon (IFN)-gamma together with IFN-gamma receptor in the rheumatoid synovial membrane compared with synovium of patients with osteoarthritis, 24**
- DOLHAIN RJEM see also Ronday HK**
- Domiciliary visits for locomotor problems, audit, (A2) 30**
- DONN RP, KAY RA, HOLT PLH, OLLIER WER. T-cell receptor variable gene segment TCRBV1254 null allele frequency in UK juvenile chronic arthritis, (A1) 80**
- DONN RP see also Stanworth SJ**
- DONNELLY S, LONG A, HALL N, MURPHY A, CASEY EB, KELLEHER D. Gold sodium thiomalate does not inhibit PKC mediated cellular activation for signalling through CD44 and LFA-1, (A1) 14**
- DORIA A see Fiocco U**
- DOUGADOS M see Ayral X; Cisman A; Ravaud P**
- DOUGHTY RN see Armstrong GP**
- DOYLE D see Cooper C**
- DOYLE DV see Arden NK; Griffiths GO; Houssien DA; Masud T; Nandra D; Taylor JC**
- DOYLE TCA see Highton J**
- DRB1*0404 association with polymyalgia rheumatica, 632**
- DRB1 associations in adult Still's disease, (A1) 78**
- DRESES-WERINGLOER U, BONK C, REISS G, ZEIDLER H, KOEHLER L. Effect of ciprofloxacin and interferon-gamma on the infection of *Chlamydia trachomatis* in the monocytic cell line THP-1, (A1) 169**
- Drug therapy, effect of second line anti-rheumatic therapy on gait characteristics in early rheumatoid arthritis, (A1) 212**
- Drug treatment, perception by patients with rheumatoid arthritis of the risk benefit ratio, (A1) 212**
- DRYLL A see Lioté F**
- DRYSDALE MJ see McNicholas MJ**
- DTPA clearance, sequential and high resolution computerized tomography in monitoring interstitial lung disease in dermatomyositis, 164**
- DU BOIS RM see Dawson JK**
- Dual-energy X-ray absorptiometry measurement of hand bone mass in rheumatoid arthritis, 1256**
- DUDDY K see Leak AM**
- DUDHIA J, WELLS TM, HARDINGHAM TE, BAYLISS MT. Age-related changes in the C-terminal of aggrecan from human cartilage, (A1) 236**
- DUDHIA J see also Bolton MC**
- DULIEU V see Maillefer JF**
- DUMAS GA see Mannion AF**
- DUNCAN M see Munro R**
- DUNCAN MR, McENTEGART A, PORTER DR, MADHOK R, THOMSON EA, CAPELL HA. Discrepancies in 5 year mortality in RA patients during second line therapy—cultural or procedural?, (A1) 234**
- DUNCAN MR see also Capell HA; McEntegart A; Morrison E**
- DUNN A see McDonagh JE**
- DUNNE CA, BEGLEY J, THOMPSON PW. Effect of intramuscular corticosteroid on markers of bone mineral metabolism in rheumatoid patients, (A1) 81**
- DUNNE E see Richmond R**
- DUNPHY J see Haigh RC; McFarlane XA**
- DURANDY A see Sibilia J**
- DURHAM B see Black AJ; MacDonald AG**
- DYCE C, BIDDLE P, HALL K, FERGUSON M, YOUNG A. Evaluation of extended role of physio and occupational therapists in rheumatology practice, (A1) 130**
- DYER PA see Haworth S**
- DZIEDZIC K see Avers HL**
- E-selectin**
 expression characterization, leucocyte traffic and clinical sequelae in urate crystal-induced inflammation: an insight into gout, 323
 treatment of rheumatoid arthritis with antibody to TNF α decreases expression and shedding, (A1) 218
- EASTELL R see Choy EHS; Cooper AM; Peel NFA**
- EBERHARDT K see Fex E**
- EBRAHIM S see Harwood RH**
- EBRINGER A see Tiwana H**
- Ed-B-containing fibronectin (an embryonic isoform of fibronectin), increased expression in human osteoarthritic cartilage, 407**
- EDELSTEIN C. Antibodies against retinal S-antigen in patients with juvenile chronic arthritis-associated uveitis, (L) 101**
- EDMONDS SE, HALL MA, GAFFNEY K, BLAKE DR. Intra-articular pressure measurement in children with juvenile chronic arthritis, (A1) 78**
- EDMUNDS S see Allen RL**
- Education**
 formal, influence on compliance and disease activity in early RA patients starting sulphasalazine, (A1) 216
 medical and rheumatology in Great Britain, 269
 patient: treatment or nice extra. The British experience, (L) 805
 pre-admission for patients undergoing total knee replacement, (A2) 31
 undergraduate in rheumatology, (E) 110, (E) 203
- Educational booklet, single-blind randomized controlled trial for patients with chronic arthritis, 775**
- EDWARDS DR see Clark IM**
- EDWARDS JCW see Leigh RD**
- EGGENS U see Braun J**
- EGGER P see Coggon D; Cooper C; Smedley J**
- EGIDO J see Gutierrez S; Miguélez R; Palacios I; Sánchez-Pernaute O**
- EHLLENZ K see Baerwald CGO**
- EHLERS S see Braun J**
- EHRENFELD M, BUSKILA D, TISHLER M, LORBER M, SHOENFELD Y. Rheumatology in Israel, 778**
- EHRENFELD et al. article, (L) 195**
- EHRENFELD MR see Ravirajan CT**
- EL RAMAHI KM, HALIM M, KARRAR A, AL DALAAN A, AL SHAIKH A. *Salmonella* infection in Saudi patients with systemic lupus erythematosus, (A1) 35**
- EL RAMAHI KM see also Kattar A**

- EL-BADAWY SA see Popert AJ
 EL-BARBARY M see Elbeialy A
 EL-BOGHADY IA, EL-SHAL EW, EL-DOSOKY ME. Primary fibromyalgia: effect of aerobic and breathing exercise on pulmonary ventilatory function, (A2) 33
 EL-BOGHADY IA see also El-Shal EW
 EL-DOSOKY ME see El-Boghdady IA
 EL-MANZALAWI E see Troughton PR
 EL-MOGY SA see El-Shal EW
 EL-SHAL EW, EL-BOGHADY IA, EL-MOGY SA, SIRAG SM. Low energy laser therapy: effect on papain-induced OA, (A2) 7
 EL-SHAL EW. Response index for monitoring efficacy of DMARDs in rheumatoid arthritis, (A2) 21
 EL-SHAL EW see also El-Boghdady IA
 ELBEIALY A, EL-BARBARY M, KAMEL M. Peripheral beta- endorphin in rheumatoid arthritis: a correlation with the disease activity, (A1) 193
 Electromagnetic field, pulsed 27 MHz, effect on fibroblast cell division cyclic AMP and inositol phosphate, (A1) 64
 Electron paramagnetic resonance study of the arthritic joint, (A1) 204
 ELIAS A see Cordonnier C
 ELISA specificity for antibody to β 2-glycoprotein I in patients with antiphospholipid syndrome, 1239
 ELKON KB see Ramos P
 ELLING H see Elling P
 ELLING P, OLSSON AT, ELLING H. Reduced CD8+ T-cell concentration in peripheral blood of patients with carotid artery stenosis: relation to arteritis temporalis, 649
 ELLIOTT M, HERRICK AL, SMITH CP, DAVIDSON JE. Polyarteritis nodosa in the North West Region, (A2) 38
 ELLIOTT M see Davis DM; Paleolog E
 ELLIOTT MJ see Davis DM
 ELLIS AJ, CURRY VA, LEAN EM, WARD D, CAWSTON TE. Interleukin-4 can prevent the release of collagen from cartilage stimulated with proinflammatory cytokines *in vitro*, (A1) 8
 ELLWOOD CN see Wong WM
 ELSLEY WAJ see Stagg AJ
 ELSON CJ see Leader KA; Perry MJ; Webb GR
 ELSON CJE see Sharif M
 ELVINS D see Minaur NJ
 EMERY D, GOW P, GRAY DH. Study of foot problems of a consecutive group of patients with rheumatoid arthritis, (A1) 198
 EMERY P, VAN DE PUTTE LBA. New perspectives on controlling progression of severe rheumatoid arthritis—summary of the potential role of cyclosporin, (S2) 1
 EMERY P see also Devlin J; Gordon C; Griffiths B; Prouse P; Pugh MT; Reece RJ; Talamo J
 EMOI I see Chevalier X
 Employment
 attrition in a community based inception cohort of rheumatoid arthritis patients, (A1) 235
 and work activities as predictors of future back pain, (A1) 141
 Endogenous opioid tone in patients with rheumatoid arthritis, 436
 beta-Endorphin in rheumatoid arthritis: a correlation with the disease activity, (A1) 193
 Endothelial activation in urate crystal-induced inflammation: *in vitro* and *in vivo* studies on the role of tumour necrosis factor, (A1) 146
 Endothelial cell derived interleukin 1 and basic fibroblast growth factor independently modulate scleroderma fibroblast properties, (A1) 41
 Endothelial cells and β 2-glycoprotein I antibodies in the antiphospholipid syndrome: prevalence and isotype distribution, 523
 Endothelial damage is an important mechanism in the pathogenesis of rheumatoid neuropathy, (A1) 170
 Endothelial proliferation index in rat and human tissues, (A1) 10
 Endothelial responses and cellular immunity, contemporaneous activation in limited cutaneous systemic sclerosis, (A2) 27
 Endothelin, differential modulation of scleroderma and control fibroblast proliferation and collagen biosynthesis, (A1) 47
 Endothelin-1 expression in serum and bronchoalveolar lavage from patients with active Behçet's disease, 357
 ENGELHARDT G. Pharmacology of meloxicam—a new non-steroidal anti-inflammatory drug with an improved safety profile through preferential inhibition of COX-2, (S1) 4
 ENGELHARDT G see also Frölich JC
 ENGLAND AJ, REID DM, CAPELL H, RALSTON SH. Effects of oral clodronate on disease activity and bone mineral density in rheumatoid arthritis, (A2) 21
 ENGLAND AJ, REID DM. Effects of bone disease duration in rheumatoid arthritis on appendicular bone mineral measurements, (A1) 86
 ENGLAND AJ see also Murray R; Reid DM
 Enterococcal arthritis with avascular necrosis in a lupus patient, 595
 Epithelitis, autoimmune, Sjögren's syndrome revisited, (E) 204
 Epstein-Barr virus, rheumatic disease patients, prone to Sjögren's syndrome and/or lymphoma, mount an antibody response to BHRF1, the Epstein-Barr viral homologue of BCL-2, 1075
 ERAUSQUIN C see Belzunegui J; Rodriguez-Valverde V
 ERHARDT CC see Bamji AN
 Erythroblast iron metabolism, evidence for abnormal in the chronic anaemia of rheumatoid arthritis, (A1) 191
 Erythropoiesis, role of TNF α as an inhibitor in rheumatoid arthritis: insights from TNF α blockade *in vivo*, (A1) 217
Escherichia coli osteomyelitis of the clavicle, chronic case, (C) 192
 ESCOLÀ A see Obach J
 ESCOLÀ A, ALEGRE C, OBACH J, PLANS P, ASENSI E. Risk factors of the hand, hip and knee osteoarthritis in Catalonia-II survey, (A1) 101
 ESCOLAR G see Navarro M
 ESCRIBANO M see Perez A
 ESPINOSA FJ see Mannion AF
 ESPINOSA G see Garcia-Carrasco M
 ESR
 at presentation is a prognostic indicator for bone loss and duration of treatment in polymyalgia rheumatica, (A1) 58
 inappropriately high results—a feature of amyloid complicating rheumatic disorders?, (A1) 77
 ESSEX R see Walker DJ
 ESTEVE M see Olivé A
 ETHERINGTON J, ARDEN N, BROAD I, BUCHANAN N, SPECTOR TD, HUGHES GRV. Serum sex hormones in pregnancy: comparison between lupus, rheumatoid and control patients, (A1) 35
 Etidronate, cyclical combined with 1,25-dihydroxyvitamin D₃ on spine and femoral neck bone density in postmenopausal women, (A1) 115
 European Study Group for Therapeutic Trials in Systemic Vasculitis see Luqmani RA
 European Union, specialist training into the new millenium: the price of conformity with Europe, (E) 607
 European Vertebral Osteoporosis Study Group see O'Neill TW
 EVANS AR see Vakalis AN
 EVANS EL see Close DR
 EVANS MD, WILLIAMS SML, PERRETT D, LUNEC J, AKIL M, NICHOL FE, HERBERT KE. Susceptibility of peripheral blood mononuclear cells from RA patients to peroxide-induced DNA damage, (A1) 184
 EVANS WD see Byrne PAC
 EVISON G see Balakrishnan C
 EXLEY A, BANBURY P, CARRUTHERS D, MOOTS R, LUQMANI R, KITAS G, BACON P. Vasculitis damage index, (A1) 53
 EXLEY A see also Luqmani RA
 Extra-articular features of benign joint hypermobility syndrome, 861
 Eye, rheumatic disorders and of the various structures involved, 258
 Facet joint injection, value of SPECT scans in identifying back pain likely to benefit from, 1269
 Factor V gene mutation increases the risk of venous thrombosis in Behçet's disease, 1178
 FALCO P see Ferri C
 Familial Mediterranean fever attacks, resolving with interferon alpha, (L) 1188
 Familial tendency to severe symptomatic osteoarthritis of the hip and knee, (A1) 100
 FAN T-PD see Walsh DA
 FAN W see Horsfall AC
 FANNING G see McHugh NJ
 FANTINI F. Is cyclosporin a disease-controlling antirheumatic therapy? review of the two-year results of the GRISAR study, (A1) 215
 FANTINI F see also Ferraccioli GF; Marchesoni A
 FARAGHER EB see Taylor DJ
 FARIS FJ see Mannion AF
 FARR M see Jawed S
 FARRELL R see McGonagle D
 Fas gene, human, polymorphism of the extracellular domain and its relationship with SLE, (A1) 25
 FATIMA Z see Gibson T
 Fatty acid plasma pattern in rheumatoid arthritis, (A1) 191
 FAVARO MA see Fiocco U
 Fc γ RIIA polymorphism in SLE and RA, (A1) 28
 FEAR J see Tennant A
 FEEST TG see Hawkins PN
 FEIGHERY C see Guerin J
 FELDMANN M see Brennan FM; Davis DM; Gould DJ; Kissnerghis M; Malfait A-M; Marinova L; Mauri C; Paleolog E; Ross S
 FELLOWES R see Kissnerghis M
 FELSENBURG D see O'Neill TW
 Felt's syndrome
 anti-TCRBV monoclonal antibodies in the detection and monitoring of large granular lymphocyte leukaemia, (A1) 182

- large granular lymphocyte expansions have an unusual phenotype of activated CD45RA+ cells, 1252
- Female patients, are we serving them less well than males? Baseline characteristics of rheumatoid arthritis patients enrolled into sulphasalazine studies, (A1) 213
- Femur, proximal, geometric measurements in UK women: secular increase between the late 1950s and early 1990s, (A1) 90
- FENN GC, MORANT S, SCOTT DL. Efficacy of misoprostol in ulcer healing with continued NSAID therapy, (A1) 222
- FERGUSON M *see* DYCE C
- FERGUSON WP *see* Kane D
- FERNAND JP *see* Sibilia J
- FERNANDES L *see* Dasgupta B; Dolan AL
- FERNANDEZ F *see* Olivé A
- FERNANDEZ-COUTO D *see* Graña J
- FERNÁNDEZ-DAPICA P, PATO E, POVEDA MJ, SANCHEZ-NISTAL MA, LARRODERA L, SERRANO L, DE LA MATA J. Spondiloeipiphyseal dysplasia associated with familial CPPD deposition disease: study of a pedigree, (A1) 147
- FERNÁNDEZ-DAPICA P *see also* De La Mata J; Poveda MJ
- FERNÁNDEZ-ECHEVARRIA MA *see* Calvo-Alén J; Sánchez-Andrada S
- FERNÁNDEZ-ESPARTERO MC *see* Abad MA; Carreño L
- FERNÁNDEZ-RODRIGUEZ A *see* Albornoz J
- FERNÁNDEZ-SUEIRO JL *see* González-Gay MA
- FERNIHOUGH JK, BILLINGHAM MEJ, CWFYAN-HUGHES S, HOLLY JMP. Disruption of the IGF system in osteoarthritic and rheumatoid patients compared to non-arthritic volunteers, (A1) 105
- FERONE D *see* Scarpa R
- FERRACCIOLI GF, DE VITA S, PEGORARO I, CASATTA L, DAMATO R. Immunological features of patients with primary Sjögren's syndrome and patients HCV positive with sicca syndrome, (A1) 15
- FERRACCIOLI GF, DELLA CASA-ALBERIGHI O, MARUBINI E, PRIOLIO F, MATHIEU A, FANTINI F, CUTOLO M, PASERO G. Is the control of disease progression within our grasp? review of the GRISAR study, (S2) 8
- FERRACCIOLI GF. GRISAR study, assessment of side effects, withdrawals and survival on treatment after 24 months, (A1) 215
- FERRAZ MB *see* Ciconelli RM
- FERRAZZI V *see* Jorgensen C
- FERRERA A *see* Muñoz-Fernández S
- FERRI C, GIUGGIOLI D, MARTINI A, LA CIVITA L, PASERO G, GIANNONI A, FALCO P, ROSSI A, CONTI L. Psychological and psychopathological evaluation of systemic sclerosis and rheumatoid arthritis patients, (A1) 47
- FERRI C, LA CIVITA L, MONTI M, LOMBARDINI F, GIANNINI C, LONGOMBARDI G, BOMBARDIERI S, ZIGNEGO AL. Mixed cryoglobulinaemia and hepatitis C virus genotypes, (A1) 50
- Ferritin H-chain mRNA is upregulated in human articular cartilage stimulated with TNF- α , (A1) 4
- Ferryl and/or hydroxyl radicals at sites of inflammation, indirect evidence to support production, (A2) 13
- FERRUCCI MG *see* Scarpa R
- FEX E, JONSSON K, JOHNSON U, EBERHARDT K. Development of radiographic damage during the first 5-6 yr of rheumatoid arthritis. Prospective follow-up study of a Swedish cohort, 1106
- Fibrinogen in systemic lupus erythematosus: an atherothrombotic link?, (A1) 122
- Fibrinolysis abnormalities in primary antiphospholipid syndrome: sex dependency and clustering with low free protein S, (A1) 38
- Fibroblast cell division cyclic AMP and inositol phosphate, effect of pulsed 27 MHz electromagnetic field, (A1) 64
- Fibroblast growth factor and endothelial cell derived interleukin 1 independently modulate scleroderma fibroblast properties, (A1) 41
- Fibromyalgia
comparison of tenoxicam and bromazepam in treatment. A randomized, double-blind, placebo-controlled trial, (A1) 138
decreased plasma amino acid concentrations, (A1) 137
nocturnal growth hormone pulsatility and sleep study in patients with fibromyalgia, (A1) 138
primary: effect of aerobic and breathing exercise on pulmonary ventilatory function, (A2) 33
syndrome, somatotroph axis, (A1) 139
- Fibronectin
40 kDa collagen-binding fragment, proteoglycan-degrading activity associated with, 506
and proteoglycan proliferation and synthesis, SAME restores the changes induced by TNF α on cultured synovial cells, (A1) 63
soluble, interactions with isolated human articular chondrocytes, (A1) 62
- FIELD M *see* Crilly A; McInnes IB
- FIELDER M *see* Tiwana H
- FIGUEROA M *see* Belzuncegui J; González-Gay MA
- FILIPPA D *see* Ramos P
- FINCH MB *see* Smedstad LM
- FINN J *see* Jones M
- FINN JD *see* Jones MA
- Finnish rheumatology, current trends, 489
- FIOTTO U, COZZI L, RIGON C, CHIECO-BIANCHI F, BALDOVIN M, CASSISI GA, GALLO C, DORIA A, FAVARO MA, PICCOLI A, DE CANDIA A, RUBATELLI L, TODESCO S. Arthroscopic synovectomy in rheumatoid and psoriatic knee joint synovitis: long-term outcome, 463
- FIOTTO U, COZZI L, RUBATELLI L, RIGON C, DE CANDIA A, TREGNAGHI A, GALLO C, FAVARO MA, CHIECO-BIANCHI F, BALDOVIN M, TODESCO S. Long-term sonographic follow-up of rheumatoid and psoriatic proliferative knee joint synovitis, 155
- FISHER J *see* Cheung NT
- FISHER P, SCOTT DL. Is homeopathy effective in rheumatoid arthritis?, (A1) 227
- FISHMAN D, WOO P. 5' regulatory region polymorphisms in the interleukin-6 gene of patients with systemic onset JCA and normal control subjects, (A1) 79
- FITER J, NOLLA JM, GÓMEZ-VAQUERO C, ROS S, ARIZA J, ROIG-ESCOFET D. Septic spondylodiscitis induced by pyogenic germs in patients without addition to intravenous drugs, (A1) 165
- FITER J *see also* Gómez-Vaquero C
- FITZGERALD O *see* Bolger T; Costello P; Cunnane G; Mulherin D
- FITZPATRICK F, BADLEY EM. Overview of disability, 184
- FITZSIMONS EJ *see* Houston T
- Flatfeet in pregnancy, (L) 396
- FLETCHER A *see* Yeap SS
- FLETT SM *see* Armstrong GP
- FLORA R *see* Mackay K
- FLORA RS, JONES SM, BALAKRISHNAN C, HALL ND, MCHUGH NJ. Prevalence of the rheumatoid arthritis shared epitope in patients with psoriatic arthritis, (A1) 159
- FLOWER RJ *see* Perretti M
- Fluoride pharmacokinetics and changes in lumbar spine and femoral neck bone mineral density, (A1) 84
- FOEY A, CRAWFORD A, HALL N. Inhibition of the Na/K-ATPase exerts differential effects on cytokine production by monocytes and fibroblasts, (A1) 67
- FOGELMAN I *see* Dolan AL; Mishra MB
- Folate supplementation and methotrexate, (L) 1031
- FOLEY-NOLAN D, CLARK B, MONTAGUE B, GOOI HC. DRB1 associations in adult Still's disease, (A1) 78
- FOLEY-NOLAN D *see also* Mullaney P
- FONT C, MIRÓ O, PEDROL E, MASANÉS F, COLL-VINENT B, CASADEMONT J, CID MC, GRAU JM. Polyarthritis nodosa in human immunodeficiency virus infection: report of four cases and review of the literature, 796
- FONT J *see* García-Carrasco M; Navarro M
- FONTAN G *see* Muñoz-Fernández S
- Food intolerance, scurvy, osteoporosis and megaloblastic anaemia, (L) 701
- Foot
deformity development in early rheumatoid arthritis, (A1) 198
valgus heel deformity and plantar pressure distribution of the forefoot in rheumatoid arthritis, (A1) 199
- Foot problems
isotope bone scans, (A2) 8
study of a consecutive group of patients with rheumatoid arthritis, (A1) 198
- FORDHAM JN *see* Leitch DN; Speed CA
- Fore-arm bone mineral density in non-steroid treated rheumatoid arthritis is reduced in patients on long-term methotrexate compared with other disease-modifying drugs, (A1) 209
- FORREST GR *see* Cooke SP
- FOSTER HE, CABRAL DA, MALLESEN PN, PETTY RE. High titre anti-RNP antibody in association with carcinoma in a child who did not have a rheumatic disease, (A2) 37
- FOSTER HE *see also* Davidson BKS
- FOURNIER M *see* Lioté F
- FOX R *see* Lim KL
- FOX RA, ATKINS M, ISENBERG DA. Human immunodeficiency virus infection in a patient with systemic lupus erythematosus, (A2) 35
- FOXWELL BMJ *see* Gould DJ
- FRAIX E *see* Calabozo M; Perez-Ruiz F
- FRANCES C *see* Le Thi Huong D
- FRANSEN MJAM *see* Creemers MCW
- FRASER SM *see* Hamilton J
- FRASER WD *see* Black AJ; MacDonald AG
- FREARS ER *see* Zhang Z
- FREDRIKSON E *see* Rydgren L
- FREEDMAN J *see* Pattison D
- FREEMONT AJ. Pathophysiology of cartilage and synovium, (S3) 11
- FREIRE M *see* Graña J
- FREIRE-MOAR J *see* Kang RY
- FRIARD H *see* Lioté F
- FRIEMANN J *see* Wedekind F
- FRIES J *see* Distel M

- FRÖLICH JC, STICHTENOTH DO, PAIRET M, ENGELHARDT G, DEGNER FL. Selective inhibition of inducible cyclooxygenase by meloxicam: summary of pharmacological data and clinical pharmacology, (A1) 221
- FRÖLICH JC *see also* Stichtenoth DO
- FRYER A *see* Hajer A
- FUCHIGAMI T *see* Kobayashi H
- FUKUKAWA Y, SHINGU M, TORISU T, ITONAGA I, MASUMI S. Bone resorption by tartrate-resistant acid phosphatase-positive multinuclear cells isolated from rheumatoid synovium, 213
- FUJISHIMA M *see* Kobayashi H
- Functional disability correlates in early rheumatoid arthritis: a cross-sectional study of 706 patients in four European countries, 746
- FURIÓ MP *see* Belmonte MA
- FURNO P *see* Patoia L
- FURST DE. Rational disease-modifying anti-rheumatic drug (DMARD) therapy: is it possible?, (E) 707
- FURST DE *see also* Seibold JR
- FURUKAWA M, SAKAMOTO A, KITA Y, OHISHI Y, MATSUMURA R, TSUBATA R, TSUBATA T, IWAMOTO I, SAITOH Y, SUMIDA T. T-cell repertoire of infiltrating T cells in lacrimal glands, salivary glands and kidneys from alymphoplasia (*aly*) mutant mice: a new model for Sjögren's syndrome, 1223
- Gadolinium-enhanced magnetic resonance imaging
correlates with synovial vascularity in rheumatoid arthritis, (A1) 204
of the finger joints in rheumatoid arthritis, (A1) 118
quantitative assessment of synovial inflammation. A study of the effect of intra-articular methylprednisolone on the rate of early synovial enhancement, 50
- GAFFNEY A. Rheumatoid arthritis and heparin, (L) 808
- GAFFNEY K. Antineutrophil cytoplasmic antibody-positive polyarthritis associated with minocycline therapy, (L) 1327
- GAFFNEY K, COOKSON J, BLADES S, COUMBE A, MAPP P, WALSH D, BLAKE DR. Gadolinium enhanced magnetic resonance imaging correlates with synovial vascularity in rheumatoid arthritis, (A1) 204
- GAFFNEY K, JAWED S, BLAKE DR. Intra-articular pressure dynamics in inflammatory arthropathies, (A1) 71
- GAFFNEY K *see also* Edmonds SE; Naughton DP; Nazhat NB; Ross EL
- GAGAN RM *see* Salih AM
- Gait characteristics, effect of second line anti-rheumatic therapy in early rheumatoid arthritis, (A1) 212
- β 1,4-Galactosyltransferase isoenzyme changes in serum of patients with rheumatoid arthritis, (A1) 174
- GALLAGHER AC, CHOW JWM, CHAMBERS TJ, TOBIAS JH. Serum cross-linked telopeptides of type I collagen are a useful marker of bone resorption in the rat, (A1) 91
- GALLAGHER AC *see also* Tobias JH
- GALLARDO D *see* Lience E
- GALLIMORE JR *see* Nandra D; Persey MR
- GALLINARI C *see* Le Thi Huong D
- GALLIVAN S *see* Talamo J
- GALLO C *see* Fiocco U
- GAMBLE GD *see* Armstrong GP
- GANDY S, HUTTON CW, JACOBY RK, VENNART W. Magnetic resonance imaging of acromegaly, (A1) 95
- GANGI M *see* Ramos P
- GANTES M *see* Hdez-Berriain JA
- GARCIA A *see* De La Iglesia JL
- GARCIA M *see* Camacho M
- GARCIA-CARRASCO M, RAMOS M, CERVERA R, FONT J, MUÑOZ FJ, VIVANCOS J, MIRET C, ESPINOSA G, INGELMO M. Liver involvement and hepatitis C virus infection in patients with primary Sjögren's syndrome, (A1) 15
- GARCIA-FUENTES M *see* Blanco R
- GARCIA-LÓPEZ A *see* Alborno J; De La Iglesia JL
- GARCIA-MONFORTE A *see* Rivera J
- GARCIA-PESQUERA F *see* De La Iglesia JL
- GARDINER DM *see* Highton J
- GARIJO F *see* Val-Bernal F
- GARNERO P *see* Keen RW
- GARRETT NE *see* Cruwys SC
- GARRETT SL *see* Jones SD; Kabasakal Y
- GARTON MJ, ISENBERG DA. Idiopathic versus lupus myositis: comparison of clinical and laboratory features, (A1) 48
- GARTON MJ, ISENBERG DA, REICHLIN MW, REICHLIN M. Clinical features and autoantibodies in black female lupus patients: cross-sectional and longitudinal studies, (A1) 30
- GASSON J, HUTTON CW, JACOBY RK, VENNART W. High-resolution magnetic-resonance imaging of erosive changes in the metacarpophalangeal joint in rheumatoid arthritis, (A1) 124
- GASTON JSH *see* Sibanda EN
- Gastrointestinal bleeding, NSAID-induced, incidence and severity, (A1) 220
- GASULL MA *see* Olivé A
- GATLEY M *see* Malfait A-M
- GAUSI C *see* Moragues-Pastor C
- GEDDES GC *see* Bowman SJ
- GELI C *see* Giralp P
- General practice requirements from the rheumatology services in Leicestershire, (A2) 29
- General practitioners
do they refer to rheumatologists on the basis of symptoms or as a result of tests?, (A2) 30
what they expect from rheumatology clinics, (E) 920
- General public's perceptions about rheumatic diseases, (A1) 126
- Genetic analysis of TAP2 in systemic lupus erythematosus patients from two ethnic groups, 529
- Genetic anticipation, does it occur in ankylosing spondylitis?, (A2) 24
- GERBER NJ *see* Beyeler C; Lehmann T
- GERLI R *see* Biagini P
- GERHARDI R *see* Guillemin L
- GHOZLAN PR, BERNHARDT M, VÉLICITAT P, BLUHMKE E. Tolerability of multiple administration of intramuscular meloxicam: comparison with intramuscular piroxicam in patients with rheumatoid arthritis or osteoarthritis, (S1) 51
- GHOZLAN R *see* Huskisson EC
- GIANNINI C *see* Ferri C
- GIANNONI A *see* Ferri C
- Giant cell arteritis
and amyloidosis, lack of association between, (A1) 57
low-dose corticosteroids and blindness, (L) 702
ocular involvement: study of 191 patients in 15 years, (A1) 57
and pericardial effusion, (L) 194
scalp necrosis, (A2) 36
three year follow-up of a case presenting with a chronic cough and upper limb ischaemic symptoms, 800
- GIBBINS J *see* Woods RJ
- GIBBONS DL *see* Brennan FM
- GIBSON K *see* Hall FC; Rudwaleit M
- GIBSON T, HAMEED K, KADIR M, SULTANA S, FATIMA Z, SYED A. Knee pain amongst the poor and affluent in Pakistan, 146
- GIBSON T *see also* Dolan AL; Hameed K; Rojkovich B
- GUÓN BAÑOS J *see* Muñoz-Fernández S
- GUÓN-BAÑOS J *see* Usón J
- GILLESPIE GMA *see* Allen RL
- GIRALT P, DIAZ C, GELI C, LLOBET JM, CASAN P, SÁNCHEZ J. Evolution of the pulmonary function in ankylosing spondylitis, (A1) 151
- GIUGGIOLI D *see* Ferri C
- GLADMAN DD *see* Koh ET
- GLANT TT *see* Brennan FR
- GLENNAS A *see* Bas S; Kvien TK
- GLEZ-GAY MA *see* Graña J
- Glomerulonephritis, cryoglobulinaemic, complicating case of primary Sjögren's syndrome, with pericardial and pleural effusions, 72
- Glucocorticoid
induced inhibition of adhesion molecule expression is reversed by IL-2, but not by TNF- α , IL-1 β /IL-4 or IL-10, (A1) 7
novel action—NF- κ B inhibition, (E) 709
treated patients, ultrasound treated patients, (A2) 5
- beta2-Glycoprotein I
antibodies recognise a conformational epitope and are highly specific for the antiphospholipid syndrome, (A1) 21
antibody specificity in patients with the antiphospholipid syndrome, (A1) 24
and endothelial cell antibodies in the antiphospholipid syndrome: prevalence and isotype distribution, 523
and human monoclonal anticardiolipin antibodies effects on the protein S / C4B-binding protein system, (A1) 19
- Glycosylation changes of IgG during fasting in patients with rheumatoid arthritis, 117
- GODEAU P *see* Le Thi Huong D
- GODOY H *see* Abad MA; Monteagudo I
- GOHCHI K *see* Matsuda J
- Gold
injectable compounds: an overview, 1154
neurological signs in RA patients, (L) 803
plated stomach—myth or reality, (A1) 225
- Gold sodium thiomalate
association of HLA with toxicity in patients with rheumatoid arthritis in a Spanish population, (A1) 211
does not inhibit PKC mediated cellular activation for signalling through CD44 and LFA-1, (A1) 14
- Gold therapy
intramuscular in 440 rheumatoid patients—results over 5 years in relation to level of social deprivation at outset, (A1) 213
parenteral, adding low-dose cyclosporin A in rheumatoid arthritis: a double-blind placebo-controlled study, 1142

- GOLDMAN J *see* Naughton M
- GOMEZ K, KENNEDY LG, CALIN A. Does genetic anticipation occur in ankylosing spondylitis? (A2) 24
- GÓMEZ-REINO JJ *see* Poveda MJ
- GÓMEZ-VAQUERO C, JUANOLA X, NAVARRO MA, MATEO L, ROS S, ROIG-ESCOFET D. Serum prolactin and salivary testosterone in young and middle-aged men with ankylosing spondylitis, (A1) 152
- GÓMEZ-VAQUERO C, RODRÍGUEZ-MORENO J, ROS S, MARCOS R, FITER J, ROIG-ESCOFET D. Termination of disease-modifying drugs in psoriatic arthritis: study of 109 courses of treatment, 564
- GÓMEZ-VAQUERO C *see also* Fiter J
- GOMEZ K *see also* Kennedy LG
- GONZALEZ C *see* Belzuncgui J
- GONZÁLEZ CM *see* Abad MA; Carreño L; González M
- GONZÁLEZ E *see* Miguélez R
- GONZALEZ E *see* Palacios I
- GONZÁLEZ J, ROLDÁN R, MARTÍNEZ-CONCHA D, CARACUEL MA. Economic nailfold capillaroscopy study in Raynaud's phenomenon: follow-up and evaluation as prognostic value, (A1) 49
- GONZÁLEZ M, GONZÁLEZ CM, LÓPEZ-LONGO FJ, MORENO AC, MINGUEZ MD, GRAU R, ARIZA A, MONTEAGUDO I, CARREÑO L. Diagnostic relevance of anti-Sm antibodies detected by ELISA in systemic lupus erythematosus, (A1) 19
- GONZÁLEZ M *see also* Carreño L; López-Longo FJ
- GONZÁLEZ MR *see* Poveda MJ
- GONZÁLEZ T *see* Hdez-Berriain JA
- GONZÁLEZ-COMPTA X *see* Moragues-Pastor C
- GONZÁLEZ-DOMÍNGUEZ J, KINDELAN JM, SALMORAL A, PERULA L, TORRE-CISNEROS J, JURADO R. Osteoarticular tuberculosis in two different populations: the HIV infection impact, (A1) 166
- GONZÁLEZ-GAY MA, BLANCO R, MARTÍNEZ-TABOADA VM, ARMONA J, FERNÁNDEZ-SÚEIRO JL, FIGUEROA M, RODRÍGUEZ-VALVERDE V. Polymyalgia rheumatica with a slightly increased erythrocyte sedimentation rate, (A1) 56
- GONZÁLEZ-GAY MA, GONZÁLEZ-VELA C, BLANCO R, SANTIAGO J, CEREJO MJ. Lack of association between biopsy-proved giant cell arteritis and amyloidosis, (A1) 57
- GONZÁLEZ-GAY MA. Low-dose corticosteroids and blindness in giant cell arteritis, (L) 702
- GONZÁLEZ-GAY MA, MARTÍNEZ-TABOADA VM, BLANCO R, FERNÁNDEZ-SÚEIRO JL, ARMONA J, FIGUEROA M, RODRÍGUEZ-VALVERDE V. Ocular involvement in giant cell arteritis: study of 191 patients in 15 years, (A1) 57
- GONZÁLEZ-GAY MA *see also* Blanco R
- GONZÁLEZ-VELA C *see* González-Gay MA; Val-Bernal F
- GOODACRE JA *see* Warnock MG
- GOODFELLOW RM, WILLIAMS AS, MORGAN BP, WILLIAMS BD. Administration of local and systemic soluble complement receptor one (SCR1) has a beneficial effect on the development of rat mono-articular arthritis, (A1) 74
- GOODFELLOW RM *see also* Williams AS
- GOOI HC *see* Foley-Nolan D; Griffiths B
- GORDON C, MATTHEWS N, SCHLESINGER BC, AKBAR AN, BACON PA, EMERY P, SALMON M. Active systemic lupus erythematosus is associated with the recruitment of naive/resting T cells, 226
- GORDON C, STOLL T, RICHARDSON K, BACON PA, ISENBERG DA. Quality of life is influenced by disease activity but is impaired in all SLE patients, (A1) 31
- GORDON C, STOLL T, RICHARDSON K, MALIK J, BACON PA, ISENBERG DA. Comparison of the SF-20 and SF-36 in the assessment of quality of life in SLE, (A1) 31
- GORDON MM, BLYTH T, HUNTER JA. Comparison of glenohumeral injections of triamcinolone hexacetonide in the painful rheumatic shoulder, (A1) 132
- GORMAN ST *see* Plant MJ
- GOTOH M *see* Matsuda J
- GOUGH A. The 1994-5 Michael Mason Travelling Fellowship: goodbye the Commonwealth, hello Australia, 1324
- GOUGH A *see also* Devlin J; Prouse P; Pugh MT; Talamo J
- GOULD DJ, FOXWELL BMJ, VITTEA E, FELDMANN M, CHERNAJOVSKY Y. Synthesis of an active cytotoxin by the cross-linking of human TNF α and ricin A-chain, (A1) 3
- GOULDING NJ *see* Biagini P; Perretti M; Pipitone N
- GOUPILLE P *see* Combe B
- Gout
chronic primary, comparative study of the efficacy of benzbromarone and allopurinol for the therapy of hyperuricemia, (A1) 145
and CPPD crystal arthropathy diagnosis, (E) 306
treatment following cardiac transplantation, (L) 1329
- GOW P *see* Emery D
- Gower's sign in children with juvenile chronic arthritis, (A1) 79
- GRABOWSKI PS, MACPHERSON H, RALSTON SH. Nitric oxide production in cells derived from the human joint, 207
- GRABOWSKI PS, WRIGHT P, RALSTON SH. Immunolocalisation of inducible nitric oxide synthase in rheumatoid synovium, (A1) 177
- Graft-versus-host disease and chronic polyarthritis, (L) 297
- GRAHAM DR *see* Dawson JK
- GRAHAME R, WEST J. Role of the rheumatology nurse practitioner in primary care: an experiment in the further education of the practice nurse, 581
- GRAHAME R *see also* Mishra MB
- Gram-negative bacteria, antibodies to four in rheumatoid arthritis which share sequences with the rheumatoid arthritis susceptibility motif, 592
- GRAN JT *see* Myklebust G
- GRAÑA J, FREIRE M, HERNÁNDEZ A, REINO S, MÉNDEZ MJ, FERNÁNDEZ-COUTO D, MARTÍNEZ A, DE LA FUENTE R, GLEZ-GAY MA, ASPE B, MADRIGAL MJ. Magnetic resonance imaging shows brain lesions in patients with Behçet's disease without central nervous system symptoms, (A1) 52
- GRANFORS K *see* Mäki-Ikola O
- GRANINGER W *see* Templ E
- GRANT M *see* Hamilton J
- GRATTAN C *see* Jolliffe VA
- GRAU JM *see* Font C
- GRAU R *see* González M; López-Longo FJ
- GRAY DH *see* Emery D
- GRAZIO S *see* O'Neill TW
- GRAVES M *see* Hill MB
- GREEN M, MASON JC, ARNOLD MM, GRIFFITHS IGH, McKENNA F. Oesophagitis from NSAIDs is not caused by pill oesophagitis, (A1) 225
- GREENWOOD B *see* Brown MA
- GRENNAN DM *see* Sanders PA
- GRENNAN S *see* Montague S
- GRIBNAU FWJ *see* Creemers MCW
- GRIEP EN *see* Bruyn GAW
- GRIFFITH J *see* Pincus T
- GRIFFITH SM *see* Dacre JA
- GRIFFITHS B, SITUNAYAKE RD, SALMON M, CLARK B, GOOI HC, EMERY P. Matched cross-sectional study of HLA DR types, rheumatoid factor and X-ray damage in North Indians and Pakistanis compared with Northern European RA patients, (A2) 11
- GRIFFITHS G *see* Arden NK
- GRIFFITHS GO, KEEN RW, HART DJ, LANCHBURY JS, DOYLE DV, SPECTOR TD. Polymorphisms of the vitamin D receptor and osteoarthritis, (A1) 117
- GRIFFITHS GO *see also* Arden NK; Keen RW
- GRIFFITHS H *see* Plant MJ
- GRIFFITHS I *see* Symmons DPM
- GRIFFITHS ID *see* Davidson BKS
- GRIFFITHS IGH *see* Green M; Mason JC
- GRIFFITHS MM *see* Terato K
- GRIGGS DC *see* Artlett CM
- GRISAR study
assessment of side effects, withdrawals and survival on treatment after 24 months, (A1) 215
is the control of disease progression within our grasp?, (S2)8
is cyclosporin a disease-controlling antirheumatic therapy? review of the two-year results, (A1) 215
- GROBBEE DE *see* Odding E
- GroEL epitope recognised by a synovial fluid T cell clone, mapping from a boy with juvenile chronic arthritis, (A1) 162
- GROOTVELD MC *see* Naughton DP
- GROSS WL *see* Johnston TW; Stoffel MP
- GROSSE R *see* Pugh MT
- GROSSMAN AB *see* Kassimos D
- GROTHE H *see* Mau W
- GROULT N *see* Chevalier X
- Growth hormone, nocturnal pulsatility and sleep study in patients with fibromyalgia, (A1) 138
- Gruppo Rheumatologi Italiani Studio Artrite Reumatoide *see* Ferraccioli GF
- GUANG LS. ANCA in RA patients, (L) 1328
- GUASCH I *see* Olivé A
- GUEDES C *see* Saroux A
- GUERIN B *see* Jorgensen C
- GUERIN J, CASEY E, FEIGHERY C, SIM B, JACKSON J. Antibodies to beta-2-glycoprotein 1 recognise a conformational epitope and are highly specific for the antiphospholipid syndrome, (A1) 21
- GUIALIS A *see* Vlachoyiannopoulos PG
- GUILLEMIN F *see* Smedstad LM
- GUILLEVIN L, LHOE F, AMOUREUX J, GHERARDI R, CALLARD P, CASASSUS P. Antineutrophil cytoplasmic antibodies, abnormal angiograms and pathological findings in polyarteritis nodosa and Churg-Strauss syndrome: indicators for the classification of vasculitides of the polyarteritis nodosa group, 958
- GÜL A, ÖZBEK U, ÖZTÜRK C, İNANÇ M, KÖNİÇ M, ÖZÇELİK T. Coagulation factor V gene mutation increases the risk of venous thrombosis in Behçet's disease, 1178

- GUNDEL RH *see* Chapman PT
 GUTIERREZ R *see* Hdez-Berian JA
 GUTIERREZ S *see* Migúlez R
 GUTIERREZ S, PALACIOS I, SÁNCHEZ-PERNAUTE O, HERNÁNDEZ P, EGIDO J, HERRERO-BEAMONT G. SÁME restores the changes in proliferation and in the synthesis of fibronectin and proteoglycans induced by TNF α on cultured synovial cells, (A1) 63
 GUTIERREZ S *see also* Palacios I
 GUTZKI F-M *see* Stichtenoth DO
- HÄBERLE J *see* Braun J
 HACKETT J, JOHNSON B, PARKIN A, SOUTHWOOD T. Physiotherapy and occupational therapy for juvenile chronic arthritis: custom and practice in five centres in the UK, USA and Canada, 695
 HAENEY MR *see* Snowden N
 HAGA H-J, JONSSON R. Rheumatology and immunology in Bergen—from basic science to clinical application, 1159
 HAIGH RC, MACK CS, KOH W-H, DUNPHY J, DIXEY J, CALIN A, MCHUGH NJ, MADDISON PJ. Atypical antineutrophil cytoplasmic antibodies (ANCA) in patients with ankylosing spondylitis and psoriasis, (A1) 161
 HAIGH RC, TAYLOR AL, CALIN A. Assessment of disease activity measures in ankylosing spondylitis: their relationship with acute phase reactants, (A1) 153
 HAJAL-HASSOUNI N *see* Cinal A
 HAIM T *see* Cordonnier C
 HAIM T *see de* Bandt M
 HAJEER A, JOHN S, OLLIER W, SILMAN A, DAWES P, HASSELL A, MATHEY D, FRYER A, STRANGE R, WORTHINGTON J. Association of extended TNF-HLA haplotypes with rheumatoid arthritis in males and females, (A1) 189
 HAJEER A *see also* Marlow A; Snowden N
 HAJEER AH, WORTHINGTON J, NAVES M, DAVIES EJ, HILLARBY MC, POULTON K, OLLIER WER. TNF A2 B3 and D2 microsatellite allele frequencies are increased in SLE and correlate with clinical outcome, (A1) 25
 HAJEER AH *see also* John SL
 HALIM M *see* El Ramahi KM
 HALL C *see* Smith G
 HALL FC, CAMILLERI J, WEEKS D, AMOS N, DARKE C, GIBSON K, PILE K, JESSOP J, WORDSWORTH BP. Relationships of HLA-DRB1 genotypes to susceptibility and severity in rheumatoid arthritis, (A1) 186
 HALL GM, LARBRE JP, SPECTOR TD, PERRY LA, DA SILVA JAP. Randomized trial of testosterone therapy in males with rheumatoid arthritis, 568
 HALL J *see* Avers HL
 HALL K *see* Dyce C
 HALL M *see* Yanni G
 HALL MA *see* Allen RL; Edmonds SE
 HALL N *see* Donnelly S; Foey A; Mack CS
 HALL ND *see* Balsa A; Flora RS
 HALLETT MB *see* Lloyds D
 HALSEY J, DODDS W, MCCALL E, MACKENZIE A. Rheumatology initiative to reduce orthopaedic waiting lists, (A1) 129
 HAMBLIN JJ *see* Ismail AA
 HAMEED K, BOWMAN S, KONDEATIS E, VAUGHAN R, LANCHBURY J, GIBSON T. Frequency of HLA class II antigens amongst Pakistani patients with rheumatoid arthritis, (A1) 185
 HAMEED K, GIBSON T. Comparison of the clinical features of hospital out-patients with rheumatoid disease and osteoarthritis in Pakistan and England, 994
 HAMEED K *see also* Gibson T
 HAMILTON I, LORY C, SHARMA B, RYAN PJ. Detection of active rheumatoid arthritis by 99m Tc human immunoglobulin (IgG), (A2) 15
 HAMILTON J, BRYDSON G, RAFFERTY D, BELL F, GRANT M, FRASER SM. Effect of second line anti-rheumatic therapy on gait characteristics in early rheumatoid arthritis, (A1) 212
 HAMMAM W *see* Dolan AL
 HAMURYUDAN V *see* Şahin (Özgün) Ş; Yazici H
 HAMZAOU A, HAMZAOU K, CHABBOU A, AYED K. Endothelin-1 expression in serum and bronchoalveolar lavage from patients with active Behçet's disease, 357
 HAMZAOU K *see* Hamzaoui A
- Hand
 bone mass, clinical associations of dual-energy X-ray absorptiometry measurement in rheumatoid arthritis, 1256
 bone mineral density, cross-sectional area of forearm musculature and grip strength in RA, (A1) 205
 MRI changes in early rheumatoid arthritis, (A1) 118
 osteoarthritis in early populations, 1291
 radiology in psoriatic arthritis: a five year follow-up study, (A1) 158
 video image analysis: development of an anatomic index as a potential outcome measure in rheumatoid arthritis, 1274
- Handicap
 in inflammatory arthritis, 891
 patient-perceived, development of a clinical tool to measure: patient-centred approach to evaluation and treatment in rheumatoid arthritis, 921
 HANFY M *see* Atia G
 HANSEN BU *see* Berglund J
 HANSEN M *see* Johansen JS; Østergaard M
 HARBUS MS, PERVEEN-GILL Z, LALIES MD, JESSOP DS, LIGHTMAN SL, CHOWDREY HS. Role of endogenous serotonin in adjuvant-induced arthritis in the rat, 112
 HARDINGHAM T *see* Venn G
 HARDINGHAM TE *see* Dudhia J
 HARDWICK L, WILLIAMS C, BUTCHER S, CONSIDINE E, NICOD A, WALSH S, PROCKOP D, REGINATO A, CAEIRO F, BRANCOLINI V, CARR A, LATHROP M, WORDSWORTH P. Genetic linkage of chondrocalcinosis to chromosome 5p15.1-15.2, (A1) 148
 HARDWICK LJ, WALSH S, BUTCHER S, NICOD A, SHATFORD J, BELL J, LATHROP M, WORDSWORTH BPW. Mapping and characterisation of the genes involved in rheumatoid arthritis, (A2) 10
 HARMER IJ, MAGEED RA, BRÜGGEMANN M, MACKWORTH-YOUNG CG. Single anti-DNA-related V_H gene encodes for multiple specificities: the influence of the light chain, (A1) 23
 HARRISON AA *see* Chapman PT
 HARRISON B *see* Brennan P
 HARRISON BDW *see* Reid AJC
 HARRISON BJ, BRENNAN P, SILMAN AJ, BANKHEAD CR, BARRETT EM, SYMMONS DPM. Synovitis is associated with future radiological erosions in individual joints of the hands and feet of patients with early RA, (A1) 229
 HARRISON BJ, SILMAN AJ, SYMMONS DPM. ARA 1987 criteria do not perform well in early RA, (A2) 18
 HARRISON BJ, SYMMONS DPM, BRENNAN P, BARRETT EM, SILMAN AJ. Natural remission in inflammatory polyarthritis: issues of definition and prediction, 1096
 HARRISON BJ *see also* Symmons DPM
 HARRISON R *see* Jawed S
 HART D *see* Cooper C
 HART DJ *see* Arden NK; Griffiths GO; Nandra D; Spector TD
 HART K *see* Leak AM
 HARVEY G *see* McHugh NJ
 HARVEY GR, RANDS A, MADDISON PJ, MCHUGH NJ. Antibodies to RNA polymerases in systemic sclerosis: clinical and serological associations, (A1) 46
 HARWOOD RH, CARR AJ, THOMPSON PW, EBRAHIM S. Handicap in inflammatory arthritis, 891
 HASKARD DO *see* Chapman PT; Mason JC
 HASLOCK I. Quality of care and patient satisfaction, 382
 HASLOCK I *see also* Speed CA
 HASSALL A *see* Stanworth SJ
 HASSAN LA *see* Acebes JC
 HASSAN WU *see* Yeap SS
 HASSELL A, DAWES P. Survey of attitudes and current practices regarding rheumatology trainees among UK rheumatologists, (A1) 131
 HASSELL A *see also* Birrell FN
 HASSELL P *see* Hajeer A
 HAWKINS CP *see* Salih AM
 HAWKINS PN, PERSEY MR, HEATON N, WILLIAMS R, YACOB M, WINEARLS CG, FEEST TG, HOLMGREN G, PEPYS MB. New roles for organ transplantation in amyloidosis, (A1) 12
 HAWKINS PN *see also* Persey MR
 HAWORTH S, RIDGEWAY J, STEWART I, DYER PA, PEPPER L, OLLIER W. Polymyalgia rheumatica is associated with both HLA-DRB1*0401 and DRB1*0404, 632
 HAY FC *see* Dalziel M; Laversuch CJ
 Hay fever, low prevalence in patients with rheumatoid arthritis, (A1) 197
 HAZES JMW *see* Simon CH; Vliet Vlieland TPM
 HAZLEMAN BL. Rheumatic disorders of the eye and the various structures involved, 258
 HAZLEMAN BL *see also* Isaacs JD; Sharma SD; Summers CA
 HDEZ-BERIAIN JA, DIAZ-FLOREZ L, GUTIERREZ R, GONZÁLEZ T, SEGURA-GARCÍA C, GANTES M, BUSTABAD S, RUEZ-LOZANO B. Response of cartilage and perichondrium implanted in the joint, (A1) 240
 HDEZ-BERIAIN JA, SEGURA GARCÍA C, NODA B, MARLES M, CHERREZ C. Incidence and severity of NSAID-induced gastrointestinal bleeding, (A1) 220
 Health Assessment Questionnaire, Spanish, comparative study of the short vs Stanford versions. Correlation between Likert and VAS scales for pain and health status, (A1) 232
 Health economics as an aspect of health outcome: basic principles and application in rheumatoid arthritis, (L) 494
 Hearing loss study in Paget's disease of bone. High frequency audiometry evaluation, (A1) 92

- HEATH P *see* Salih AM
 HEATON N *see* Hawkins PN
 HECKER H *see* Mau W
 HEINEGÅRD D *see* McAlinden A
 HELDERS PJM *see* van der Net J
 HELIÖVAARA M, AHO K, KNEKT P, REUNANEN A, AROMAA A. Serum cholesterol and risk of rheumatoid arthritis in a cohort of 52 800 men and women, 255
 HELLIO LE GRAVERAND MP, TRON AM, ICHOU M, DALLARD MC, RICHARD M, UEBELHART D, VIGNON E. Assessment of urinary hydroxypyridinium cross-links measurement in osteoarthritis, 1091
 HELLIWELL PS
 Audit of shared care between hospital and general practice in the management of rheumatoid arthritis, (L) 398
 Combination therapy with sulphasalazine and azathioprine, (L) 493
 Comparison of a community clinic with a hospital out-patient clinic in rheumatology, 385
 Diagnostic criteria for work-related upper limb disorders, (E) 1195
 Serial measurement of joint stiffness following intra-articular steroid: an early increase may indicate sub-clinical crystal synovitis, (A1) 211
 HELLIWELL PS *see also* Woodburn J
 HENSEN MMA *see* Verschure PJ
 HENKLEIN P *see* Riemekasten G
 Henoch-Schönlein purpura in adulthood: a more severe syndrome with a good outcome, (A1) 55
 HENRIKSEN O *see* Østergaard M
 Heparin
 low molecular weight, pilot study of treatment in Raynaud's phenomenon and scleroderma, (A1) 43
 and rheumatoid arthritis, (L) 808
 Hepatitis C virus
 associated with mixed cryoglobulinaemia, broncheolar lavage, 978
 chronic infection, presence of rheumatic diseases, (A1) 168
 infection, rheumatic manifestations, (A1) 168
 infection and liver involvement in patients with primary Sjögren's syndrome, (A1) 15
 and mixed cryoglobulinaemia genotypes, (A1) 50
 positive patients with sicca syndrome and primary Sjögren's syndrome, immunological features, (A1) 15
 Hepatitis lupoid or lupus with mesenteric vasculitis, 1309
 Herbal medicine, proprietary, effect on the relief of chronic arthritic pain: a double-blind study, 874
 HERBERT KE *see* Hill CM
 HERMAN C, CHERNAJOVSKY Y. Mutational analysis of the membrane proximal cleavage site of human p75 tumour necrosis factor receptor, (A1) 109
 HERMAN C *see also* Chernajovsky Y
 HERNÁNDEZ A *see* Graña J
 HERNÁNDEZ P *see* Gutierrez S; Palacios I; Sánchez-Pernaute O
 HERNÁNDEZ-MUNOZ H *see* Leirisalo-Repo M
 HERNANZ A *see* De Miguel E; Rodríguez R
 HERRERO-BEAMONT G *see* Gutierrez S; Miguélez R; Palacios I; Sánchez-Pernaute O
 HERRICK A *see* Blann AD; Bukhari M; Chaloner C
 HERRICK AL *see* Anderson ME; Bukhari M; Elliott M; Jayanetti S; Teh LS
 HESLOP P *see* Speed CA; Walker DJ
 HESLOP PS *see* Chatterjee S; Kay LJ; Walker DJ
 Heterotopic ossification complicating long-term sedation, (L) 700
 HEWLETT S, KIRWAN JR. Dissatisfaction with disability: repeated associations with mood and beliefs about function, (A1) 196
 HICKLING P, JACOBY R, KIRWAN J. Joint destruction recommences after glucocorticoids are withdrawn in early rheumatoid arthritis, (A2) 22
 HICKLING P *see also* Suzuki H
 HIEPE F *see* Riemekasten G
 HIETARINTA M. Antinuclear and antinucleolar antibodies in patients with scleroderma-polymyositis overlap syndrome, (L) 1326
 HIGHTON J, SOLOMON C, GARDINER DM, DOYLE TCA. Video image analysis of hands: development of an anatomic index as a potential outcome measure in rheumatoid arthritis, 1274
 HILL C, ROMAS E, KIRKHAM B. Use of sequential DTPA clearance and high resolution computerized tomography in monitoring interstitial lung disease in dermatomyositis, 164
 HILL CM, LUNEC J, HERBERT KE. Release of tumour necrosis factor α from human monocytes: a common mechanism of lupus-inducing drug immunomodulation, (A1) 3
 HILL MB, PHIPPS JL, MILFORD-WARD A, GREAVES M, HUGHES P. Further characterization of anti-endothelial cell antibodies in systemic lupus erythematosus by controlled immunoblotting, 1231
 HILLARBY C *see* Sanders PA
 HILLARBY MC *see* Hajeer AH
 HILLIQUIN P, RENOUX M, PERROT S, PUÉCHAL X, MENKES CJ. Occurrence of pulmonary complications during methotrexate therapy in rheumatoid arthritis, 441
 HILLMAN M *see* Tennant A
 HILLS BA *see* Schwarz IM
 HILSON AJW *see* MacGregor AJ
 Hip
 fractures, discrimination by peripheral quantitative computed tomography (pQCT), (A2) 6
 and knee, association of abnormalities on physical examination with locomotor disability in the Rotterdam study, 884
 pain, patients presenting with in the community: comparing those with and without radiographic changes of osteoarthritis, (A1) 102
 and wrist fracture incidence reduction of an Asian population in Leicester, (A2) 3
 Histiophagocytic syndrome presenting as juvenile chronic arthritis, (A2) 37
 HIV infection
 clubbing, 292
 musculoskeletal manifestations, (A1) 167
 osteoarticular tuberculosis in two different populations, (A1) 166
 in a patient with systemic lupus erythematosus, (A2) 35
 polyarthritis nodosa in: report of four cases and review of the literature, 796
 HIV-infected patient, bone bacillary angiomatosis, 901
 HIZLI N *see* Yazici H
 HLA association
 and disease expression of RA: comparison of Egypt and the UK, (A1) 185
 with toxicity in patients with rheumatoid arthritis receiving gold sodium thiomalate in a Spanish population, (A1) 211
 HLA B27, spondylarthritis in the Gambia—evidence for a non-HLA B27 protective effect, (A1) 113
 HLA class I antigens in serum and synovial fluid from patients with rheumatoid arthritis and other arthropathies, (A1) 176
 HLA class II antigens
 frequency amongst Pakistani patients with rheumatoid arthritis, (A1) 185
 and synovial fluid cytology in rheumatoid arthritis, (A2) 12
 HLA DR4, hormonal associations with rheumatoid arthritis, (A1) 119
 HLA DR types, rheumatoid factor and X-ray damage, matched cross-sectional study in North Indians and Pakistanis compared with Northern European RA patients, (A2) 11
 HLA-B27 typing in the diagnosis of early, oligosymptomatic spondylarthropathies, (C) 192
 HLA-B27-restricted peptide presentation to cytotoxic T cells in reactive arthritis, (A1) 114
 HLA-DQB1 polymorphism and antibodies to RNA polymerases in systemic sclerosis, (A1) 45
 HLA-DRB*0401 association with polymyalgia rheumatica, 632
 HLA-DRB1
 and complement C4 null alleles in systemic lupus erythematosus in a Spanish population, (A1) 26
 genotype relationship to susceptibility and severity in rheumatoid arthritis, (A1) 185
 genotype role in the susceptibility to and severity of inflammatory polyarthritis, (A1) 186
 Ho C *see* Mok CC
 HO M, LAVERY B, PULLAR T. Perception by patients with rheumatoid arthritis of the risk benefit ratio of drug treatment, (A1) 212
 HODGKINS PR *see* Vakalis AN
 HOEFAKKER S *see* Dolhain RJEM
 HOFFMEYER F, WITTE K, SCHMIDT RE. Survival and functional properties of INF γ -treated PMN, (A1) 61
 HOFMAN A *see* Odding E
 HOGARTH MB, MORLEY BJ, ALLEN PJ, SLINGSBY JH, DAVIES KA, WALPORT MJ. Effect of dominant disease susceptibility genes on phenotype in BXS lupus-prone mice, (A1) 112
 HOGG J *see* Smith RW
 HOLDER R *see* Devlin J
 HOLLINGWORTH P. Back pain in children, 1022
 HOLLIS S *see* Anderson ME
 HOLLY JMP *see* Fernihough JK
 HOLMGREN G *see* Hawkins PN
 HOLT PJL, BALDAM E, LENDON M, KEEN J. Histiophagocytic syndrome presenting as juvenile chronic arthritis, (A2) 37
 HOLT PJL *see also* Baldam EM; Stathopoulou E
 HOLT PLH *see* Donn RP
 HOMAEI H, ALAVI A, AXFORD JS. Rheumatoid arthritis associated glycosylation changes may not be idiosyncratic to IgG, (A1) 174
 Homeopathy, is it effective in rheumatoid arthritis?, (A1) 227
 HOPKINS A *see* Houssien DA
 HORDON LD. Massive pericardial effusion in scleroderma, (L) 807
 Hormonal associations with rheumatoid arthritis due to HLA DR4, (A1) 119
 Hormone replacement therapy, is it protective for osteoarthritis in women: the Chingford Study, (A1) 124
 HORNEBECK W *see* Chevalier X

- HORSFALL AC, LI J-M, FAN W, MAINI RN, VENABLES PJW. Autoimmunisation to an endogenous retroviral envelope protein during pregnancy—an autoantigen in congenital heart block, (A1) 112
- HORST G *see* Spronk PE
- HOSIE G *see* Nahit ES
- HOSIE J, DISTEL M, BLUHMKI E. Meloxicam in osteoarthritis: a six-month, double-blind comparison with diclofenac sodium, (S1) 39
- HOSKING DJ *see* Patel S; Yeap SS
- Hospital out-patients, comparison of the clinical features with rheumatoid disease and osteoarthritis in Pakistan and England, 994
- HOUGH D *see* Mack CS
- HOUSSEIN DA, BERRY H, MCKENNA S, SCOTT DL. Measuring disease activity and outcome in rheumatoid arthritis using the Nottingham Health Profile, (A1) 231
- HOUSSEIN DA, LEFEBVRE C, DEPRESSEUX G, LAMBERT M, DEVOGELAER J-P, NAGANT DE DEUXCHAISNES C. Trabecular and cortical bone loss in systemic lupus erythematosus, 244
- HOUSSEIN D, VINCENT C, SERRE G, SCOTT D. Anti-keratin antibodies, disease activity and radiological damage in rheumatoid arthritis, (A2) 14
- HOUSSEIN DA, CHOY EHS, BERRY H, SCOTT DL. Differences between the clinical and radiological progression of rheumatoid arthritis, (A1) 199
- HOUSSEIN DA, COMER M, DOYLE DV, HOPKINS A, SCOTT DL. Early referral and outcome in rheumatoid arthritis, (A1) 229
- HOUSSEIN DA, JONSSON T, SCOTT DL
Rheumatoid factor isotypes, an association with radiological progression in patients with RA, (A1) 207
Rheumatoid factor isotypes and radiological progression in rheumatoid arthritis, (A2) 14
- HOUSSEIN DA *see also* Scott DL
- HOUSTON T, MUNRO R, BROCK JH, STURROCK RD, FITZSIMONS EJ. Evidence for abnormal erythroblast iron metabolism in the chronic anaemia of rheumatoid arthritis, (A1) 191
- HOUTMAN PM. Folate supplementation and methotrexate, (L) 1031
- HOUTMAN PM *see also* Riemsma RP
- HOWARD Y *see* Bryan C
- HOWAT S, BOLTON M, DAY A, BAYLISS M. TSG-6: a hyaluronan binding matrix protein upregulated by cytokines in joint disease, (A1) 236
- HOWELL K *see* Denton CP
- HOWELL WM *see* Wong WM
- HTLV-1, tax gene is not present in the salivary glands of patients with idiopathic Sjögren's syndrome, (A1) 16
- HU DE *see* Walsh DA
- HUANG FP *see* McInnes IB
- HUANG J-L. Multiple cryptococcal brain abscesses in systemic lupus erythematosus, (L) 1334
- HUGHES FJ *see* Sahinoglu T
- HUGHES G *see* Rudwaleit M
- HUGHES GRV *see* Amengual O; Ames PRJ; Atsumi T; Etherington J; Khamashta MA; Méndez MJ; Ruiz-Irastorza G
- HUGHES P *see* Hill MB
- HUGHES RA *see* Stagg AJ
- Hughes syndrome: a common problem in Kuwait hospitals, 1132
- HUISOON A *see* Devlin J
- HUITEMA MG *see* Spronk PE
- HUKUDA S *see* Tiwana H
- HULL RG *see* Sinclair D
- HULL RJ *see* Vakalis AN
- HUMPHRIES J. Search for infective agents in undifferentiated oligoarthritis in Papua New Guinea, (L) 492
- HUMPHRIES M *see* Ng T
- HUNT BJ *see* Khamashta MA
- HUNT M *see* Paleolog E
- HUNTER JA, BLYTH TH. Inappropriately high ESR results—a feature of amyloid complicating rheumatic disorders?, (A1) 77
- HUNTER JA *see also* Gordon MM; Porter DR
- HUNTER M *see* Hurst NP
- HUNZIKER T *see* Lehmann T
- HUPPERTZ H-I, MICHELS H. Pattern of joint involvement in children with Lyme arthritis, 1016
- HURLEY MV, NG TS. Evaluation of reliability, reproducibility and validity of two methods of assessing proprioceptive acuity in the lower limb, (A1) 140
- HURLEY MV, REES J. Quadriceps strength, activation and functional performance in healthy young, middle-aged and elderly subjects, (A1) 140
- HURST NP, KIND P, RUTA D, HUNTER M, STUBBINGS A
Construct validity of EuroQol in 240 patients with rheumatoid arthritis stratified by functional class, (A2) 18
Quality of life in rheumatoid arthritis: EuroQol is as responsive as ACR disease activity core measures to self-reported change in RA, (A2) 17
Testing the validity of the MOS SF36 generic health profile in rheumatoid arthritis, (A2) 17
- HUSKISSON EC, GHOZLAN R, KURTHEN R, DEGNER FL, BLUHMKI E. Long-term study to evaluate safety and efficacy of meloxicam therapy in patients with rheumatoid arthritis, (S1) 29
- HUSSAIN MAS *see* Malaviya AN
- HUSSEY R *see* Tunn AJ
- HUTTON CW *see* Gandy S; Gasson J
- HVOLRIS J *see* Johansen JS
- Hyaluronan binding matrix protein TSG-6 upregulated by cytokines in joint disease, (A1) 236
- Hydrotherapy, acute physiological effect in patients with ankylosing spondylitis, (A1) 156
- Hydroxyl and/or ferryl radicals at sites of inflammation, indirect evidence to support production, (A2) 13
- Hydroxypropylidinium cross-links measurement in osteoarthritis, 1091
- Hyper-IgM syndrome associated with rheumatoid arthritis: report of RA in a patient with primary impaired CD40 pathway, 282
- Hypermobility Special Interest Group report, British Society for Rheumatology meeting in Glasgow, April 1995, 392
- Hyperparathyroidism, tertiary after long-term phosphate supplementation in adult-onset hypophosphataemic osteomalacia, 598
- Hyperuricaemia
changes in apolipoproteins levels following low-calorie dieting, (A1) 145
comparative study of the efficacy of benzbromarone and allopurinol for therapy in chronic primary gout, (A1) 145
risk factors in Catalonia-II survey, (A1) 146
- Hypokaemic paralysis, thyrotoxicosis and renal tubular acidosis presenting as, 289
- Hypophosphataemia, chronic: possible osteoporosis risk-factor, (A1) 96
- IANNACCONE L *see* Ames PRJ
- IANNONE F, CAULI A, YANNI G, KINGSLEY GH, ISENBERG DA, CORRIGALL V, PANAYI GS. T-lymphocyte immunophenotyping in polymyositis and dermatomyositis, 839
- IANNONE F. CD69 on synovial T cells in rheumatoid arthritis correlates with disease activity, (L) 397
- ICHIKAWA K *see* Atsumi T
- ICHOU M *see* Helio le Graverand MP
- Idiopathic carpal tunnel syndrome management: survey of rheumatologists, (A2) 34
- IgA1 and IgA2 subclass antibody, similarly increased serum levels against *Klebsiella pneumoniae* bacteria in ankylosing spondylitis patients with/without extra-articular features, 125
- IgA
and α 1-antitrypsin, normal urine provides an environment that augments complex formation between, (A2) 13
immunoglobulins in rheumatoid arthritis and sacroiliitis, (A1) 201
- IGF system disruption in osteoarthritic and rheumatoid patients compared to non-arthritic volunteers, (A1) 105
- IgG
anti-DNA antibody, sequence analysis and pathogenic properties, (A1) 111
Fc γ IIA polymorphism in SLE: genotypic analysis in Caucasian, Chinese and Afro-Caribbean patients, (A1) 27
glycosylation changes during fasting in patients with rheumatoid arthritis, 117
rheumatoid arthritis associated glycosylation changes may not be idiosyncratic to, (A1) 174
sialylation in rheumatoid arthritis in relation to galactosylation and rheumatoid factor expression, (A1) 175
- IgM
anti-U1RNP antibodies, predominance in patients with systemic lupus erythematosus, 534
rheumatoid factor, interference by on ELISA for antibodies to mycobacterial 65 kDa heat shock protein in rheumatoid arthritis, (L) 300
- IIDA M *see* Kobayashi H
- IL-1 antagonist, novel, double blind study assessing safety and efficacy of IL-1Ra for treatment of rheumatoid arthritis, (A1) 214
- IL-1 α , TGF β and retinoic acid effects on homeostasis of aggrecan and biglycan from articular cartilage, (A1) 7
- IL-4, can prevent the release of collagen from cartilage stimulated with proinflammatory cytokines *in vitro*, (A1) 8
- IL-6
5' regulatory region gene polymorphisms in patients with systemic onset JCA and normal control subjects, (A1) 79
may be responsible for increased bone turnover in polymyalgia rheumatica, (A2) 4
receptor, serum and synovial fluid in patients with different arthropathies, (A1) 9
- IL-8 and neutrophil function of synovial fluid in rheumatoid arthritis and seronegative polyarthritis, 1244
- IL-10, T cell clones from a subset of patients with primary Sjögren's syndrome produce high levels, (A1) 16

- IL-12
antibodies to, modification of murine collagen-induced arthritis, (A1) 108
expression and quantitation by competitive PCR in the synovium of patients with rheumatoid arthritis and osteoarthritis, (A1) 180
- IL-15 expression in rheumatoid arthritis, (A1) 180
- Iliopsoas bursitis, an unusual presentation of metastatic bone disease, 285
- Illness perception vs disease activity as predictors of health status and psychological well-being in patients with systemic lupus erythematosus: longitudinal analysis, (A1) 34
- Iloprost
improved outcome in scleroderma renal crisis: association with anti RNA polymerase, (A1) 41
intermittent infusion therapy of pulmonary hypertension in scleroderma—a pilot study, (L) 1187
- IM HOF V see Beyeler C
- Imaging inflammation in rheumatoid arthritis: comparisons between ¹¹¹In-anti-E-selectin MAB 1.2B6 and ^{99m}Tc- and ¹¹¹In-immunoglobulin, (A1) 203
- IMHOF H see Kainberger F
- Immune complexes, complement and systemic lupus erythematosus, 5
- Immune response to the degraded and nondegraded (native) Sa antigens, dissociation in rheumatoid arthritis and systemic lupus erythematosus, (A1) 18
- Immunoglobulin
intravenous, failure of first-line therapy in a child with scleromyositis, (L) 1029
intravenous as sole therapy for systemic vasculitis, 1150
lambda light chains, somatic mutation and gene fingerprinting in the synovium of patients with rheumatoid arthritis, (A1) 172
- Immunoglobulin-Tc99 scan in the assessment of joint inflammation in rheumatoid arthritis, (A1) 203
- Immunology and rheumatology in Bergen—from basic science to clinical application, 1159
- Immunosuppressive therapy
does exposure increase the 10 year malignancy and mortality risks in rheumatoid arthritis? A matched cohort study, 738
does exposure increase the 10-year malignancy and mortality risks in rheumatoid arthritis? a matched cohort study, (A1) 227
- IMURA S see Wada M
- INANÇ M see Gül A
- Index, simple of range of movement of large joints as an outcome measure in RA, (A1) 194
- Indomethacin and morphine, synergistic interaction between attenuates pain sensation and tissue destruction in adjuvant arthritis, (A1) 75
- Infective agents in undifferentiated oligoarthritis, search in Papua New Guinea, (L) 492
- Infertility may sometimes be associated with NSAID consumption, 76
- Inflammation
imaging in rheumatoid arthritis: comparisons between ¹¹¹In-anti-E-selectin MAB 1.2B6 and ^{99m}Tc- and ¹¹¹In-immunoglobulin, (A1) 203
in the joint, why does it hurt?, (E) 405
urate crystal-induced, endothelial activation: *in vitro* and *in vivo* studies on the role of tumour necrosis factor, (A1) 146
- Inflammatory polyarthritis, natural remission in: issues of definition and prediction, 1096
- Inflammatory response, intra-articular, silica glass fragments from steroid containing ampoules augment, (A1) 121
- Informal care for patients with rheumatoid arthritis, (A1) 127
- Information leaflets, randomised controlled study for people with rheumatoid arthritis, (A2) 32
- INGELMO M see García-Carrasco M; Navarro M
- Insulin-like growth factor-1
binding protein 3 role in human articular cartilage explants, 515
receptor localization in human normal and osteoarthritic cartilage in relation to proteoglycan synthesis and content, 1044
- Integrin-induced T-cell signaling in systemic lupus erythematosus: a novel tool for disease monitoring, (A1) 38
- Interferon alpha, resolving familial Mediterranean fever attacks, (L) 1188
- Interferon alpha-2b, severe osteoporosis due to systemic mast cell disease: successful treatment, 898
- Interferon-gamma
and ciprofloxacin, effect on the infection of *Chlamydia trachomatis* in the monocytic cell line THP-1, (A1) 169
together with IFN-gamma receptor, increased expression in the rheumatoid synovial membrane compared with synovium of patients with osteoarthritis, 24
- Intestinal microflora, increased bacterial urease activity in faeces in juvenile chronic arthritis, 689
- Intestinal permeability, high in patients with inflammatory arthritis is more strongly associated with NSAID therapy than with other factors, (A1) 221
- Intra-articular pressure
dynamics in inflammatory arthropathies, (A1) 71
measurement in children with juvenile chronic arthritis, (A1) 78
- IRVINE AE see Bell AL
- ISAACS JD, MANNA VK, RAFSON N, BULPITT KJ, HAZLEMAN BL, MATTESON EL, ST. CLAIR EW, SCHNITZER TJ, JOHNSTON JM. CAMPATH-1H in rheumatoid arthritis—an intravenous dose-ranging study, 231
- ISASI C, PEREZ A, TABUENCA MJ, DIEZ ILE, LARREA A, ORTIZ J. Scintigraphic uptake of metabolic activity tracers in Paget's disease of bone, (A1) 94
- ISASI C see also Perez A
- ISENBERG D see Bodman-Smith K; Pincus T
- ISENBERG DA, MAINI RN. Consequences of the Calman Report on academic rheumatology, (E) 608
- ISENBERG DA see also Choy EHS; Cohen MR; Davies KA; Devi B; Fox RA; Garton MJ; Gordon C; Iannone F; McLure CE; Moore PM; Newkirk MM; Öcal L; Ravirajan CT; Sharma A; Shortall E; Stoll T
- ISMAIL AA, DASGUPTA B, TANQUERAY AB, HAMBLIN JJ. Ultrasonographic features of diabetic cheiroarthropathy, 676
- ISOLAURI E see Malin M
- ISOMÄKI P, SÖDERSTRÖM K-O, PUNNONEN J, ROIVAINEN A, LUUKKAINEN R, MERILAHTI-PALO R, NIKKARI S, LASSILA O, TOIVANEN P. Expression of bcl-2 in rheumatoid arthritis, 611
- Isonitrite-Tc99 bone scan vs radiologic evaluation in the assessment of the malignant nature of a tumour-like bone lesion, (A1) 241
- Isotope bone scans for foot problems, (A2) 8
- Isotype determination of serum antibodies against *Chlamydia*, value for the diagnosis of *Chlamydia* reactive arthritis, 542
- ISSLER H see Slingsby JH
- ITONAGA I see Fujikawa Y
- IWAMOTO I see Furukawa M
- IWASAKI K see Okano K
- IWASHITA A see Kobayashi H
- IX207-887, double blind study assessing safety and efficacy of novel IL-1 antagonist for treatment of rheumatoid arthritis, (A1) 214
- JACKSON G see Mishra MB
- JACKSON J see Guerin J
- JACOBY R see Hickling P
- JACOBY RK see Gandy S; Gasson J; Mills SY
- JAFFEY MN see Vakalis AN
- JAHR O see Jevtic V
- JAMAR F see Chapman PT
- JAMES D see Prouse P; Talamo J
- JAMES DW. Accidental overdose of prednisolone and methotrexate, (L) 704
- JAMES IT, WALNE AJ, PERRETT D. Relationships between bone formation and degradation markers in serum and synovial fluid in rheumatoid patients, (A1) 91
- JAMES IT see also Walne AJ
- JARVIS L see Smith RW
- JAWAD ASM. Tumoral calcinosis: a case report, (A2) 35
- JAWAD ASM see also Saced IT
- JAWAHEER D see Brennan P
- JAWED S, STEVENS CR, ZHANG Z, FARR M, SAHINOGLU T, CHIKANZA I, HARRISON R, BLAKE DR. Plasma NADH oxidase activity in inflammatory synovitis measured by chemiluminescence, (A1) 69
- JAWED S see also Gaffney K; Masud T
- JAYANETTI S, SMITH C, MOORE T, JAYSON MIV, HERRICK AL. Thermal stress testing and video nail-fold capillaroscopy in children with Raynaud's phenomenon, (A2) 26
- JAYNE DRW, LOCKWOOD CM. Intravenous immunoglobulin as sole therapy for systemic vasculitis, 1150
- JAYSON MIV see Anderson ME; Blann AD; Bukhari M; Jayanetti S; Macfarlane GJ; Teh LS
- Jejunal lesions in ankylosing spondylitis are due to non-steroidal anti-inflammatory drug therapy, (A1) 157
- JEPSON A see Brown MA
- JESSOP DS see Harbuz MS
- JESSOP J see Hall FC
- JEVITIC V, WATT I, ROZMAN B, PRESETNIK M, LOGAR D, PRAPROTNIK S, TOMSIC M, SIPEK A, KOS-GOLJA M, SEPE A, JAHR O, DEMSAR F, MUSIKIC P, CAMPION G. Prognostic value of contrast enhanced Gd-DTPA MRI for development of bone erosive changes in rheumatoid arthritis, (S3) 27
- JIMENEZ M see De La Mata J
- JIMENEZ MD see Sanchez J
- JOBANPUTRA P, LIN H, BRENNAN F, PINDER S, BAVINGTON C, SALTER DM, NUKI G. Interactions of soluble fibronectin with isolated human articular chondrocytes, (A1) 62
- JOBANPUTRA P see also Bavington C; Brennan FR
- JOHANSEN JS, HVOLRIS J, HANSEN M, BACKER V, LORENZEN I, PRICE PA. Serum YKL-40 levels in healthy children and adults. Comparison with

- serum and synovial fluid levels of YKL-40 in patients with osteoarthritis or trauma of the knee joint, 553
- JOHANSON NA *see* Sakkas LI
- JOHN S, MYERSCOUGH A, MARLOW A, OLLIER W, SILMAN A, WORTHINGTON J. Linkage analysis studies of candidate genes on affected sib pair RA families, (A1) 190
- JOHN S *see also* Hajer A; Marlow A
- JOHN SL, HAJER AH, MARLOW A, OLLIER WER, SILMAN A, WORTHINGTON J. Linkage and association studies of the natural resistance associated macrophage protein locus in rheumatoid arthritis, (A1) 188
- JOHNS CW *see* Teh LS
- JOHNSON AS *see* Coombes GM
- JOHNSON B *see* Hackett J
- JOHNSON J, YANNI G, KINGSLEY GH. Nature of the synovial infiltrate in the synovial membrane in reactive arthritis, (A1) 163
- JOHNSON N *see* Newkirk MM
- JOHNSON U *see* Fex E
- JOHNSTON JM *see* Isaacs JD
- JOHNSTON R *see* Munro R
- JOHNSTON TW, SZYMOWIAK CH, CSERNOK E, GROSS WL. Expression of proteinase 3 (Wegener's autoantigen) in a baculovirus system, (A1) 51
- Joint
- aseptically failed prostheses, osteolytic properties of the synovial-like tissue, 943
 - assessment in rheumatoid arthritis, (S2) 14
 - benign hypermobility syndrome, extra-articular features, 861
 - destruction recommences after glucocorticoids are withdrawn in early rheumatoid arthritis, (A2) 22
 - human, nitric oxide production in cells derived from, 207
 - involvement in children with Lyme arthritis, 1016
 - involvement in connective tissue disease and its associations with the third hypervariable region (HVR3), (A1) 30
 - rheumatic disease, contrast-enhanced MRI, (S3) 44
 - rheumatoid, agalactosyl IgG in aggregates, 335
 - serial measurement of stiffness following intra-articular steroid: an early increase may indicate sub-clinical crystal synovitis, (A1) 211
 - why does an inflammation hurt?, (E) 405
- JOLLIFFE VA, WATTS RA, GRATTAN C, SCOTT DGI. Incidence and clinical associations of biopsy proven localised cutaneous vasculitis in the Norwich Health Authority, (A1) 54
- JOLLIFFE VA *see also* Cruwys SC
- JOLLY BC *see* Dacre JA
- JONAS D *see* Baerwald CGO
- JONES A *see* Lanyon P; Ledingham J; O'Reilly SC
- JONES AC *see* Yeap SS
- JONES AKP *see* Coombes GM
- JONES G, BROOKS PM. Injectable gold compounds: an overview, 1154
- JONES M, SYMMONS D, FINN J, WOLFE F. Does exposure to immunosuppressive therapy increase the 10 year malignancy and mortality risks in rheumatoid arthritis? A matched cohort study, 738
- JONES M *see also* Williamson L
- JONES MA, SYMMONS DPM, FINN JD, WOLFE F. Does exposure to immunosuppressive therapy increase the 10-year malignancy and mortality risks in rheumatoid arthritis? a matched cohort study, (A1) 227
- JONES PJ *see* Popert AJ
- JONES PW *see* Avers HL; Cheung NT; Plant MJ
- JONES S *see* Koh WH
- JONES SD, STEINER A, GARRETT SL, CALIN A. Bath ankylosing spondylitis patient global score (BAS-G), 66
- JONES SM, BALAKRISHNAN C, MCHUGH NJ. Progression of peripheral joint disease in psoriatic arthritis: a five year clinical follow-up study, (A1) 158
- JONES SM *see also* Balakrishnan C; Flora RS
- JONSSON K *see* Fex E
- JONSSON R *see* Haga H-J
- JONSSON T *see* Houssien DA
- JOOSTEN LAB *see* Verschure PJ
- JORDI B *see* Beyeler C
- JORGENSEN C, GUERIN B, FERRAZZI V, BOLOGNA C, SANY J. Arthritis associated with monoclonal gammopathy: clinical characteristics, 241
- JORGENSEN C *see also* Bologna C
- JOUSSEAUME C *see* Conrozier T
- JUANOLA X *see* Gomez-Vaquero C; Ros S; Valverde J
- JUBB RW *see* Maggs FM
- JULKUNEN H *see* Mack CS
- JURADO R *see* Gonzalez-Dominguez J
- Juvenile chronic arthritis
- 5' regulatory region polymorphisms in the interleukin-6 gene, (A1) 79
 - Gower's sign in children, (A1) 79
 - histiophagocytic syndrome presenting as, (A2) 37
 - increased bacterial urease activity in faeces: evidence of altered intestinal microflora?, 689
 - intra-articular pressure measurement in children, (A1) 78
 - mapping the GroEL epitope recognised by a synovial fluid T cell clone from a boy, (A1) 162
 - methotrexate treatment every other week, (L) 1030
 - pauciarticular, neurofibromatosis masquerading as, (A2) 39
 - physiotherapy and occupational therapy for: custom and practice in five centres in the UK, USA and Canada, 695
 - polyarticular
 - disablement correlates—a cross-sectional study, 91
 - multiple joint corticosteroid injections: a single-blind one year follow-up study, (A1) 121
 - prevalence of chronic anterior uveitis in New Zealand children, (A2) 25
 - T-cell receptor variable gene segment TCRBV12S4 null allele frequency, (A1) 80
 - uveitis, antibodies against retinal S-antigen, (L) 101
 - young adults with arthritis: meeting their transitional needs, 84
- KABASAKAL Y, GARRETT SL, CALIN A. Epidemiology of spondylodiscitis in ankylosing spondylitis—a controlled study, 660
- KADIOGLU A, SHELTON P. Adhesion of rheumatoid lymphocytes to mucosal endothelium: the gut revisited, 218
- KADIOGLU A *see also* Sheldon P
- KADIR M *see* Gibson T
- KAHN M-F *see* de Bandt M
- KAHN MF *see* Cordonnier C
- KAINBERGER F, TRATTNIG S, CZERNY C, SEIDL G, KRITZ H, IMHOF H. MRI assessment of the systemic manifestations of rheumatological disease, (S3) 39
- KAIPIAINEN-SEPPÄNEN O. Incidence of psoriatic arthritis in Finland, 1289
- KALDEN JR *see* Manger K
- KALDEN RW *see* Burkhardt H
- KALLENBERG CGM *see* Spronk PE
- KAMEL M *see* Elbeialy A
- KANZLER JM *see* Ali NN
- KANE D, KERR JR, MCQUAID S, CURRAN MD, MOORE JE, CROWLEY B, LEONARD N, COYLE PV, O'BRIEN S, MULCAHY FM, ALLEN IV, FERGUSON WP. Low incidence of arthritis in persistent human parvovirus B19 infection in immunocompromised HIV-infected patients, (A1) 167
- KANG RY, FREIRE-MOAR J, SIGAL E, CHU C-Q. Expression of cyclooxygenase-2 in human and an animal model of rheumatoid arthritis, 711
- KANIS A *see* O'NEILL TW
- KANIS J *see* Coombes G
- KANIS JA *see* O'Neill TW
- KANNER S *see* Ng T
- KAPILA S *see* Damsar F
- KARLSON EW. *Re: Ehrenstein et al.* (L) 195
- KARRAR A, EL RAMAHI KM, AL JISHI F, AL DALAAN A, AL SHAIKH A. Neurological manifestations in Saudi patients with systemic lupus erythematosus, (A1) 32
- KARRAR A *see also* El Ramahi KM
- KASSIMOS D, CHOY EHS, GROSSMAN AB, CHIKANZA IC, PANAYI GS. Endogenous opioid tone in patients with rheumatoid arthritis, 436
- KASSIMOS D *see also* Choy EHS
- KAUPPI M, ANTILA P. Stiff collar for the treatment of rheumatoid atlantoaxial subluxation, 771
- KAWASUGI K *see* Matsuda J
- KAY L *see* Atiea G
- KAY LJ, HESLOP PS, CHANDLER C, WALKER DJ. Objective measurement of patient activity before and after steroid injection of the knee in rheumatoid arthritis, (A1) 208
- KAY RA *see* Donn RP
- KEAT ACS *see* Stagg AJ
- KEEN J *see* Holt PJL
- KEEN M *see* Creamer P
- KEEN R *see* Masud T
- KEEN RW, GRIFFITHS GO, MAJOR PJ, GARNERO P, LANCHBURY JS, SPECTOR TD. Polymorphisms of the vitamin D receptor and bone turnover in female twins, (A1) 88
- KEEN RW, WOODFORD-RICHENS KL, LANCHBURY JS, SPECTOR TD. Polymorphism at the oestrogen receptor gene and bone mass in pre- and post-menopausal women, (A1) 89
- KEEN RW *see also* Griffiths GO
- KEENAN J *see* Pal B
- KELLEHER D *see* Donnelly S; McGonagle D
- KELSEY A *see* Baildam EM
- KEMM JR *see* Maggs FM
- KENNEDY LG, GOMEZ K, TAYLOR A, CALIN A. Do childhood infections affect the development of ankylosing spondylitis, (A2) 23
- KENNEDY LG *see also* Gomez K; Roussou E
- KENNY J *see* Dawson JK
- KENT RM, WRIGHT D, CHANDLER BJ. Survey of disability associated with rheumatic disease in a rural population, (A1) 131

- Keratan sulphate and chondroitin markers in normal and arthritic synovial fluid, 951
- Keratoconjunctivitis sicca in rheumatoid arthritis: prevalence in an outpatient population and role of Opticare eye drop dispenser, (A2) 20
- KERIN AJ *see* Adams MA
- KERR A *see* Rudge SR
- KERR J. Parvovirus B19 in rheumatoid arthritis, (L) 494
- KERR JR *see* Kane D
- KERSLAKE S *see* Khamashta MA
- KERSTENS P *see* Stolk J
- KHAMASHTA M *see* Méndez MJ; Rudwaleit M
- KHAMASHTA MA, BUCHANAN NMM, LIMA F, KERSLAKE S, HUNT BJ, HUGHES GRV. Obstetric outcome in systemic lupus erythematosus and antiphospholipid syndrome: the St Thomas' Hospital Lupus in Pregnancy clinic experience, (A1) 123
- KHAMASHTA MA *see also* Amengual O; Ames PRJ; Atsumi T; Ruiz-Irastorza G
- KHAMIS A *see* Malaviya AN
- KHATTAK FH. Sulphasalazine-induced systemic lupus erythematosus in a patient with rheumatoid arthritis, (L) 104
- KHOA ND *see* An TN
- KHURANA S *see* Leitch DN
- KIDD BL *see* Cruwys SC; Morris VH
- KIELY PDW. Lymphatic function in inflammatory arthritis, (L) 396
- KIELY PDW, OLIVEIRA DBG. Anti-CD8 treatment reduces the severity of inflammatory arthritis, but not vasculitis, in mercuric chloride-induced autoimmunity, (A1) 10
- KIFF PS *see* Walker DJ
- KILFEATHER SA *see* Mullaney P
- KIND P *see* Hurst NP
- KINDELAN JM *see* Gonzalez-Dominguez J
- KINGSLEY GH *see* Campbell F; Choy EHS; Iannone F; Johnson J; Wilkinson NZ
- KINKAD-WEEKS R *see* Leak AM
- KIRK BA *see* DEMSAR F
- KIRKHAM B *see* Hill C
- KIRWAN J *see* Hickling P; Taylor DJ
- KIRWAN JR. Rheumatology outpatient workload increases inexorably, (A2) 29
- KIRWAN JR *see also* Carr AJ; Cooper AM; Hewlett S; Leader KA; Reid DM; Sharif M
- KISS I *see* Makula É
- KISSLING RO *see* Weber M
- KISSONERGHIS M, FELLOWS R, FELDMANN M, CHERNAJOVSKY Y. Murine p75 TNF receptor promoter region: DNA sequence and characterization of a cis-acting silencer, (A1) 9
- KITA Y *see* Furukawa M
- KITAS G *see* Exley A
- KITAS GD *see* Koutedakis Y; Luqmani RA
- KJELDSEN-KRAGH J, SUMAR N, BODMAN-SMITH K, BROSTOFF J. Changes in glycosylation of IgG during fasting in patients with rheumatoid arthritis, 117
- KLAASSEN J *see* Olivé A
- KLEBSIELLA, Proteus antibodies and the peptides HLA-B27 and HLA-DR4 (DW15) in Japanese patients with ankylosing spondylitis and rheumatoid arthritis, (A2) 12
- Klebsiella pneumoniae*, similarly increased serum IgA1 and IgA2 subclass antibody levels against in ankylosing spondylitis patients with/without extra-articular features, 125
- KLEIN G *see* Riemsma RP
- KLINKE J *see* Rudwaleit M
- Knee
- and hip, association of abnormalities on physical examination with locomotor disability in the Rotterdam study, 884
 - laxity in patients with osteoarthritis and rheumatoid arthritis, 560
 - MRI evaluation in rheumatoid arthritis, (S3) 35
 - steroid injection, objective measurement of patient activity before and after in rheumatoid arthritis, (A1) 208
 - total replacement
 - effects in patients with rheumatoid versus osteoarthritis, (A2) 32
 - pre-admission education for patients, (A2) 31
- Knee joint
- evaluation of the efficacy of radiosynovectomy with gadolinium-enhanced magnetic nuclear resonance, (A1) 224
 - human meniscus: studies of proteoglycan and matrix protein synthesis and composition, (A1) 238
 - MRI can differentiate effects of intra-articular therapy on synovial proliferation and acute inflammatory changes, (A1) 205
 - proliferative synovitis, rheumatoid and psoriatic, long-term sonographic follow-up, 155
 - space width measurement: an experimental study of the influence of radiographic procedure and joint positioning, 761
 - synovitis, rheumatoid and psoriatic, arthroscopic synovectomy: long-term outcome, 463
 - trauma, serum YKL-40 levels in healthy children and adults. Comparison with serum and synovial fluid levels of YKL-40 in patients with osteoarthritis or trauma of the knee joint, 553
- Knee pain
- acute and chronic, patterns of symptom impact, (A1) 133
 - amongst the poor and affluent in Pakistan, 146
 - and disability, factors affecting in the community: the importance of obesity and psychological factors, (A1) 134
 - impact of a common symptom: validation of the disease repercussion profile, (A1) 133
 - psychological impact, (A1) 135
 - quadriceps strength and depression are important factors in and in associated disability, (A1) 135
- KNEKT P *see* Heliövaara M
- KNIGHT J, MCCAFFERTY A, NOON E, BELL A. Indirect evidence to support the production of hydroxyl and/or ferryl radicals at sites of inflammation, (A2) 13
- KNIGHT SC *see* Stagg AJ
- KNOWLES RG *see* McNicholas MJ
- KO S *see* Mok CC
- KO TC *see* Szeto CC
- KOBAYASHI H, TADA S, FUCHIGAMI T, OKUDA Y, TAKASUGI K, MATSUMOTO T, IIDA M, AOYAGI K, IWASHITA A, DAIMARU Y, FUJISHIMA M. Secondary amyloidosis in patients with rheumatoid arthritis: diagnostic and prognostic value of gastroduodenal biopsy, 44
- KOEHLER L *see* Dreses-Werringloer U
- KOELLER M *see* Templ E
- KOH ET, LEE P, GLADMAN DD, ABU-SHAKRA M. Pulmonary hypertension in systemic sclerosis: an analysis of 17 patients, 989
- KOH W-H *see* Haigh RC
- KOH WH, PANDE I, JONES S, SAMUELS A, CALIN A. Placebo-controlled study of low dose amitriptyline in ankylosing spondylitis, (A1) 155
- KOHLMAN T, RASPE H, DECK R, BÜTZOW C. Pain coping strategies as predictors for graded back pain status: results of a population based follow-up study, (A1) 142
- KOIKE T *see* Amengual O; Atsumi T
- KONDEATIS E *see* Hameed K
- KONİÇE M *see* Gül A
- KORDOSSIS T *see* Moutsopoulos HM
- KORTEKANGAS P *see* Roivainen A
- KOS-GOLJA M *see* Jevtic V
- KOUTEDAKIS Y, BEGUM S, LABIB M, DELAMERE JP, KITAS GD. Causes of cachexia in rheumatoid arthritis, (A1) 200
- KOVÁCS A *see* Makula É
- KOVÁCS L *see* Makula É
- KRAUSE A *see* Baerwald CGO
- KRETSCHMER G *see* Mau W
- KRITZ H *see* Kainberger F
- KRUSS B *see* Stei P
- KUIS W *see* van der Net J
- KUNTZ D *see* Lioté F
- KUNTZ JL *see* Combe B
- KURTHEN R *see* Huskisson EC
- KVIEN TK, GLENNÄS A, MELBY K. Prediction of diagnosis in acute and subacute oligoarthritis of unknown origin, 359
- KVIEN TK *see also* Bas S; Smedstad LM
- L-selectin, soluble, reduced levels in systemic sclerosis and vasculitis, (A1) 42
- LA CIVITA L *see* Ferri C
- LAASONEN L *see* Pettersson T
- LABIANO I *see* Andreu JL; Diez E; Mulero J
- LABIB M *see* Koutedakis Y
- LAFABER FPJG *see* van der Veen MH; Van Valburg AA; Verhoef CM; Verschure PJ
- LAKOMEK H-J *see* Schade C
- LALIES MD *see* Harbuz MS
- LAMB A *see* Luqmani RA
- LAMBERT M *see* Houssiau FA
- Lamellar bodies in type B synoviocytes and proteolipid in synovial fluid and the articular lining, 821
- LÄMMLÉ B *see* Lehmann T
- LAMPREAVE JL *see* Carreño L
- LANCHBURY J *see* Hameed K; Yanni G
- LANCHBURY JS *see* Baerwald CGO; Bowman SJ; Coakley G; Griffiths GO; Keen RW; Öcal L
- LANYON P, JONES A, DOHERTY M. Skyline are better than lateral radiographs for detecting progression of patellofemoral osteoarthritis, (A1) 98
- LANZA FL, CARLSON MR. Effects of an amino-versus and alkyl-bisphosphonate on the healing of aspirin-induced stomach erosions in healthy subjects, (A2) 7
- LAPHAM G *see* Bevis PJR

- LARBRE JP *see* Hall GM
- Large granular lymphocyte
expansions in Felty's syndrome have an unusual phenotype of activated CD45RA+ cells, 1252
- leukaemia, anti-TCRBV monoclonal antibodies in the detection and monitoring in Felty's syndrome, (A1) 182
- LARREA A *see* Andreu JL; Díez E; Isasi C; Mulero J; Perez A
- LARRODERA L *see* Fernández-Dapica P
- Laser therapy, low energy: effect on papain-induced OA, (A2) 7
- LASSILA O *see* Isomäki P
- LASTRA GC *see* Leader KA
- LATCHMAN DS *see* Ravirajan CT
- LATHROP M *see* Hardwick L; Hardwick LJ
- LAU CS *see* Mok CC
- LAU YL *see* Mok CC
- LAUFENBERG M *see* Baerwald CGO
- LAVERSUCH CJ, TRIBBICK G, WESTWOOD OMR, BRUCKNER FE, HAY FC
Rheumatoid factor epitope profiles in rheumatoid arthritis, (A1) 173
Using the Pepscan technique to map binding of rheumatoid factor to IgG, (A1) 173
- LAVERSUCH CJ *see also* Tobias JH
- LAVERY B *see* Ho M
- LAWRENCE R *see* Şahin (Özgün) Ş
- LAWSON TM, WANG EY, BORYSIEWICZ LK. Anti-TCRBV monoclonal antibodies in the detection and monitoring of large granular lymphocyte leukaemia in Felty's syndrome, (A1) 182
- LAWSON TM, WANG EY, VEDHARA K, BORYSIEWICZ LK. CD8+ T cells and their T cell receptor repertoires in rheumatoid arthritis, (A1) 183
- LAWSON TM, WILLIAMS BD. Silent myocardial infarction in Wegener's granulomatosis, 188
- LAYCOCK CA *see* Warnock MG
- Lba5 and ribosomal P autoantibodies in NP-SLE, (A1) 111
- LE GOFF P *see* Sarau A
- LE LOET X *see* Wojtulewski JA
- LE THI HUONG D, WECHSLER B, PIETTE J-C, ARFI S, GALLINARI C, DARBOIS Y, FRANCES C, GODEAU P. Risks of ovulation-induction therapy in systemic lupus erythematosus, 1184
- LEADER KA, LASTRA GC, KIRWAN JR, ELSON CJ. Agalactosyl IgG in aggregates from the rheumatoid joint, 335
- LEAK AM, WOODS Z, BERTI A, HART K, KINKEAD-WEEKS R, DUDDY K. Multidisciplinary study of outpatient education in patients with lower limb osteoarthritis, (A1) 103
- LEAL-CERRO A *see* De La Iglesia JL
- LEAN EM *see* Ellis AJ
- LEARMONTH ID *see* Perry MJ
- LEDINGHAM J, REGAN M, JONES A, DOHERTY M. Symmetry of radiographic change in hospital-referred knee osteoarthritis, (A1) 97
- LEE A *see* Biagini P
- LEE FD *see* Morris AJ
- LEE P *see* Koh ET
- LEES C, CARTY H, WALSH HPJ, DAVIDSON JE. SAPHO syndrome in an eleven year old girl, (A2) 36
- LEFEBVRE C *see* Houssiau FA
- Leflunomide, eruption of vasculitis during treatment for rheumatoid arthritis, (A1) 226
- Left ventricular function in scleroderma, 983
- LEHMANN T, BEYELER C, LÄMMLER B, HUNZIKER T, VOCK P, OLAH AJ, DAHINDEN C, GERBER NJ. Severe osteoporosis due to systemic mast cell disease: successful treatment with interferon alpha-2b, 898
- LEHTINEN K *see* Mäki-Ikola O
- LEIGH RD, CAMBRIDGE G, EDWARDS JCW. Facilitated expression by synovial fibroblasts of factors involved in B lymphocyte maturation and survival, (A1) 110
- LEIRISALO-REPO M, ROOK GAW, HERNANDEZ-MUNOZ H. Agalactosyl IgG is a marker of severe spondylitis, (A1) 153
- LEIRISALO-REPO M *see also* Allen RL; Pettersson T
- LEITCH DN, KHURANA S, FORDHAM JN. Incidence of osteoporosis in patients attending fracture clinic, (A2) 6
- LEITCH DN. Reactive arthritis, β -haemolytic *Streptococcus* and *Staphylococcus aureus*, (L) 912
- LEMARCHAND P *see* Touray M
- LONDON M *see* Holt PJL
- LEONARD N *see* Kane D
- Letter from
Japan, 579
Kuwait, 380
- Leukotriene biosynthesis, study of the cooperation between chondrocytes and polymorphonuclear leucocytes, (A1) 62
- LEUNG B *see* McInnes IB
- Lewis X trisaccharide, expression on serum IgG in rheumatoid arthritis, (A1) 175
- LHOTÉ F *see* Guillevin L
- LI J-M *see* Horsfall AC
- LI KY *see* Szeto CC
- LIANG HY. Overexpression of proto-oncogene bcl-2 in rheumatoid synovium, (L) 803
- LIDDICOTT AJ *see* Yeap SS
- LIENCE E, MARSAL S, ARMADANS L, GALLARDO D, MARTINEZ M, RIBERA A. Clinical, radiographic and HLA associations as markers for psoriatic arthritis patterns, (A1) 159
- LIEW FY *see* McInnes IB
- LIGHTMAN SL *see* Harbuz MS
- LILES JH, PINDER IM. Effects of total knee replacements in patients with rheumatoid versus osteoarthritis, (A2) 32
- LILLEY J *see* Devlin J
- LIM KKT. Interference by IgM rheumatoid factor on ELISA for antibodies to mycobacterial 65 kDa heat shock protein in rheumatoid arthritis, (L) 300
- LIM KL. Cervical spondylodiscitis in a patient with ankylosing spondylitis, (L) 1034
- LIM KL, FOX R, ABDUL-WAHAB R, POWELL RJ. Previously undiagnosed systemic lupus erythematosus in acute psychiatric hospital admissions, (A1) 36
- LIM KL, MOUG SJ, STURROCK RD. Acute physiological effect of hydrotherapy in patients with ankylosing spondylitis, (A1) 156
- LIMA F *see* Khamashta MA; Ruiz-Irastorza G
- LIMBURG PC *see* Spronk PE
- LIN H *see* Bavington C; Brennan FR; Jobanputra P
- LINDEN B, DISTEL M, BLUHMKE E. Double-blind study to compare efficacy and safety of meloxicam 15 mg with piroxicam 20 mg in patients with osteoarthritis of the hip, (S1) 35
- LINGARD E *see* Chatterjee S
- LINGARD EA, SOLAN SK, PINDER IM, SMITH SR, MORAN CG. Pre-admission education for patients undergoing total knee replacement, (A2) 31
- Link protein and aggrecan mRNA quantitation in human articular cartilage of different ages by competitive RT-PCR, (A1) 237
- Linkage analysis studies of candidate genes on affected sib pair RA families, (A1) 190
- Linoleic acid is a better substrate for PGH-synthase-2 than for PGH-synthase-1 tested in whole endothelial cells in culture, (A1) 219
- LIOTÉ F, BROUSSE C, FRIARD S, VIGNON D, PAUNET M, STERN M. Calcitropic hormones and osteocalcin levels in lung transplant recipients, (A1) 86
- LIOTÉ F, BROUSSE S, FRIARD H, MAL H, SLIEMAN C, FOURNIER M, STERN M. Osteoporosis and bone morbidity in lung transplant recipients, (A1) 85
- LIOTÉ F, PIZZUTI P, WEYL-CLERC D, CERF I, BELLAICHE L, DRYLL A, KUNTZ D. Rheumatoid arthritis, serum IgA immunoglobulins and sacroiliitis, (A1) 201
- LIOTÉ F *see also* Combe B
- Lipocortin, does it mediate the inhibition of adhesion molecule expression induced by glucocorticoids?, (A1) 64
- Lipocortin-1, intracellular, neutrophil adhesion to endothelial cells is a signal for loss, (A1) 60
- Lipopolysaccharide, *Escherichia coli*, induction of chronic autoimmune arthritis in DBA/1 mice by oral administration, 828
- LISSE JR *see* Acebes JC
- Liver involvement and hepatitis C virus infection in patients with primary Sjögren's syndrome, (A1) 15
- Liver transplant recipients, rheumatic manifestations, (A1) 87
- LLOBET JM *see* Giral P
- LLOYDS D, DAVIES EV, WILLIAMS BD, HALLETT MB. Tyrosine phosphorylation in neutrophils from synovial fluid of patients with rheumatoid arthritis, 846
- Loading
sustained
makes articular cartilage stronger, (A1) 240
reduces the stress-equalising property of cartilage, (A1) 239
- Local tissue tolerability of meloxicam, a new NSAID: indications for parenteral, dermal and mucosal administration, (S1) 44
- LOCKWOOD CM *see* Jayne DRW
- Locomotor disability, association of abnormalities on physical examination of the hip and knee in the Rotterdam study, 884
- LOGAR D *see* Jevtic V
- LOIZOU S, CAZABON JK, SO AKL, TAIT D, WALPORT MJ. Antiphospholipid antibodies from patients infected with human B19 parvovirus have similar specificity and co-factor dependence to those found in SLE patients, (A1) 22
- LOMBARDI G *see* Scarpa R
- LOMBARDINI F *see* Ferri C
- LONDEI M *see* Chu CQ
- LONG A *see* Donnelly S
- LONGOMBARDO G *see* Ferri C
- LÓPEZ A *see* Collantes E
- LÓPEZ E *see* Collantes E
- LÓPEZ-ARMADA MJ *see* Palacios I; Sánchez- Pernaute O

- LÓPEZ-LONGO FJ, GRAU R, MONTEAGUDO I, GONZÁLEZ M, MINGUEZ MD, ABAD MA, ALONSO J, CARREÑO L. Dissociation of immune response to the degraded and nondegraded (native) Sa antigens in rheumatoid arthritis and systemic lupus erythematosus, (A1) 18
- LÓPEZ-LONGO FJ *see also* Abad MA; Carreño L; González M; Monteagudo I
- LÓPEZ-SOTO A *see* Navarro M
- LORBER M *see* Ehrenfeld M
- LORENZEN I *see* Johansen JS; Østergaard M
- LORENZO JC *see* Olivé A
- LORENZO P *see* McAlinden A
- LORY C *see* Hamilton I
- LOTTI T *see* Matucci Cerinic M
- LOUGHLIN JA *see* Spector TD
- LOVAT LB *see* Persey MR
- Low back pain
developing a care path from primary through secondary to tertiary care, (A1) 143
incident in nurses, risk factors for in nurses, (A1) 123
occurrence of in relation to psychosocial factors in the workplace, (A1) 141
- Low back school, prospective controlled study in the general population, 178
- Lower gastrointestinal and renal side effects of the COX-2 inhibitor meloxicam, compared to naproxen in rheumatoid arthritis patients, (A1) 223
- LOWRY PJ *see* Woods RJ
- LUBRANO E, CIACCI C, AMES PRJ, MAZZACCA G, ORIENTE P, SCARPA R. Arthritis of coeliac disease: prevalence and pattern in 200 adult patients, 1314
- LUBRANO E *see also* Scarpa R
- LUGER A *see* Templ E
- Lumbar spine bone density measurements: the effect of posterior facet joint osteoarthritis in a middle age female population, (A1) 98
- LUNEC J *see* Evans MD; Hill CM
- Lung disease, HRCT diagnosed interstitial, clinical features associated with rheumatoid arthritis, (A2) 16
- Lung transplant recipients
calcitropic hormones and osteocalcin levels, (A1) 86
osteoporosis and bone morbidity, (A1) 85
- Lupoid hepatitis or lupus with mesenteric vasculitis, 1309
- LUPOLI S *see* Ames PRJ
- Lupus anticoagulant, phosphatidyl serine-dependent antiprotease antibody is exclusive to patients with, 589
- Lupus flare, increased rate during pregnancy and the puerperium: a prospective study of 78 pregnancies, 133
- Lupus or lupoid hepatitis with mesenteric vasculitis, 1309
- Lupus patients
black female, clinical features and autoantibodies: cross-sectional and longitudinal studies, (A1) 30
enterococcal arthritis with avascular necrosis, 595
- Lupus-prone mice, BXSb, effect of dominant disease susceptibility genes on phenotype, (A1) 112
- LUQMANI R *see* Exley A
- LUQMANI RA, TALBOT H, LAMB A, ARMSTRONG C, KITAS GD, EXLEY A, MOOTS RJ, BACON PA, RASMUSSEN N, European Study Group for Therapeutic Trials in Systemic Vasculitis. Vasculitis integrated total assessment log (vital): computerised system for assessment of vasculitis, (A1) 54
- LUUKKAINEN R *see* Isomäki P
- Lyme arthritis, pattern of joint involvement, 1016
- Lyme borreliosis in rheumatological practice: identification of Lyme arthritis and diagnostic aspects in a Swedish county with high endemicity, 853
- Lymphatic function in inflammatory arthritis, (L) 396
- Lymphocyte adhesion, is the corticosteroid mediated change to mucosal endothelium by mononuclear cells from rheumatoid arthritis patients due to a quantitative change in lymphocyte subsets?, (A1) 14
- Lymphocytes, rheumatoid, adhesion to mucosal endothelium: the gut revisited, 218
- Lymphoid cells capable of transferring disease as carriers of immunomodulators—a rheumatoid arthritis model, (A1) 182
- LYNCH JA *see* Buckland-Wright JC
- LYNCH MP *see* Dawson JK
- LYONS CBA *see* Suzuki H
- MCALINDEN A, BAYLISS MT, LORENZO P, HEINEGÅRD D. Human meniscus: studies of proteoglycan and matrix protein synthesis and composition, (A1) 238
- MCAFFERTY A *see* Knight J
- MCCALL E *see* Halsey J
- MCCANN BG *see* Reid AJC
- MCCARTHY D *see* Bolger T
- MCCARTHY DA *see* Carbon RJ
- MCCCHIONI P *see* Boiardi L
- MCCLOSKEY DA *see* Seibold JR
- MCCLOSKEY E *see* Coombes G
- MCCLOSKEY EV *see* O'NEILL TW
- MCCREADY S *see* Mack CS
- MCDONAGH JE, DUNN A, OLLIER WER, WALKER DJ. Compound heterozygosity for the shared epitope and the risk and severity of rheumatoid arthritis, (A1) 187
- MCDONAGH JE, WALKER DJ. Mortality in extended pedigree multicase RA families, (A2) 11
- MCDONAGH JE *see also* Davidson BKS
- MACDONALD AG, BIRKENSHAW G, DURHAM B, OVER K, BUCKNALL RC, FRASER WD
Biochemical markers of bone turnover in ankylosing spondylitis, (A1) 151
Effects of prednisolone and deflazacort on bone metabolism in polymyalgia rheumatica, (A1) 58
- MACDONALD AG *see also* Over KE
- MCENTEGART A, DUNCAN MR, CAPELL HA, MADHOK R, PORTER DR, THOMSON EA. Intramuscular gold therapy in 440 rheumatoid patients—results over 5 years in relation to level of social deprivation at outset, (A1) 213
- MCENTEGART A *see also* Capell HA; Duncan MR; Morrison E
- MACEY MG *see* Carbon RJ
- MACFARLANE GJ, PAPAGEORGIOU AC, THOMAS E, CROFT PR, JAYSON MIV, SILMAN AJ. Occurrence of low back pain in relation to psychosocial factors in the workplace, (A1) 141
- MACFARLANE GJ, PRITCHARD CM, PAPAGEORGIOU AC, SCHOLLUM J, CROFT PR, SILMAN AJ. Chronic widespread pain: is a better classification possible?, (A1) 136
- MACFARLANE GJ, THOMAS E, BURT J, CROFT PR, SILMAN AJ. Follow-up study of chronic widespread pain in the community: a better prognosis than in the clinic, (A1) 137
- MACFARLANE GJ, THOMAS E, PAPAGEORGIOU AC, CROFT PR, JAYSON MIV, SILMAN AJ. Employment and work activities as predictors of future back pain, (A1) 141
- MACFARLANE GJ *see also* Nahit ES; Pope DP; Pope DR
- MACFARLANE L *see* Pal B
- MACFARLANE XA, DIXEY J, DUNPHY J, BHALLA AK, ROBERTSON DAF. Bone mineral density and bone turnover in treated coeliac disease, (A1) 85
- MCGONAGLE D, TODD A, FARRELL R, ORMOND P, MONTAGUE S, KELLEHER D, CASEY EB. Rheumatic manifestations of hepatitis C virus infection, (A1) 168
- MCGONAGLE D *see also* Montague S; Ormond P
- MACGREGOR AJ, BUSCOMBE J, HILSON AJW, AGNEW JE, BLACK CM. Risk factors for progression of oesophageal dysmotility in systemic sclerosis, (A1) 44
- MCGURTY DW *see* McNicholas MJ
- MACHEIN U, CONCA W. Rapid RT-PCR based matrix metalloproteinase gene family screening reveals expression of several metalloproteinases in monocytic cells, (A1) 63
- MACHTEY I. Neurological signs in RA patients receiving gold, (L) 803
- MCUGH N *see* Mackay K
- MCUGH NJ, HARVEY G, MADDISON PJ, BLACK C, FANNING G, WELSH K. HLA-DQB1 polymorphism and antibodies to RNA polymerases in systemic sclerosis, (A1) 45
- MCUGH NJ *see also* Balakrishnan C; Flora RS; Haigh RC; Harvey GR; Jones SM; Patel YI
- MACIEWICZ R *see* Creamer P
- MCINTOSH E. Cost of rheumatoid arthritis, 781
- MCINNES I *see* Porter DR
- MCINNES IB, FIELD M, AL-MUGHAIJES J, LEUNG B, HUANG FP, WILKINSON PC, STURROCK RD, LIEW FY. Interleukin-15 expression in rheumatoid arthritis, (A1) 180
- MACK CS, MCCREADY S, HALL N, HOUGH D, JULKUNEN H, MADDISON PJ. Anti-p57, sero-epidemiology and correlations with HLA, (A1) 24
- MACK CS, MACKAY K, CALIN A. Radiology in ankylosing spondylitis: comparison of two scoring systems, (A2) 23
- MACK CS, WHITELAW D. Clinical audit. Isotope bone scans for foot problems, (A2) 8
- MACK CS *see also* Haigh RC
- MACKAY K, FLORA R, BARNES J, PATEL Y, MCUGH N. Joint involvement in connective tissue disease and its associations with the third hypervariable region (HVR3), (A1) 30
- MACKAY K *see also* Mack CS
- MCKENNA F *see* Green M; Mason JC
- MCKENNA LA, BOLTON MC, MASON RM, BAYLISS MT. Ferritin H-chain mRNA is upregulated in human articular cartilage stimulated with TNF α , (A1) 4
- MCKENNA S *see* Houssein DA
- MACKENZIE A *see* Halsey J
- MCKENZIE JF *see* Morris AJ
- MACKILLOP N *see* Nandra D
- MACKWORTH-YOUNG CG *see* Harmer IJ
- MCLAREN M *see* Coggon D

- McLURE CE, MENON S, ISENBERG DA. Causes of death in two hundred patients with systemic lupus erythematosus, (A1) 37
- McMICHAEL AJ see Allen RL; Bowness P
- McNALLY J see Bell AL
- McNICHOLAS MJ, ALDBERTH T, DRYSDALE MJ, ROWLEY DI, MCGURTY DW, ABDON P, KNOWLES RG. Synovial fluid from knees with secondary osteoarthritis has much higher nitrite concentrations than serum, suggesting local production of nitric oxide, (A1) 105
- MACPHERSON H see Grabowski PS
- MCQUAID S see Kane D
- 27E10 Macrophages, distribution in synovium of inflammatory and degenerative joint disease, (A1) 178
- McRORIE ER see Richmond R
- MADDISON PJ see Balsa A; Haigh RC; Harvey GR; McHugh NJ; Mack CS
- MADHOK R see Capell HA; Crilly A; Duncan MR; McEntegart A; Morrison E; Porter DR
- MADHOO S see Persey MR
- MADRIGAL MJ see Graña J
- MAGEED RA see Burd GP; Harmer IJ; Moyes SP
- MAGGS FM, JUBB RW, KEMM JR. Single-blind randomized controlled trial of an educational booklet for patients with chronic arthritis, 775
- MAGILL K see Bell AL
- Magnetic resonance imaging of acromegaly, (A1) 95
- in the assessment of synovium and cartilage, (S3) 3
- assessment of the systemic manifestations of rheumatological disease, (S3) 39
- can differentiate effects of intra-articular therapy on synovial proliferation and acute inflammatory changes in the knee joint, (A1) 205
- contrast-enhanced of rheumatic joint disease, (S3) 44
- evaluation of the knee in rheumatoid arthritis, (S3) 35
- gadolinium-enhanced of the finger joints in rheumatoid arthritis, (A1) 118
- hand changes in early rheumatoid arthritis, (A1) 118
- high-resolution of erosive changes in the metacarpophalangeal joint in rheumatoid arthritis, (A1) 124
- intravenous contrast enhancement of inflammatory synovium: a dose study, (S3) 31
- mapping abnormal synovial vascular permeability in temporomandibular joint arthritis in the rabbit, (S3) 24
- and plain radiography, comparative study in the assessment of sacroiliitis in early spondylarthritis, (A1) 117
- prognostic value of contrast enhanced Gd-DTPA MRI for development of bone erosive changes in rheumatoid arthritis, (S3) 27
- quantitative assessment of the synovial membrane in the rheumatoid wrist: an easily obtained MRI score reflects the synovial volume, 965
- role in the assessment of rheumatological disease, (S3) 1
- shows brain lesions in patients with Behçet's disease without central nervous system symptoms, (A1) 52
- MAGNUSSON S. Treatment of rheumatoid arthritis—does it affect society's cost for the disease?, 791
- MAHMUD T, WRIGGLESWORTH JM, RAFI S, SCOTT DL, BJARNASON I. Non-acidic pro-NSAIDs and COX-2 inhibitors do not uncouple mitochondrial oxidative phosphorylation, (A1) 222
- MAHWINNEY B see Billingham MEJ
- MAILLEFERT JF, MAYNADIE M, TEBB JG, AHO S, WALKER P, CHATARD C, DULIEU V, BOUVIER M, CARLI PM, TAVERNIER C. Expression of the multidrug resistance glycoprotein 170 in peripheral blood lymphocytes of rheumatoid arthritis patients, 430
- MAINI R see Paleolog E
- MAINI RN see Allard SA; Brennan FM; Burd GP; Davis DM; Horsfall AC; Isenberg DA; Malfait A-M; Marinova L; Moyes SP; Ross S; Taylor PC
- MAJOR PJ see Keen RW
- MAJORI M see Manganelli P
- MÄKI-IKOLA O, LEHTINEN K, GRANFORS K. Similarly increased serum IgA1 and IgA2 subclass antibody levels against *Klebsiella pneumoniae* bacteria in ankylosing spondylitis patients with/without extra-articular features, 125
- MAKULA É, POKORNY G, RAJTA M, KISS I, KOVÁCS A, KOVÁCS L. Parotid gland ultrasonography as a diagnostic tool in primary Sjögren's syndrome, 972
- MAL H see Lioté F
- MALAVIYA AN, AL-AWADI A, AL-SAJED K, AL-JARALLAH K, SHEHAB D. Letter from Kuwait, 380
- MALAVIYA AN, MAROUF R, AL-JARALLAH K, AL-AWADI A, AL-SAJED K, AL-GAURER S, KHAMIS A, AL-SALEM I, RAGHUPATHY R, AL-MAZIDI M, SEREBOUR F, UMAMAHESWARAN I, MOUROU M, MOKHTAR M, QURTOM M, RAOOF AL-SHAHEB A, TARASEVICIUS T, HUSSAIN MAS. Hughes syndrome: a common problem in Kuwait hospitals, 1132
- MALCOLM A see Billingham MEJ
- MALFAIT A-M, BUTLER DM, BRENNAN FM, GATLEY M, MAINI RN, FELDMANN M. Modification of murine collagen-induced arthritis with antibodies to interleukin 12, (A1) 108
- MALIK J see Gordon C; Stoll T
- MALIN M, VERRONEN P, MYKKÄNEN H, SALMINEN S, ISOLAURI E. Increased bacterial urease activity in faeces in juvenile chronic arthritis: evidence of altered intestinal microflora?, 689
- MALLESON PN see Foster HE
- MALLIA C see Matucci Cerinic M
- MANCINI R see Boiardi L
- MANGANELLI P, SALAFFI F, SUBIACO S, CAROTTI M, CERVINI C, CONSIGLI G, MAJORI M, PESTI A. Bronchoalveolar lavage in mixed cryoglobulinaemia associated with hepatitis C virus, 978
- MANGER B see Manger K
- MANGER K, KALDEN JR, MANGER B. Cyclosporin A in the treatment of systemic lupus erythematosus: results of an open clinical study, 669
- MANJON P see Poveda MJ
- MANN J see Demsar F
- MANNA VK see Isaacs JD
- MANNING J see Bukhari M
- MANNION AF, DUMAS GA, STEVENSON JM, ESPINOSA FJ, FARIS MW, COOPER RG. Muscle fibre size and distribution of the back extensors: relationship with muscle fatigability, (A1) 144
- Normal muscle fibre size and type distribution of the human back extensors, (A1) 144
- Mannose binding protein gene dysfunctional allele in rheumatoid arthritis, (A1) 189
- polymorphism and rheumatoid arthritis, (A1) 188
- MANZANEQUE L see Sanchez J
- MANZI C see Nandra D
- MAPP P see Gaffney K
- MAPP PJ see Blake DR; Walsh DA
- Mapping and characterisation of the genes involved in rheumatoid arthritis, (A2) 10
- MARCHESONI A, CERAVOLO GP, ROSSETTI A, FANTINI F, TOSI S. Cyclosporin A in the treatment of adult Still's disease, (A1) 77
- MARCOS R see Gómez-Vaquero C
- MARELL J see Riemekasten G
- MARENCO JL see Sanchez J
- MARINI MG see Patoia L
- MARINOVA L, WILLIAMS RO, MASON LJ, FELDMANN M, MAINI RN. Pro-inflammatory cytokine expression in early collagen-induced arthritis, (A1) 6
- MARINOVA L see also Ross S
- Markers of bone metabolism, changes in early rheumatoid arthritis, (A2) 3
- MARKOV G, DOBRO J, SHANKMAN S, BELMONT HM. Enterococcal arthritis with avascular necrosis in a lupus patient, 595
- MARKS R see Rudge SR
- MARLES M see Hdez-Berian JA
- MARLOW A, JOHN S, HAJEER A, OLLIER W, SILMAN A, WORTHINGTON J. Sensitivity of different analytical methods to detect disease susceptibility genes in RA sibling pair families, (A1) 190
- MARLOW A see also John S; John SL
- MAROUF R see Malaviya AN
- MARSAL S see Lience E
- MARTEL-PELLETIER J, PELLETIER J-P. Wanted—the collagenase responsible for the destruction of the collagen network in human cartilage, (E) 818
- MARTIN JC. Case of Zieve's syndrome presenting with myalgia: not to be confused with polymyalgia rheumatica, (L) 495
- MARTIN JC, DESOYA R, O'SULLIVAN MM, SILVERSTONE E, WILLIAMS H. Chronic recurrent multifocal osteomyelitis: spinal involvement and radiological appearance, 1019
- MARTIN JC, O'SULLIVAN MM, BUTLER RC, DIXEY JJ, SHARP CA, DAVIE MWJ. Changes in markers of bone metabolism in early rheumatoid arthritis, (A2) 3
- MARTIN JC, STEWART A, REID DM. Discrimination of hip fractures by peripheral quantitative computed tomography (pQCT), (A2) 6
- MARTIN MOLA E see Muñoz-Fernández S
- MARTIN-MOLA E see De Miguel E
- MARTINEZ A see Graña J
- MARTINEZ M see Lience E
- MARTINEZ-CONCHA D see González J
- MARTÍNEZ-TABOADA V see Blanco R
- MARTÍNEZ-TABOADA VM see González-Gay MA; Rodríguez-Valverde V
- MARTINI A see Ferri C
- MARUBINI E see Ferraccioli GF
- MARZULLO P see Scarpa R
- MASANÉS F see Font C
- MASON H see Over KE
- MASON JC, ARNOLD MM, GRIFFITHS IGH, MCKENNA F. Gold plated stomach—myth or reality, (A1) 225
- MASON JC, YARWOOD H, SUGARS K, HASKARD DO. Human umbilical vein and dermal microvascular display heterogeneity in expression of surface antigens following activation of protein kinase, (A1) 11
- MASON JC see also Green M
- MASON LJ see Marinova L; Ross S

- MASON RM *see* McKenna LA
- MASUD T, KEEN R, NANDRA D, JAWED S, DOYLE DV, SPECTOR TD. Lumbar spine bone density measurements: the effect of posterior facet joint osteoarthritis in a middle age female population, (A1) 98
- MASUD T, MULCAHY B, THOMPSON AV, KEEN R, DOYLE DV, SPECTOR TD. 1,25-Dihydroxyvitamin D₃ combined with cyclical etidronate on spine and femoral neck bone density in postmenopausal women, (A1) 115
- MASUD T *see also* Cooper C
- MASUMI S *see* Fujikawa Y
- MAT C *see* Yazici H
- MATEO L *see* Gomez-Vaquero C
- MATHEY D *see* Hajeer A
- MATHIEU A *see* Ferraccioli GF
- MATHIEU P *see* Conrozier T
- MATHIS C *see* Maugars Y
- Matrix metalloproteinase gene family screening reveals expression of several metalloproteinases in monocytic cells, (A1) 63
- MATSUDA J, SAITO N, GOTOH M, KAWASUGI K, GOHCHI K, TSUKAMOTO M. Phosphatidyl serine-dependent antiprotease antibody is exclusive to patients with lupus anticoagulant, 589
- MATSUMOTO T *see* Kobayashi H
- MATSUMURA R *see* Furukawa M
- MATTESON EL *see* Isaacs JD
- MATTEY DL *see* Cheung NT; Salih AM; Scott LJ
- MATTHEWS N *see* Gordon C
- MATTHIS Ch *see* Raspe H
- MATUCCI CERINIC M, LOTTI T, VALENTINI G, PACE J, MALLIA C. Rheumatology and dermatology: sister specialties in Italy and Malta, 1312
- MAU W, BORNHANN M, WEBER H, WEIDEMANN HF. Deficient rehabilitation during the course of early RA, (A1) 228
- MAU W, BORNHANN M, WEBER H, WEIDEMANN HF, HECKER H, RASPE HH. Prediction of permanent work disability in a follow-up study of early rheumatoid arthritis: results of a tree structured analysis using RECPAM, 652
- MAU W, KRETSCHMER G, GROTHE H, ZEIDLER H. Slow radiographic progression in a 15-year follow-up of patients with undifferentiated spondylarthritis, (A1) 157
- MAUGARS Y, MATHIS C, BERTHOLET J-M, CHARLIER C, PROST A. Assessment of the efficacy of sacroiliac corticosteroid injections in spondylarthropathies: a double-blind study, 767
- MAURI C, WILLIAMS RO, WALMSLEY M, FELDMANN M. Pattern of cytokine secretion during the development of collagen induced arthritis, (A1) 5
- MAURI C *see also* Ross S
- MAXWELL S *see* Pal B
- MAYER B *see* Ryan SJ
- MAYNADIE M *see* Maillefert JF
- MAYORGA M *see* Val-Bernal F
- MAYOUX-BENHAMOU A *see* Ayral X
- MAZZACCA G *see* Lubrano E
- MEADOWS K *see* Sharif M
- Mechanism of action of NSAIDs, (A1) 1
- Medical education and rheumatology in Great Britain, 269
- MEDRANO C, SALAS E. Chronic hypophosphataemia: possible osteoporosis risk-factor, (A1) 96
- Megaloblastic anaemia, scurvy and osteoporosis due to alleged food intolerance, (L) 701
- MEIERS T *see* Billingham MEJ
- MELBY K *see* Kvien TK
- Meloxicam
- 4-week, double-blind, parallel-group study to compare gastrointestinal effects of meloxicam 7.5 mg and 15 mg, piroxicam 20 mg and placebo, (S1) 61
 - does not inhibit platelet aggregation and renal prostaglandin E₂ synthesis in healthy volunteers, (A1) 220
 - double-blind study to compare efficacy and safety of meloxicam 15 mg with piroxicam 20 mg in patients with osteoarthritis of the hip, (S1) 35
 - double-blind, three-week study to compare the efficacy and safety of meloxicam 7.5 mg and meloxicam 15 mg in patients with rheumatoid arthritis, (S1) 17
 - local tissue tolerability: indications for parenteral, dermal and mucosal administration, (S1) 44
 - long-term study to evaluate safety and efficacy of therapy in patients with rheumatoid arthritis, (S1) 29
 - lower gastrointestinal and renal side effects of the COX-2 inhibitor, compared to naproxen in rheumatoid arthritis patients, (A1) 223
 - new non-steroidal anti-inflammatory drug with an improved safety profile through preferential inhibition of COX-2, pharmacology, (S1) 4
 - open study to assess safety and tolerability of meloxicam 15 mg in subjects with rheumatic disease and mild renal impairment, (S1) 56
 - in osteoarthritis: a six-month, double-blind comparison with diclofenac sodium, (S1) 39
 - review of clinical pharmacokinetics, (S1) 13
 - safety: global analysis of clinical trials, (S1) 68
 - safety of treatment with a new NSAID which shows selectivity towards COX-2: a global analysis, (A1) 219
 - selective inhibition of inducible cyclooxygenase: summary of pharmacological data and clinical pharmacology, (A1) 221
 - six-month double-blind trial to compare the efficacy and safety of meloxicam 7.5 mg and naproxen 750 mg in patients with rheumatoid arthritis, (S1) 22
 - study to assess the tolerability and safety in patients with rheumatoid arthritis who have mild renal impairment, (A1) 223
 - tolerability of multiple administration of intramuscular: comparison with intramuscular piroxicam in patients with rheumatoid arthritis or osteoarthritis, (S1) 51
- MÉNDEZ MJ, AMES P, KHAMASHTA M, HUGHES GRV. Pulse cyclophosphamide in the treatment of neuropsychiatric systemic lupus erythematosus, (A1) 32
- MÉNDEZ MJ *see also* Graña J
- MENÉN-NAVARRO J *see* Moragues-Pastor C
- Meniscus: studies of proteoglycan and matrix protein synthesis and composition, (A1) 238
- MENKES CJ *see* Hilliquin P
- MENON S *see* McLure CE; Sharma A
- Mercuric-chloride-induced autoimmunity, anti-CD8 treatment reduces the severity of inflammatory arthritis, but not vasculitis, (A1) 10
- MERILÄHTI-PALO R *see* Isomäki P; Roivainen A
- MEROLA B *see* Scarpa R
- Metabolic activity tracers
- scintigraphic uptake in Paget's disease of bone, (A1) 94
 - scintigraphic uptake in Paget's disease of bone can be modified by treatment of the disease, (A1) 93
- Metalloproteinase inhibitor, synthetic, which blocks both p55 and p75 TNF receptor shedding and TNF α processing in RA synovial membrane cell cultures, (A1) 181
- Metalloproteinases, rapid RT-PCR based matrix metalloproteinase gene family screening reveals expression of several in monocytic cells, (A1) 63
- Methotrexate
- age effects on efficacy and tolerance in rheumatoid arthritis, 453
 - comparison with placebo in the treatment of systemic sclerosis: a 24 week randomized double-blind trial, followed by a 24 week observation trial, 364
 - and folate supplementation, (L) 1031
 - fore-arm bone mineral density in non-steroid treated rheumatoid arthritis is reduced in patients on long-term treatment compared with other disease-modifying drugs, (A1) 209
 - in vitro* effects on human articular cartilage and bone-derived osteoblasts, 342
 - low-dose treatment, pulmonary function in rheumatoid arthritis: a longitudinal study, 446
 - and myociclin therapies, central nervous system side effects in rheumatoid arthritis, (A1) 216
 - occurrence of pulmonary complications during therapy in rheumatoid arthritis, 441
 - and prednisolone, accidental overdose, (L) 704
 - in rheumatoid arthritis: toxicity issues, (E) 403
 - in severe ankylosing spondylitis: clinical and serological response compared with rheumatoid arthritis and psoriatic arthritis, (A1) 155
 - single intra-articular injection of liposomally conjugated suppresses joint inflammation in rat antigen-induced arthritis, 719
 - treatment every other week in patients with juvenile chronic arthritis, (L) 1030
- Methylprednisolone
- intramuscular vs oral prednisolone, controlled trial in polymyalgia rheumatica, (A1) 59
 - quantitative assessment of synovial inflammation by dynamic gadolinium-enhanced magnetic resonance imaging. A study of the effect of intra-articular methylprednisolone on the rate of early synovial enhancement, 50
- MEYER O *see* Cordonnier C; de Bandt M
- MEYER P *see* Burkhardt H
- MHC-encoded TAP genes in rheumatic diseases, (E) 501
- Michael Mason Travelling Fellowship, 1994-5: goodbye the Commonwealth, hello Australia, 1324
- MICHELS H *see* Huppertz H-I
- MIGUELEZ R *see* Sánchez-Pernaute O
- MIKECZ K *see* Brennan FR
- MILFORD-WARD A *see* Hill MB
- MILLS SY, JACOBY RK, CHACKSFIELD M, WILLOUGHBY M. Effect of a proprietary herbal medicine on the relief of chronic arthritic pain: a double-blind study, 874
- MILTENBURG AMM *see* Dolhain RJEM

- MINAUR NJ, RING F, ELVINS D, BERESFORD JN, BHALLA AK. Fore-arm bone mineral density in non-steroid treated rheumatoid arthritis is reduced in patients on long-term methotrexate compared with other disease-modifying drugs, (A1) 209
- MINGUEZ MD *see* Carreño L; González M; López-Longo FJ
- Minocycline therapy, antineutrophil cytoplasmic antibody-positive polyarthritis associated with, (L) 1327
- MIQUELÉZ R, PALACIOS I, NAVARRO F, GUTIÉRREZ S, EGIDO J, GONZÁLEZ E, HERRERO-BEAMONT G. Antiinflammatory effect of two new platelet activating factor receptor antagonists in experimental models of acute and chronic arthritis, (A1) 13
- MIRET C *see* García-Carrasco M
- MIRÓ O *see* Font C
- MISHRA MB, RYAN P, ATKINSON P, TAYLOR H, BELL J, CALVER D, FOGELMAN I, CHILD A, JACKSON G, CHAMBERS JB, GRAHAME R. Extra-articular features of benign joint hypermobility syndrome, 861
- Misoprostol efficacy in ulcer healing with continued NSAID therapy, (A1) 222
- MISRA A *see* Cox NL
- MITCHELL TJ *see* Cooke SP
- MITRA A *see* Sakkas LI
- Mixed cryoglobulinaemia
associated with hepatitis C virus, broncheolar lavage, 978
and hepatitis C virus genotypes, (A1) 50
- MIYAHARA H *see* Terato K
- MOK CC, CHAN E, KO S, LAU CS. Peripheral lymphocyte apoptosis and bcl-2 expression in systemic lupus erythematosus: correlation with disease activity, (A1) 39
- MOK CC, CHAN SY, LAU YL, CHAN D, LAU CS. Dysfunctional allele of the mannose binding protein gene in rheumatoid arthritis, (A1) 189
- MOK CC, HO C, CHAN E, LAU CS. Red cell complement receptor 1 deficiency in systemic lupus erythematosus in southern Chinese: correlation with disease activity, (A1) 26
- MOK CC. Polymyalgia rheumatica as the rheumatological manifestation of myelodysplastic syndrome in a Chinese patient, (L) 496
- MOKHTAR M *see* Malaviya AN
- MOLING O. *Chlamydia pneumoniae* reactive arthritis and persistent infection, (L) 1189
- MONIZ C *see* Dolan AL
- Monoclonal gammopathy, arthritis associated with: clinical characteristics, 241
- Monocyte activity in Behçet's disease, 424
- MONTAGUE B *see* Foley-Nolan D
- MONTAGUE S, MCGONAGLE D, ROHAN D, ORMOND P, GRENNAN S, CASEY EB. Continued low level occurrence of rheumatic fever in an Irish population, (A1) 164
- MONTAGUE S *see also* McGonagle D
- MONTEAGUDO I, LÓPEZ-LONGO FJ, MORENO AC, ABAD MA, GODOY H, RODRÍGUEZ-MAHOU M, ALONSO J, CARREÑO L. Anticardiolipin antibodies and renal disease in patients with systemic lupus erythematosus, (A1) 20
- MONTEAGUDO I *see also* Carreño L; González M; López-Longo FJ
- MONTEAGUDO J *see* Navarro M
- MONTI M *see* Ferri C
- MONTILLA C *see* Collantes E
- MOODY AR *see* Yeap SS
- MOORE JE *see* Kane D
- MOORE PM, ISENBERG DA. Autoantibodies to Lba5 and ribosomal P in NP-SLE, (A1) 111
- MOORE T *see* Anderson ME; Bukhari M; Jayanetti S
- MOORE TL *see* Anderson ME
- MOOTS R *see* Devlin J; Exley A
- MOOTS RJ *see* Luqmani RA
- MORAGUES C *see* Narváez J
- MORAGUES-PASTOR C, GONZÁLEZ-COMPTA X, VERA-LLAÓ R, NOLLA-SOLÉ JM, MENÉN-NAVARRO J, ROIG-ESCOFFET D, DICENTA-SOUSA M. Hearing loss study in Paget's disease of bone. High frequency audiometry evaluation, (A1) 92
- MORAGUES-PASTOR C, SABATÉ M, UGARTEMENDIA C, NOLLA-SOLÉ JM, GAUSÍ C, ROIG-ESCOFFET D. Echocardiographic assessment of cardiac function in Paget's disease of bone, (A1) 93
- MORAN A *see* Snowden N
- MORAN CG *see* Lingard EA
- MORANT S *see* Fenn GC
- MORANT SV *see* Walker DJ
- MORENO AC *see* González M; Monteagudo I
- MORENO I *see* De La Iglesia JL
- MORGAN AW. Report on the Hypermobility Special Interest Group, British Society for Rheumatology meeting in Glasgow, April 1995, 392
- MORGAN BP *see* Goodfellow RM
- MORLA RM *see* Obach J
- MORLEY BJ *see* Hogarth MB; Slingsby JH
- Morphine and indomethacin, synergistic interaction between attenuates pain sensation and tissue destruction in adjuvant arthritis, (A1) 75
- MORRIS AJ, LEE FD, CAPELL HA, STURROCK RD, MCKENZIE JF. Jejunal lesions in ankylosing spondylitis are due to non-steroidal anti-inflammatory drug therapy, (A1) 157
- MORRIS IM. School Christmas shows... 'look out the backache is behind you', (L) 1335
- MORRIS J *see* Pal B
- MORRIS VH, CRUWYS SC, KIDD BL. Evidence for a central component to pain in rheumatoid arthritis, (A1) 125
- MORRISON E, CAPELL H. Corticosteroids in management of rheumatoid arthritis, (E) 2
- MORRISON E, DUNCAN MR, MCENTEGART A, CAPELL HA, MADHOK R, PORTER DR, THOMSON EA. Social deprivation profile in patients commencing sulphasalazine therapy for rheumatoid arthritis 1990-1993, (A1) 226
- MORRISON E *see also* Capell HA
- MORRISON PT *see* Bavington C; Brennan FR
- MORROW WJW *see* Ng T; Ross EL
- Mortality
5 year, discrepancies in RA patients during second line therapy—cultural or procedural?, (A1) 234
in extended pedigree multicase RA families, (A2) 11
- MORTUZA FY *see* Perry MJ
- MOUG SJ *see* Lim KL
- MOUM T *see* Smedstad LM
- MOUROU M *see* Malaviya AN
- MOUSSA KMK *see* Sanders PA
- MOUTSOPOULOS HM, KORDOSSIS T. Sjögren's syndrome revisited: autoimmune epithelitis, (E) 204
- MOUTSOPOULOS HM *see also* Vlachoyiannopoulos PG
- Movement range
frequency of restricted in individuals with self-reported shoulder pain: results from a population-based survey, (A1) 132, 1137
of large joints, simple index as an outcome measure in RA, (A1) 194
- MOWAT AG *see* Williamson L
- MOYES SP, MAGEED RA, MAINI RN. Somatic mutation and gene fingerprinting of immunoglobulin lambda light chains in the synovium of patients with rheumatoid arthritis, (A1) 172
- Mseleni joint disease, (A1) 106
- MT-MMPI expression by human fibroblast-like synoviocytes and its regulation by inflammatory cytokines, (A1) 67
- MUDDU B *see* Pal B
- MUELLER C *see* Distel M
- MUIR A, PULLAR T. Combined sodium aurothiomalate and sulphasalazine patients with rheumatoid arthritis who showed a partial response to sulphasalazine alone, (A1) 224
- MUIR KR *see* O'Reilly SC
- MULCAHY B *see* Masud T; Taylor JC
- MULCAHY FM *see* Kane D
- MULERO J, LABIANO I, RAMOS J, ORTIZ-BERROCAL J, LARREA A, ANDREU JL. Role of human immunoglobulin-Tc99 scan in the assessment of joint inflammation in rheumatoid arthritis, (A1) 203
- MULERO J *see also* Andreu JL; Diez E
- MULHERIN D, FITZGERALD O, BRESNIHAN B
Mechanisms of articular destruction in the feet and hands may differ, (A1) 192
Walking time may be a very useful measure of clinical laboratory and radiological status in patients with rheumatoid arthritis, (A1) 195
- MULHERIN D, FITZGERALD O, BRESNIHAN B. Clinical improvement and radiological deterioration in rheumatoid arthritis: evidence that the pathogenesis of synovial inflammation and articular erosion may differ, 1263
- MULLANEY P, COLLUM L, KILFEATHER SA, FOLEY-NOLAN D. Effect of a pulsed 27 MHz electromagnetic field on fibroblast cell division cyclic AMP and inositol phosphate, (A1) 64
- Multidisciplinary treatment, in-patient, versus routine out-patient care, randomized clinical trial in active rheumatoid arthritis, 475
- Multidrug resistance glycoprotein 170 expression in peripheral blood lymphocytes of rheumatoid arthritis patients, 430
- Multiple dysplasia in ochronosis: a non-rheumatological series of 16 patients, (A1) 95
- MUÑOZ E *see* Císal A
- MUÑOZ FJ *see* García-Carrasco M
- MUÑOZ-FERNÁNDEZ S. Antiperinuclear factor in spondylarthropathies, (L) 804
- MUÑOZ-FERNÁNDEZ S, FERREIRA A, FONTÁN G, MARTÍN MOLA E, GUON BAÑOS J. Soluble HLA class I antigens in serum and synovial fluid from patients with rheumatoid arthritis and other arthropathies, (A1) 176
- MUNRO R, DUNCAN M, CAPELL HA, JOHNSTON R. Unusual complications of cervical spine surgery for cervical myelopathy in rheumatoid arthritis, 682
- MUNRO R *see also* Capell HA; Houston T; Porter DR

- MURPHY A see Donnelly S
 MURPHY G see Zhang Z
 MURRAY R, ENGLAND AJ, REID R, REID DM. Value of high mineral content water in the short-term management of rheumatoid arthritis, (A1) 228
 Muscle biopsies, audit, (A1) 128
 Muscle fibre size
 and distribution of the back extensors: relationship with muscle fatigability, (A1) 144
 and type distribution, normal of the human back extensors, (A1) 144
 Musculoskeletal care package, (A1) 126
 MUSIKIC P see Jevtic V; Poleksic L
 Mutational analysis of the membrane proximal cleavage site of human p75 tumour necrosis factor receptor, (A1) 109
Mycobacterium xenopi—an unusual presentation as tenosynovitis of the wrist in an immunocompetent patient, 1008
 Myelodysplastic syndrome, polymyalgia rheumatica as the rheumatological manifestation in a Chinese patient, (L) 496
 Myelopathy functional disability index—a step towards objectivity, (A2) 19
 MYERSCOUGH A see John S
 MYKKÄNEN H see Malin M
 MYKLEBUST G, GRAN JT. Prospective study of 287 patients with polymyalgia rheumatica and temporal arteritis, 1161
 Myocardial infarction, silent in Wegener's granulomatosis, 188
 Myocrisin and methotrexate therapies, central nervous system side effects in rheumatoid arthritis, (A1) 216
 Myopathies, idiopathic inflammatory, distinctive features in French Canadians, (A1) 49
 Myositis, idiopathic versus lupus: comparison of clinical and laboratory features, (A1) 48
 MYSZOR MF see Turner HE

 Na/K-ATPase inhibition exerts differential effects on cytokine production by monocytes and fibroblasts, (A1) 67
 NADH oxidase activity in inflammatory synovitis measured by chemiluminescence, (A1) 69
 NAGANT DE DEUCHAISNES C see Houssiau FA
 NAHIT ES, MACFARLANE GJ, CROFT PR, COOPER C, HOSIE G, SILMAN AJ. Patients presenting with hip pain in the community: comparing those with and without radiographic changes of osteoarthritis, (A1) 102
 Nailfold capillaroscopy
 economic study in Raynaud's phenomenon: follow-up and evaluation as prognostic value, (A1) 49
 video and thermal stress testing in children with Raynaud's phenomenon, (A2) 26
 Nailfold capillary dimensions, increased in primary Raynaud's phenomenon and systemic sclerosis, 1127
 NAMBA H see Okano K
 NANDRA D, HART DJ, DOYLE DV, SPECTOR TD. Is hormone replacement therapy protective for osteoarthritis in women: the Chingford Study, (A1) 124
 NANDRA D, HART DJ, MANZI C, GALLIMORE JR, MACKILLOP N, DOYLE DV, PEPYS MB, SPECTOR TD. Low level increases in serum CRP are present in early osteoarthritis of the knee and predict progressive disease, (A1) 99
 NANDRA D see also Masud T
 Naproxen, six-month double-blind trial to compare the efficacy and safety of meloxicam 7.5 mg and naproxen 750 mg in patients with rheumatoid arthritis, (S1) 22
 NARANJO-HERNÁNDEZ A. Chronic polyarthritis and graft-versus-host disease, (L) 297
 NARVÁEZ FJ see Valverde J
 NARVÁEZ J, RODRIGUEZ-MORENO J, MORAGUES C, CAMPOY E, CLAYAGUERA T, ROIG-ESCOFET D. Tertiary hyperparathyroidism after long-term phosphate supplementation in adult-onset hypophosphataemic osteomalacia, 598
 NARVÁEZ J. Subarachnoid haemorrhage secondary to ruptured cerebral aneurysm in a patient with osteogenesis imperfecta, (L) 1332
 NASH EJ see Pal B
 Natural resistance associated macrophage protein locus, linkage and association studies in rheumatoid arthritis, (A1) 188
 NAUGHTON DP, GAFFNEY K, GROOTVELD MC, RHODES CJ, BLAKE DR, NAZHAT NB, SYMONS MCR. Electron paramagnetic resonance study of the arthritic joint, (A1) 204
 NAUGHTON M, BOTTO M, CARTER M, ALEXANDER G, GOLDMAN J, WALPORT MJ. Extrahepatic secreted complement C3 contributes to circulating C3 levels in humans, (A1) 66
 NAVARRO E see Olivé A
 NAVARRO F see Miguélez R; Sánchez-Pernaute O
 NAVARRO FJ, SEMPÉRIZ JM, TOVAR JV. Modification of the inflammatory activity of psoriatic arthritis in patients treated with extract of *Polipodium leucotomos* (Anapsos), (A1) 161
 NAVARRO JF. New NSAIDs—inhibitors of cyclooxygenase 2—and renal damage, (L) 1332
 NAVARRO M, CERVERA R, TEIXIDÓ M, REVERTER JC, FONT J, LÓPEZ-SOTO A, MONTEAGUDO J, ESCOLAR G, INGELMO M. Antibodies to endothelial cells and to β_2 -glycoprotein I in the antiphospholipid syndrome: prevalence and isotype distribution, 523
 NAVARRO MA see Gomez-Vaquero C
 NAVES M see Hajeer AH
 NAZHAT NB, SAADALLA-NAZHAT RA, GAFFNEY K, WINYARD P, BLAKE DR. Increased catalase activity in synovial fluid of rheumatoid arthritis patients compared to osteoarthritis patients, (A1) 70
 NAZHAT NB see also Naughton DP
 Needle arthroscopy service, audit, (A1) 127
 NEILD G see Devi B
 Nephropathy, rheumatoid, endothelial damage is an important mechanism in the pathogenesis, (A1) 170
 Nerve antigens, antibodies to in rheumatoid peripheral neuropathy, (A1) 171
 Neurofibromatosis masquerading as pauciarticular juvenile chronic arthritis, (A2) 39
 Neurological examination, quantified with emphasis on motor and sensory functions in patients with rheumatoid arthritis and controls, 1116
 Neurological manifestations in Saudi patients with systemic lupus erythematosus, (A1) 32
 Neurological signs in RA patients receiving gold, (L) 803
 Neuropeptides stimulate *in vitro* cytokine release in human leucocytes in rheumatoid arthritis patients, (A1) 8
 Neutrophil
 adhesion to endothelial cells is a signal for loss of intracellular lipocortin-1, (A1) 60
 and interleukin-8 function of synovial fluid in rheumatoid arthritis and seronegative polyarthritis, 1244
 tyrosine phosphorylation in synovial fluid of patients with rheumatoid arthritis, 846
 NEVE R see Chernajovsky Y
 NEWKIRK MM, SHIROKY JB, JOHNSON N, DANOFF D, ISENBERG DA, SHUSTIK C, PEARSON GR. Rheumatic disease patients, prone to Sjögren's syndrome and/or lymphoma, mount an antibody response to BHRF1, the Epstein-Barr viral homologue of BCL-2, 1075
 NEWMAN J, SILMAN AJ. Comparison of disease status in rheumatoid arthritis patients attending and not attending a specialist clinic, 1169
 NEWMAN SP see Shortall E
 NF- κ B inhibition, novel action of glucocorticoids, (E) 709
 NG T, COLLINS I, D'CRUZ D, KANNER S, HUMPHRIES M, NYE K, MORROW WJW. Integrin-induced T-cell signaling in systemic lupus erythematosus: a novel tool for disease monitoring, (A1) 38
 NG TS see Hurley MV
 NICAISE P see Cordonnier C
 NICHOL FE see Evans MD
 NICOD A see Hardwick L; Hardwick LJ
 NIKKARI S. Parvovirus B19 in rheumatoid arthritis, (L) 494
 NIKKARI S see also Isomäki P
 NISHIKAI M, AKIYA K, TOJO T, ONODA N, TANI M, SHIMIZU K. Seronegative Sjögren's syndrome manifested as a subset of chronic fatigue syndrome, 471
 NISHIOKA J see Tiwana H
 Nitration of tyrosine in the inflamed human synovium: evidence for the generation of peroxynitrite *in vivo*, (A1) 70
 Nitric oxide
 local production, synovial fluid from knees with secondary osteoarthritis has much higher nitrite concentrations than serum, (A1) 105
 production in cells derived from the human joint, 207
 Nitric oxide synthase, inducible, immunolocalisation in rheumatoid synovium, (A1) 177
 NIXON NB see Salih AM; Scott LJ
 NODA B see Hdez-Berain JA
 NOLLA JM see Fiter J
 NOLLA-SOLÉ JM see Moragues-Pastor C
 Non-steroidal anti-inflammatory drugs (NSAIDs)
 compliance monitoring in ankylosing spondylitis, experiences with an electronic monitoring device, 60
 efficacy of misoprostol therapy in ulcer healing, (A1) 222
 gastrointestinal bleeding, incidence and severity, (A1) 220
 high intestinal permeability in patients with inflammatory arthritis is more strongly associated with than with other factors, (A1) 221
 infertility may sometimes be associated with consumption, 76
 influence on the evolution of abdominal scintigraphy in patients with spondylarthropathies, (A1) 154
 jejunal lesions in ankylosing spondylitis are due to therapy, (A1) 157
 mechanism of action, (S1) 1
 new, safety of treatment with meloxicam, which shows selectivity towards COX-2: a global analysis, (A1) 219
 new—inhibitors of cyclooxygenase 2—and renal damage, (L) 1332
 oesophagitis from is not caused by pill oesophagitis, (A1) 225

- pharmacology of meloxicam—a new non-steroidal anti-inflammatory drug with an improved safety profile through preferential inhibition of COX-2, (A1) 4
- related serious gastrointestinal bleeding, survey of NSAID usage in hospital and community, (A1) 218
- reversible ovulatory failure associated with development of luteinized unruptured follicles in women with inflammatory arthritis, 458
- survey of usage in hospital and community and serious NSAID related gastrointestinal bleeding, (A1) 218
- NOON E *see* Knight J
- NORSWORTHY P *see* Athanassiou P
- NORSWORTHY PJ *see* Slingsby JH
- NORTH N *see* Wojtulewski JA
- NUKI G *see* Bavington C; Brennan FR; Jobanputra P; Richmond R; Smith G
- NYE K *see* Ng T
- OBACH J, MORLÀ RM, ESCOLÀ A, PÉREZ A, PLANS P, ASENSI E. Risk factors of hyperuricaemia in Catalonia-II survey, (A1) 146
- OBACH J *see also* Escolà A
- Obesity, affecting knee pain and disability in the community, (A1) 134
- O'BRIAN S *see* Kane D
- Obstetric outcome in systemic lupus erythematosus and antiphospholipid syndrome: the St Thomas' Hospital Lupus in Pregnancy clinic experience, (A1) 123
- ÓCAL L, RUSSELL K, BEYNON H, CRUICKSHANK K, LANCHBURY JS, WALPORT M, ISENBERG DA, BRIGGS D. Genetic analysis of TAP2 in systemic lupus erythematosus patients from two ethnic groups, 529
- Occupational lifting and osteoarthritis of the hip, (A1) 100
- Occupational therapists and physiotherapists, evaluation of extended role in rheumatology practice, (A1) 130
- Occupational therapy and physiotherapy for juvenile chronic arthritis: custom and practice in five centres in the UK, USA and Canada, 695
- Ochronosis, multiple dysplasia in: a non-rheumatological series of 16 patients, (A1) 95
- Ochronotic arthropathy: a non-rheumatological series of 16 patients, (A1) 95
- O'CONNELL JP *see* Zhang Z
- O'CONNOR KBR *see* Plant MJ
- Ocular involvement in giant cell arteritis: study of 191 patients in 15 years, (A1) 57
- ODDING E, VALKENBURG HA, ALGRA D, VANDENOUWELAND FA, GROBBEE DE, HOFMAN A. Association of abnormalities on physical examination of the hip and knee with locomotor disability in the Rotterdam study, 884
- Oesophageal dysmotility, risk factors for progression in systemic sclerosis, (A1) 44
- Oesophagitis from NSAIDs is not caused by pill oesophagitis, (A1) 225
- Oestrogen, postmenopausal use and osteoarthritis of the hip, (A1) 101
- Oestrogen receptor gene polymorphism and bone mass in pre- and postmenopausal women, (A1) 89
- Oestrogen-gestagen therapy, aminohehexane bisphosphonate suppresses bone turnover in postmenopausal women more rapidly, 636
- O'FARRELLY C *see* Costello P
- OHISHI Y *see* Furukawa M
- OHTSURU A *see* Okano K
- OJANGUREN I *see* Olivé A
- OKANO K, TSUKAZAKI T, OHTSURU A, NAMBA H, OSAKI M, IWASAKI K, YAMASHITA S. Parathyroid hormone-related peptide in synovial fluid and disease activity of rheumatoid arthritis, 1056
- OKUDA Y *see* Kobayashi H
- OLAH AJ *see* Lehmann T
- OLAJOS A. Value of HLA-B27 typing in the diagnosis of early, oligosymptomatic spondylarthropathies, (C) 192
- Oligoarthritis
- prediction of diagnosis in acute and subacute oligoarthritis of unknown origin, 359
- undifferentiated, search for infective agents in Papua New Guinea, (L) 492
- OLIVÉ A, ESTEVE M, NAVARRO E, KLAASSEN J, CABRÉ E, TENA X, FERNANDEZ F, PASTOR C, GASULL MA. Plasma fatty acid pattern in rheumatoid arthritis, (A1) 191
- OLIVÉ A, PÉREZ-ANDRÉS R, TENA X, GUASCH I, ROMEU J, CLOTET B. Musculoskeletal manifestations of HIV infection, (A1) 167
- OLIVÉ A, TENA X, RAVENTOS A, ROMEU J, LORENZO JC, OJANGUREN I, PÉREZ R. Bone bacillary angiomatosis in an HIV-infected patient, 901
- OLIVÉ A *see also* Acebes JC
- OLIVEIRA DBG *see* Kiely PDW
- OLIVEIRA M *see* Ciconelli RM
- OLIVER C, WATT I. Intravenous MRI contrast enhancement of inflammatory synovium: a dose study, (S3) 31
- OLIVIERI I
- Osteitis condensans ilii, (L) 295
- Undifferentiated seronegative spondylarthropathy in females, (L) 395
- OLLIER B *see* Brennan P
- OLLIER W *see* Hajeer A; Haworth S; John S; Marlow A
- OLLIER WER *see* Davies EJ; Donn RP; Hajeer AH; John SL; McDonagh JE; Plant MJ; Snowden N; Stanworth SJ
- OLSSON AT *see* Elling P
- Oncoprotein expression in human synovial tissue: an immunohistochemical study of different types of arthritis, 933
- Oncostatin M is raised in rheumatoid synovial fluids and interacts with IL-1 to promote cartilage collagen destruction *in vitro*, (A1) 109
- O'NEILL TW, GRAZIO S, SPECTOR TD, SILMAN AJ. Geometric measurements of the proximal femur in UK women: secular increase between the late 1950s and early 1990s, (A1) 90
- O'NEILL TW, MCCLOSKEY EV, REEVE J, REID DM, TODD C, WOOLF AD, KANIS A, SILMAN AJ. Relationship between vertebral osteophytosis and back pain, (A1) 143
- O'NEILL TW, VARLOW J, SILMAN AJ, FELSENBURG D, COOPER C, KANIS JA, European Vertebral Osteoporosis Study Group
- Effect of physical activity on the risk of vertebral deformity, (A1) 89
- Hormonal and gynaecological factors and risk of vertebral deformity, (A1) 90
- ONODA N *see* Nishikai M
- OOI L *see* Wang CL
- Opioid tone, endogenous in patients with rheumatoid arthritis, 436
- OPPENHEIM B *see* Pal B
- ORDI J *see* Davies EJ
- O'REILLY S, JONES A, MUIR KR, DOHERTY. Quadriceps strength and depression are important factors in knee pain and associated disability, (A1) 135
- O'REILLY S, MUIR KR, DOHERTY M. Factors affecting knee pain and disability in the community: the importance of obesity and psychological factors, (A1) 134
- Organ transplantation, new roles in amyloidosis, (A1) 12
- ORIENTE P *see* Lubrano E; Scarpa R
- ORIOL A *see* Acebes JC
- ORMOND P, MCGONAGLE D, BERGIN C, TODD A, CASEY EB. Removal of patients from rheumatology waiting list, (A1) 129
- ORMOND P *see also* McGonagle D; Montague S
- Orthopaedic medicine, the remit of the rheumatologist, (E) 1
- ORTIZ J *see* Isasi C; Perez A
- ORTIZ-BERROCAL J *see* Andreu JL; Mulero J
- OSAKI M *see* Okano K
- OSBORNE D *see* Sharif M
- OSBORNE DJ *see* Sharif M
- Ossification, heterotopic complicating long-term sedation, (L) 700
- Osteitis condensans ilii, (L) 295
- OSTENDORF B, SPECKER Ch, SCHNEIDER M. Methotrexate in severe ankylosing spondylitis: clinical and serological response compared with rheumatoid arthritis and psoriatic arthritis, (A1) 155
- Osteoarthritic cartilage, increased expression of the Ed-B-containing fibronectin (an embryonic isoform of fibronectin), 407
- Osteoarthritic synovium induces an up-regulation of p55 TNF α receptor expression on non-arthritis chondrocytes, (A1) 237
- Osteoarthritis
- actual repair of cartilage by Ilizarov joint distraction tested in the Pond-Nuki dog model, (A1) 104
- assessment of urinary hydroxypyridinium cross-links measurement, 1091
- C-reactive protein is elevated in, (A1) 99
- disruption of the IGF system in osteoarthritic and rheumatoid patients compared to non-arthritis volunteers, (A1) 105
- effects of total knee replacements in patients with rheumatoid versus osteoarthritis, (A2) 32
- generalised in women: pattern of joint involvement and approach to definition, (A1) 97
- genetic influences: a study of twins, (A1) 116
- hand, hip and knee, risk factors in Catalonia-II survey, (A1) 101
- of the hands in early populations, 1291
- of the hip, double-blind study to compare efficacy and safety of meloxicam 15 mg with piroxicam 20 mg, (S1) 35
- hip
- factors affecting the outcome, (A1) 102
- and knee, familial tendency to severe symptomatic, (A1) 100
- occupational lifting, (A1) 100
- postmenopausal oestrogen use, (A1) 101
- subchondral sclerosis and fractal signature analysis, (A1) 103
- IL-12 expression and quantitation by competitive PCR in the synovium, (A1) 180
- increased catalase activity in synovial fluid of patients compared to rheumatoid arthritis patients, (A1) 70
- increased expression of interferon (IFN)-gamma together with IFN-gamma receptor in the synovium of patients with osteoarthritis compared with the rheumatoid synovial membrane, 24
- is hormone replacement therapy protective in women: the Chingford Study, (A1) 124

- knee
 hospital-referred, symmetry of radiographic change, (A1) 97
 laxity, 560
 low level increases in serum CRP are present and predict progressive disease, (A1) 99
 pain amongst the poor and affluent in Pakistan, 146
 proposed scoring system for assessing synovial membrane abnormalities at arthroscopy, (S3) 15
 lower limb, multidisciplinary study of outpatient education, (A1) 103
 meloxicam in: a six-month, double-blind comparison with diclofenac sodium, (S1) 39
 and osteoporotic fracture association: the Chingford study, 1299
 papain-induced, low energy laser effect, (A2) 7
 patellofemoral, skyline are better than lateral radiographs for detecting progression, (A1) 98
 patients presenting with hip pain in the community: comparing those with and without radiographic changes, (A1) 102
 and polymorphisms of the vitamin D receptor, (A1) 117
 posterior facet joint, lumbar spine bone density measurements: the effect in a middle age female population, (A1) 98
 radiographic, association with osteoporotic fractures: the Chingford study, (A1) 116
 and rheumatoid arthritis
 clinical and laboratory assessments, (S3) 6
 difference in expression of the plasminogen activation system in synovial tissue, 416
 and rheumatoid disease, comparison of the clinical features of hospital out-patients in Pakistan and England, 994
 serum YKL-40 levels in healthy children and adults. Comparison with serum and synovial fluid levels of YKL-40 in patients with osteoarthritis or trauma of the knee joint, 553
 spontaneous in guinea pigs: cartilage degeneration is preceded by loss of subchondral trabecular bone, (A1) 104
 tolerability of multiple administration of intramuscular meloxicam: comparison with intramuscular piroxicam, (S1) 51
 Osteoarthritis-osteoporosis, inverse relationship: what is the evidence? What are the consequences?, (E) 813
 Osteoarthritis, secondary, synovial fluid from knees has much higher nitrite concentrations than serum, suggesting local production of nitric oxide, (A1) 105
 Osteoarticular tuberculosis in two different populations: the HIV infection impact, (A1) 166
 Osteoblastic bone nodules and bone morphogenetic proteins are induced bone growth stimulating pulsed electromagnetic fields, (A1) 115
 Osteoblasts, bone-derived, *in vitro* effects of methotrexate, 342
 Osteocalcin levels and calciotropic hormones in lung transplant recipients, (A1) 86
 Osteogenesis imperfecta, subarachnoid haemorrhage secondary to ruptured cerebral aneurysm, (L) 1332
 Osteolytic properties of the synovial-like tissue from aseptically failed joint prostheses, 943
 Osteomalacia, adult-onset hypophosphataemic, tertiary hyperparathyroidism after long-term phosphate supplementation, 598
 Osteomyelitis
 chronic recurrent multifocal: spinal involvement and radiological appearance, 1019
Escherichia coli of the clavicle, chronic case, (C) 192
 Osteophytosis, vertebral, relationship with back pain, (A1) 143
 Osteoporosis
 audit of guidelines for treatment and prevention, (A1) 81
 and bone morbidity in lung transplant recipients, (A1) 85
 chronic hypophosphataemia as possible risk-factor, (A1) 96
 incidence in patients attending fracture clinic, (A2) 6
 inverse relationship with osteoarthritis, what is the evidence? What are the consequences?, (E) 813
 management of related fractures of femoral neck and vertebrae: survey of orthopaedic surgeons' practice, (A1) 82
 postmenopausal, morphometric X-ray absorptiometry and radiographic vertebral morphometry, (A1) 83
 scurvy and megaloblastic anaemia due to alleged food intolerance, (L) 701
 severe due to systemic mast cell disease: successful treatment with interferon alpha-2b, 898
 timed therapy with calcium and phosphate supplements restores phosphate and parathyroid hormone rhythms in patients with osteoporosis, with an anabolic effect on spinal bone mineral density, (A1) 88
 Osteoporotic fracture
 association with radiographic osteoarthritis: the Chingford study, (A1) 116
 and osteoarthritis association: the Chingford study, 1299
- ØSTERGAARD M, HANSEN M, STOLTENBERG M, LORENZEN I. Quantitative assessment of the synovial membrane in the rheumatoid wrist: an easily obtained MRI score reflects the synovial volume, 965
 ØSTERGAARD M, STOLTENBERG M, HENRIKSEN O, LORENZEN I. Quantitative assessment of synovial inflammation by dynamic gadolinium-enhanced magnetic resonance imaging. A study of the effect of intra-articular methylprednisolone on the rate of early synovial enhancement, 50
 O'SULLIVAN MM see Martin JC
 ÖTÜRK C see Gül A
 Out-patient care versus in-patient multidisciplinary treatment, randomized clinical trial in active rheumatoid arthritis, 475
 Out-patient clinic, comparison with a community clinic in rheumatology, 385
 Out-patient education, multidisciplinary study in patients with lower limb osteoarthritis, (A1) 103
 Outcome
 and disease activity measurement in rheumatoid arthritis using the Nottingham Health Profile, (A1) 231
 measure, video image analysis of hands: development of an anatomic index in rheumatoid arthritis, 1274
 measures in rheumatoid arthritis: relationship with disease activity and anatomic damage, (A1) 231
 measuring in rheumatoid arthritis—which measures are suitable for routine clinical use, (L) 301
 prediction in 537 patients with newly diagnosed rheumatoid arthritis examining the effect of age at onset and gender, (A1) 233
 OVER K see MacDonald AG
 OVER KE, STEVENSON A, YAQOUB M, MACDONALD AG, MASON H, BUCKNALL RC. Renal tubular dysfunction is associated with endothelial damage in systemic sclerosis, (A1) 44
 Overdose, accidental of prednisolone and methotrexate, (L) 704
 Ovulation-induction therapy, risks in systemic lupus erythematosus, 1184
 Ovulatory failure, reversible, associated with development of luteinized unruptured follicles in women with inflammatory arthritis taking non-steroidal anti-inflammatory drugs, 458
 ÖVYAZGAN Y see Yazici H
 OXTORY J see Avers HL
 ÖZBEK U see Gül A
 ÖZÇELİK T see Gül A
- p55 TNF α receptor expression, osteoarthritic synovium induces an up-regulation on non-arthritis chondrocytes, (A1) 237
 PACE J see Matucci Cernic M
 Paget's disease
 in a patient with ankylosing spondylitis—a diagnostic dilemma, 1011
 use of the alkaline phosphatase half-life in the comparison of potency between bisphosphonate compounds in treatment, (A1) 92
 Paget's disease of bone
 echocardiographic assessment of cardiac function, (A1) 93
 hearing loss study. High frequency audiometry evaluation, (A1) 92
 scintigraphic uptake of metabolic activity tracers, (A1) 94
 scintigraphic uptake of metabolic activity tracers bone can be modified by treatment of the disease, (A1) 93
 Pain
 chronic arthritic, effect of a proprietary herbal medicine on relief: a double-blind study, 874
 chronic widespread: is a better classification possible?, (A1) 136
 chronic widespread in the community, follow-up study: a better prognosis than in the clinic, (A1) 137
 coping strategies as predictors for graded back pain status: results of a population based follow-up study, (A1) 142
 evidence for a central component in rheumatoid arthritis, (A1) 125
 relevance of daytime and nocturnal measurements in rheumatoid arthritis, (A1) 193
 why does an inflammation in the joint hurt?, (E) 405
 Painful shoulder: can consultants agree?, 1172
 PAIRET M see Frölich JC
 PAL B. Audit of domiciliary visits for locomotor problems, (A2) 30
 PAL B, MORRIS J
 Alternative/complementary medicine, a survey of the attitudes of rheumatologists and private medical insurers, (A1) 128
 Survey of ambulatory patients' attitudes towards a proposed telephone follow-up service, (A1) 130
 PAL B, MORRIS J, KEENAN J. Management of idiopathic carpal tunnel syndrome: survey of rheumatologists, (A2) 34
 PAL B, MUDDU B, MORRIS J. Management of osteoporosis related fractures of femoral neck and vertebrae: survey of orthopaedic surgeons' practice, (A1) 82
 PAL B, NASH EJ, OPPENHEIM B, MAXWELL S, MCFARLANE L. Routine synovial fluid culture: is it necessary? Lessons from an audit, (A2) 34
 PALACIOS I, SÁNCHEZ-PERNAUTE O, HERNÁNDEZ P, LÓPEZ-ARMADA MJ, GUTIERREZ S, GONZÁLEZ E, EGIDO J, HERRERO-BEAUMONT G. Tenidap reduces cell infiltration and MCP-1 mRNA expression in a rabbit model of acute arthritis, (A1) 73

- PALACIOS I *see also* Gutierrez S; Miguélez R; Sánchez-Pernaute O
 PALAZZO E *see* Cordonnier C
 PALEOLOG E, TAYLOR P, HUNT M, ELLIOTT M, FELDMAN M, TAK P, BREEDVELD F, WOODY J, MAINI R. Treatment of rheumatoid arthritis with antibody to TNF α decreases expression and shedding of E-selectin, (A1) 218
 PALEOLOG EM. Angiogenesis: a critical process in the pathogenesis of RA—a role for VEGF, (E) 917
 PALOMBO-KINNE E *see* Burkhardt H
 PANAYI G *see* Yanni G
 PANAYI GS *see* Baerwald CGO; Biagini P; Bowman SJ; Campbell F; Choy EHS; Dasgupta B; Dolan AL; Iannone F; Kassimos D; Pipitone N
 Pancreatic function in systemic sclerosis, (L) 298
 PANDE I *see* Koh WH
 PANTOJA L *see* De Miguel E
 PAOLOZZI L *see* Ravaut P
 PAPAGEORGIOU AC *see* Macfarlane GJ
 Parathyroid hormone-related peptide in synovial fluid and disease activity of rheumatoid arthritis, 1056
 PARKIN A *see* Hackett J
 PARNABY-PRICE A *see* Vakalis AN
 PAROLI M. Cardiac abnormalities in a patient with localized scleroderma, (L) 703
 Parotid gland ultrasonography as a diagnostic tool in primary Sjögren's syndrome, 972
 PARSONS M *see* Patel YI
 Parvovirus B19
 infection, low incidence of arthritis in immunocompromised HIV-infected patients, (A1) 167
 in rheumatoid arthritis, (L) 494
 PASCUAL C *see* Rodriguez R
 PASCUAL E. Diagnosis of gout and CPPD crystal arthropathy, (E) 306
 PASCUAL E *see also* Román JA
 PASCUAL-SALCEDO D *see* Uson J
 PASERO G *see* Ferraccioli GF; Ferri C
 PASTOR C *see* Olivé A
 PATEL S, CHAN JK-M, HOSKING DJ. Fluoride pharmacokinetics and changes in lumbar spine and femoral neck bone mineral density, (A1) 84
 PATEL S, COUPLAND CA, STONE M, HOSKING DJ. Use of the alkaline phosphatase half-life in the comparison of potency between bisphosphonate compounds in the treatment of Paget's disease, (A1) 92
 PATEL Y *see* Mackay K
 PATEL YI, PARSONS M, WHYTE J, MCHUGH NJ, SANSOM DM. Polymorphism of the extracellular domain of the human Fas gene and its relationship with SLE, (A1) 25
 Patient
 activity objective measurement before and after steroid injection of the knee in rheumatoid arthritis, (A1) 208
 education: treatment or nice extra. The British experience, (L) 805
 satisfaction and quality of care, 382
 Patient-centred approach to evaluation and treatment in rheumatoid arthritis: the development of a clinical tool to measure patient-perceived handicap, 921
 PATO E *see* Fernández-Dapica P
 PATOJA L, SANTUCCI L, FURNO P, DIONISI MS, DELL'ORSO S, ROMAGNOLI M, SATTARINIA A, MARINI MG. 4-week, double-blind, parallel-group study to compare gastrointestinal effects of meloxicam 7.5 mg and 15 mg, piroxicam 20 mg and placebo, (S1) 61
 PATTON D, FREEDMAN J, YOUNG A. Audit of guidelines for treatment and prevention of osteoporosis, (A1) 81
 PAULY T *see* Wedekind F
 PAUNET M *see* Lioté F
 PAZIANIS M *see* Tobias JH
 PCR, multi-bacterial nested PCR, use to screen synovial fluid and membrane samples of patients with inflammatory arthritis and sacroiliitis, (A1) 113
 PEARCE GJ *see* Slingsby JH
 PEARCE S *see* Pincus T
 PEARSON D *see* Yeap SS
 PEARSON GR *see* Newkirk MM
 PEARSON JD *see* Denton CP
 PEDRO-BOTET J. Pericardial effusion and giant cell arteritis, (L) 194
 PEDROL E *see* Font C
 PEEL NFA, BARRINGTON NA, AUSTIN CA, EASTELL R. Paget's disease in a patient with ankylosing spondylitis—a diagnostic dilemma, 1011
 PEGORARO I *see* Ferraccioli GF
 PELLETIER J-P *see* Martel-Pelletier J-P
 PEÑA F *see* Andreu JL
 PEÑA JL *see* Calvo-Alén J; Sánchez-Andrada S
 D-Penicillamine
 induced myopathy: reductions in total muscle RNA and copper, (A1) 72
 reductions in the synthesis rates of skeletal muscle proteins in treated rats, (A1) 71
 PENNINGTON DC *see* Barlow JH
 PEPPER L *see* Davies EJ; Haworth S
 PEPYS MB *see* Hawkins PN; Nandra D; Persey MR
 PEREIRA FP *see* Carbon RJ
 PEREZ A *see* Isasi C
 PEREZ A *see* Obach J; Valverde J
 PEREZ A, TABUENCA MJ, ISASI C, TORRESANO M, ESCRIBANO M, LARREA A, ORTIZ J. Scintigraphic uptake of metabolic activity tracers in Paget's disease of bone can be modified by treatment of the disease, (A1) 93
 PÉREZ G *see* Collantes E
 PÉREZ J *see* Sanchez J
 PEREZ R *see* Olivé A
 PEREZ S *see* Román JA
 PÉREZ-ANDRÉS R *see* Olivé A
 PÉREZ-PÁRAMO M *see* Andreu JL
 PEREZ-RUIZ F, ALONSO RUIZ A, CALABOZO RALUY M, RUIZ-LUCEA E, ARETXABALA I, FRAIX E. Comparative study of the efficacy of benzbromarone and allopurinol for the therapy of hyperuricemia in chronic primary gout, (A1) 145
 PEREZ-RUIZ F *see also* Calabozo M
 PÉREZ-VILCHEZ D *see* Albornoz J; De La Iglesia JL
 Periarthritis, calcific involving multiple sites. A familial condition, (A1) 133
 Pericardial effusion
 and giant cell arteritis, (L) 194
 massive in scleroderma, (L) 807
 Perichondrium and cartilage implanted in the joint, (A1) 240
 Peripheral blood mononuclear cells from RA patients, susceptibility to peroxide-induced DNA damage, (A1) 184
 Peripheral joint disease
 immunohistologic analysis in ankylosing spondylitis, (A1) 154
 in psoriatic arthritis: a five year clinical follow-up study, (A1) 158
 Peripheral lymphocyte apoptosis and bcl-2 expression in systemic lupus erythematosus: correlation with disease activity, (A1) 39
 Peripheral neuropathy
 complicating rheumatoid arthritis, anti-ganglioside antibodies, 725
 rheumatoid
 antibodies to nerve antigens, (A1) 171
 sera from patients recognise a common antigen in neuroblastoma cells, (A1) 170
 PERKINS P *see* Devlin J
 Permanent work disability prediction in a follow-up study of early rheumatoid arthritis: results of a tree structured analysis using RECPAM, 652
 Peroxynitrite, evidence for *in vivo* generation, nitration of tyrosine in the inflamed human synovium, (A1) 70
 PERRETT D *see* Evans MD; James IT; Walne AJ
 PERRETTI M, GOULDING NJ, WHEELER SK, FLOWER RJ. Neutrophil adhesion to endothelial cells is a signal for loss of intracellular lipocortin-1, (A1) 60
 PERROT S *see* Hilliquin P
 PERRY JD *see* Carbon RJ
 PERRY LA *see* Hall GM
 PERRY MJ, PONSFORD FM, MORTUZA FY, LEARMONTH ID, ATKINS RM, ELSON CJ. Osteolytic properties of the synovial-like tissue from aseptically failed joint prostheses, 943
 PERSEY MR, LOVAT LB, APPERLEY JF, MADHOO S, PEPYS MB, HAWKINS PN. Intensive chemotherapy for AL amyloidosis, (A1) 12
 PERSEY MR, LOVAT LB, MADHOO S, GALLIMORE JR, PEPYS MB, HAWKINS PN. Dynamics of AL amyloidosis in RA and JCA, (A1) 76
 PERSEY MR, LOVAT LB, MADHOO S, PEPYS MB, HAWKINS PN. Adrenal function in AL amyloidosis, (A1) 76
 PERSEY MR *see also* Hawkins PN
 PERULA L *see* Gonzalez-Dominguez J
 PERVEEN-GILL Z *see* Harbuz MS
 PESCI A *see* Manganelli P
 PETERA P *see* Studnicka-Benke A
 PETERFY CG. MRI in the assessment of synovium and cartilage, (S3) 3
 PETERFY CG *see also* Demars F
 PETERS AM *see* Chapman PT
 PETTERSSON T, LAASONEN L, LEIRISALO-REPO M, TERVAHARTIALA P. Spinal pseudoarthrosis complicating ankylosing spondylitis: a report of two patients, 1319
 PETTY RE *see* Foster HE
 PGH-synthase-2, linoleic acid is a better substrate for than for PGH-synthase-1 tested in whole endothelial cells in culture, (A1) 219
 Phenytoin-induced pseudo-Sjögren's syndrome, (L) 1033
 PHIPPS JL *see* Hill MB
 Phosphatidyl serine-dependent antithrombin antibody is exclusive to patients with lupus anticoagulant, 589
 Physical activity effect on the risk of vertebral deformity, (A1) 89
 Physical examination, association of abnormalities of the hip and knee with locomotor disability in the Rotterdam study, 884
 Physiotherapists and occupational therapists, evaluation of extended role in rheumatology practice, (A1) 130

- Physiotherapy and occupational therapy for juvenile chronic arthritis: custom and practice in five centres in the UK, USA and Canada, 695
- PICCOLI A *see* Fiocco U
- PICKERING A *see* Tennant A
- PICKERING M *see* Ames PRJ
- PIETTE J-C *see* Le Thi Huong D
- PILE K *see* Hall FC
- PINA SALVADOR JM, ROYO SERRANO M, ZIAD F, DIEZ MANGLANO J. Importance of cervical narrow disc spaces in patients with rheumatoid arthritis, (A1) 206
- PINCUS T, GRIFFITH J, PEARCE S, ISENBERG D. Prevalence of self-reported depression in patients with rheumatoid arthritis, 879
- PINDER IM *see* Liles JH; Lingard EA
- PINDER S *see* Brennan FR; Jobanputra P
- PINILLOS V *see* Rodriguez-Valverde V
- PIPERNO M *see* Conrozier T
- PIPTONE N, GOULDING NJ, BIAGINI P, PANAYI GS, PITZALIS C. Glucocorticoid induced inhibition of adhesion molecule expression is reversed by IL-2, but not by TNF- α , IL-1 β IL-4 or IL-10, (A1) 7
- PIPTONE N *see also* Biagini P
- PIRILÄ L *see* Roivainen A
- Piroxicam
- 4-week, double-blind, parallel-group study to compare gastrointestinal effects of meloxicam 7.5 mg and 15 mg, piroxicam 20 mg and placebo, (S1) 61
 - double-blind study to compare efficacy and safety of meloxicam 15 mg with piroxicam 20 mg in patients with osteoarthritis of the hip, (S1) 35
 - tolerability of multiple administration of intramuscular meloxicam: comparison with intramuscular piroxicam in patients with rheumatoid arthritis or osteoarthritis, (S1) 51
- PITZALIS C. Role of adhesion mechanisms in the pathogenesis of chronic synovitis, (A1) 125, 1198
- PITZALIS C *see also* Biagini P; Pipitone N; Yanni G
- PIZZUTI P *see* Lioté F
- PLANCHENAU T *see* Chevalier X
- PLANS P *see* Escolà A; Obach J
- PLANT MJ, JONES PW, DAWES PT. Tenosynovitis of the hand: major determinant of functional ability in rheumatoid arthritis, (A1) 202
- PLANT MJ, JONES PW, OLLIER WER, DAWES PT. Early rheumatoid arthritis: 50% of longterm radiological progression can be predicted in the first year, (A1) 206
- PLANT MJ, STEWART WR, WILLIAMS GP, O'CONNOR KBR, GORMAN ST, GRIFFITHS H, WOODCOCK JP, WILLIAMS BD. Gadolinium-enhanced magnetic resonance imaging of the finger joints in rheumatoid arthritis, (A1) 118
- Plasma cells, clonally expanded in rheumatoid synovium, (A1) 172
- Plasminogen activation system, difference in expression in synovial tissue of patients with rheumatoid arthritis and osteoarthritis, 416
- Platelet activating factor receptor antagonists, new, antiinflammatory effect in experimental models of acute and chronic arthritis, (A1) 13
- PLATER-ZYBERK C *see* Taylor PC
- PLATSOUKAS CD *see* Sakkas LI
- PLATT R *see* Troughton PR
- PLAZA MA *see* De Miguel E; Rodriguez R
- PLUMMER CJ *see* Chatterjee S
- POKORNY G *see* Makula E
- POLEKSIĆ L, MUSIKIĆ P, ZDRAVKOVIĆ D, WATT I, BACIĆ G. MRI evaluation of the knee in rheumatoid arthritis, (S3) 35
- Polipodium leucotomos* (Anapsos), modification of the inflammatory activity of psoriatic arthritis, (A1) 161
- Polyarthritis nodosa
- antineutrophil cytoplasmic antibodies, abnormal angiograms and pathological findings, 958
 - in human immunodeficiency virus infection: report of four cases and review of the literature, 796
 - isolated of the prostate, (A1) 51
 - in the North West Region, (A2) 38
 - presenting as ischaemic colitis, (L) 1334
 - in two brothers presenting at the same age, (A2) 38
- Polyarthritides
- chronic and graft-versus-host disease, (L) 297
 - inflammatory
 - natural remission in: issues of definition and prediction, 1096
 - role of the HLA-DRB1 genotype in the susceptibility to and severity, (A1) 186
 - seronegative, synovial fluid interleukin-8 and neutrophil function, 1244 - Polymorphonuclear leucocytes and chondrocytes, study of cooperation in leukotriene biosynthesis, (A1) 62
 - Polymorphonuclear neutrophils, INF γ -treated, survival and functional properties, (A1) 61
 - Polymyalgia rheumatica
 - associated with both HLA-DRB1*0401 and DRB1*0404, 632
 - CD8 lymphocyte subsets in: comparison with elderly-onset and adult rheumatoid arthritis and influence of prednisone therapy, 642
 - controlled trial of intramuscular methylprednisolone vs oral prednisolone, (A1) 59
 - effects of prednisolone and deflazacort on bone metabolism, (A1) 58
 - ESR at presentation is a prognostic indicator for bone loss and duration of treatment, (A1) 58
 - as the rheumatological manifestation of myelodysplastic syndrome in a Chinese patient, (L) 496
 - with a slightly increased erythrocyte sedimentation rate, (A1) 56
 - systemic IL6 may be responsible for increased bone turnover, (A2) 4
 - and temporal arteritis, prospective study of 287 patients, 1161
- Polymyositis, T-lymphocyte immunophenotyping, 839
- Polymyositis-associated overlap syndrome, (E) 305
- PONSFORD FM *see* Perry MJ
- POOL AJ, ALAVI A, AXFORD JS. β 1,4-Galactosyltransferase isoenzyme changes in serum of patients with rheumatoid arthritis, (A1) 174
- POPE DP, CROFT PR, PRITCHARD CM, MACFARLANE GJ, SILMAN AJ. Frequency of restricted range of movement in individuals with self-reported shoulder pain: results from a population-based survey, 1137
- POPE DR, CROFT PR, PRITCHARD CM, SILMAN AJ, MACFARLANE GJ. Frequency of restricted range of movement in individuals with self-reported shoulder pain: results from a population based survey, (A1) 132
- POPERT AJ, JONES PJ, EL-BADAWY SA. Ankylosing spondylitis: treatment of chronic painful focal lesions with low-dose radiotherapy, (A1) 156
- PORTER DR, CAPELL HA, MCINNES I, MUNRO R, MADHOK R, HUNTER JA, THOMSON EA. Is RA becoming a milder disease? Or are we starting second-line therapy in patients with milder disease?, 1305
- PORTER DR *see also* Capell HA; Duncan MR; McEntegart A; Morrison E
- PORTIOLI I *see* Boiardi L
- PORZIO F. Cyclosporin A in the long-term treatment of psoriatic arthritis, (L) 1331
- POTTER A *see* Davis DM
- POULTON K *see* Hajer AH
- POVEDA MJ, FERNÁNDEZ-DAPICA P, MANJÓN P, RODRIGUEZ M, GONZÁLEZ MR, GÓMEZ-REINO JJ. Calcific periarthritis involving multiple sites. A familial condition, (A1) 133
- POVEDA MJ *see also* De La Mata J; Fernández-Dapica P
- POVEDANO J *see* Albornoz J; De La Iglesia JL
- POWELL RJ *see* Lim KL
- POWER C *see* Bolger T
- Practice nurse, experiment in further education: rheumatology nurse practitioner role in primary care, 581
- PRAKKEN ABJ *see* van der Net J
- PRAPOTNIK S *see* Jevtic V
- Prediction of diagnosis in acute and subacute oligoarthritis of unknown origin, 359
- Predictors of mortality in a cohort of patients with newly diagnosed scleroderma, (A1) 122
- Prednisolone
- and cyclophosphamide, intermittent pulse for the treatment of interstitial lung disease in systemic sclerosis, (A2) 27
 - and deflazacort effects on bone metabolism in polymyalgia rheumatica, (A1) 58
 - and methotrexate, accidental overdose, (L) 704
 - oral vs intramuscular methylprednisolone, controlled trial in polymyalgia rheumatica, (A1) 59
 - serum markers of articular cartilage and synovial tissue metabolism in patients treated for early rheumatoid arthritis, (A1) 238
- Prednisone comparison with CD8 lymphocyte subsets in active polymyalgia rheumatica, 642
- PREEDY VR, BALDWIN D, WASSIF WS, SCOTT DL. D-penicillamine induced myopathy: reductions in total muscle RNA and copper, (A1) 72
- PREEDY VR, SCOTT DL. Reductions in the synthesis rates of skeletal muscle proteins in D-penicillamine treated rats, (A1) 71
- Pregnancy
- autoimmunisation to an endogenous retroviral envelope protein during—an autoantigen in congenital heart block, (A1) 112
 - flatfoot in, (L) 396
 - increased rate of lupus flare during: a prospective study of 78 pregnancies, 133
 - serum sex hormones in: comparison between lupus, rheumatoid and control patients, (A1) 35
- PRESETNIK M *see* Jevtic V
- PREVOO MLL, VAN GESTEL AM, VAN'T HOF MA, VAN RIJSWIJK MH, VAN DE PUTTE LBA, VAN RIEL PLCM. Remission in a prospective study of patients with rheumatoid arthritis. American Rheumatism Association preliminary remission criteria in relation to disease activity score, 1101
- PRICE EJ, COHEN SBA, BROOKES SM, WEBB LMC, VENABLES PJW. T cell clones from a subset of patients with primary Sjögren's syndrome produce high levels of IL-10, (A1) 16
- PRICE EJ *see also* Rigby S
- PRICE PA *see* Johansen JS

- PRICE TR *see* Bamji AN
- PRIOLO F *see* Ferracioli GF
- PRITCHARD CM *see* Macfarlane GJ; Pope DP; Pope DR
- Pro-collagenase and pro-stromelysin levels, effect of 6 months treatment on in rheumatoid arthritis patients, (A1) 120
- Pro-NSAIDs, non-acidic and COX-2 inhibitors do not uncouple mitochondrial oxidative phosphorylation, (A1) 222
- Pro-stromelysin and pro-collagenase levels, effect of 6 months treatment on in rheumatoid arthritis patients, (A1) 120
- PROCKOP D *see* Hardwick L
- Prognostic markers, development and validation of response criteria in rheumatoid arthritis: steps towards an international consensus, (S2) 4
- Prolactin, serum and salivary testosterone in young and middle-aged men with ankylosing spondylitis, (A1) 152
- Propricoceptive acuity assessment in the lower limb, evaluation of reliability, reproducibility and validity of two methods, (A1) 140
- PROST A *see* Maugars Y
- Protein kinase activation, human umbilical vein and dermal microvascular display heterogeneity in expression of surface antigens following, (A1) 11
- Proteinase 3 (Wegener's autoantigen) expression in a baculovirus system, (A1) 51
- Proteoglycan
and fibronectin proliferation and synthesis, SAME restores the changes induced by TNF α on cultured synovial cells, (A1) 63
markers and chondroitin sulphate isomers in synovial fluids from healthy volunteers and patients with arthritis, (A1) 239
synthesis accelerated in human chondrocytes following cyclical pressurisation is mediated by β 1 integrin and stretch activated ion channels, (A1) 110
synthesis and content, insulin-like growth factor-1 receptor localization in human normal and osteoarthritic cartilage, 1044
- Proteoglycan-degrading activity associated with the 40 kDa collagen-binding fragment of fibronectin, 506
- Proteus, Klebsiella antibodies and the peptides HLA-B27 and HLA-DR4 (DW15) in Japanese patients with ankylosing spondylitis and rheumatoid arthritis, (A2) 12
- Proto-oncogene bcl-2 overexpression in rheumatoid synovium, (L) 803
- PROUSE P, COX N, DAVIES P, DEVLIN J, DIXEY J, EMERY P, GOUGH A, JAMES D, WILLIAMS P, WINFIELD J, YOUNG A. Prediction of outcome in 537 patients with newly diagnosed rheumatoid arthritis examining the effect of age at onset and gender, (A1) 233
- PROUSE P *see also* Talamo J
- PRUSZCZYNSKI MSM *see* Ronday HK
- Pseudo-Sjögren's syndrome, phenytoin-induced, (L) 1033
- Pseudarthrosis, spinal complicating ankylosing spondylitis: a report of two patients, 1319
- Psoriasis and ankylosing spondylitis, atypical antineutrophil cytoplasmic antibodies (ANCA), (A1) 161
- Psoriatic arthritis
arthroscopic synovectomy in knee joint synovitis: long-term outcome, 463
CD1 expression in skin and synovial membrane, (A1) 179
characterisation of CD8+ T cells in synovial effusions, (A1) 160
clinical, radiographic and HLA associations as markers for patterns, (A1) 159
cyclosporin A in the long-term treatment, (L) 1332
hand radiology: a five year follow-up study, (A1) 158
incidence in Finland, 1289
modification of the inflammatory activity in patients treated with extract of *Polypodium leucotomos* (Anapso), (A1) 161
peripheral joint disease: a five year clinical follow-up study, (A1) 158
prevalence of the rheumatoid arthritis shared epitope, (A1) 159
restricted TCR expression on synovial fluid CD4 and CD8 T cells, (A1) 160
serum tenascin levels, (A1) 176
sulphasalazine in: a randomized, multicentre, placebo-controlled study, 664
termination of disease-modifying drugs: study of 109 courses of treatment, 564
- Psychiatric hospital admissions, previously undiagnosed systemic lupus erythematosus, (A1) 36
- Psychological factors, affecting knee pain and disability in the community, (A1) 134
- Psychological impact of knee pain, (A1) 135
- Psychological and psychopathological evaluation of systemic sclerosis and rheumatoid arthritis patients, (A1) 47
- Psychosocial factors, occurrence of low back pain in relation to in the workplace, (A1) 141
- PUECHAL X *see* Hilliquin P
- Puerperium, increased rate of lupus flare during: a prospective study of 78 pregnancies, 133
- PUGH MT, GROSSE R, SOUTHWOOD TR. Multiple joint corticosteroid injections in polyarticular juvenile chronic arthritis: a single-blind one year follow-up study, (A1) 121
- PUGH MT, SHERRAN T, GOUGH A, EMERY P, BACON PA. Intermittent pulse prednisolone and cyclophosphamide for the treatment of interstitial lung disease in systemic sclerosis, (A2) 27
- PUGH MT, SITUNAYAKE RD, EMERY P. Double blind study assessing safety and efficacy of novel IL-1 antagonist, IX207-887 for treatment of rheumatoid arthritis, (A1) 214
- PUIG N *see* Román JA
- PUJOL M *see* Ros S
- PULLAR T *see* Ho M; Muir A
- Pulmonary abnormalities in patients with early rheumatoid arthritis, (A2) 16
- Pulmonary complications, occurrence during methotrexate therapy in rheumatoid arthritis, 441
- Pulmonary function
evolution in ankylosing spondylitis, (A1) 151
in rheumatoid arthritis treated with low-dose methotrexate: a longitudinal study, 446
- PUNNONEN J *see* Isomäki P
- PURDIE DW *see* Doherty SM
- Purine enzyme activities in early rheumatoid arthritis, (A1) 192
- PYKE S *see* Ames PRJ; Stoll T
- Quadriceps strength
activation and functional performance in healthy young, middle-aged and elderly subjects, (A1) 140
and depression are important factors in knee pain and associated disability, (A1) 135
- Quality of care and patient satisfaction, 382
- Quality of life
comparison of the SF-20 and SF-36 in the assessment in SLE, (A1) 31
influenced by disease activity but is impaired in all SLE patients, (A1) 31
measures, 275
in rheumatoid arthritis: EuroQol is as responsive as ACR disease activity core measures to self-reported change in RA, (A2) 17
- QUIJADA J *see* Alborno J; De La Iglesia JL
- QURTOM M *see* Malaviya AN
- Radiographic change, symmetry in hospital-referred knee osteoarthritis, (A1) 97
- Radiographic damage development during the first 5-6 yr of rheumatoid arthritis. Prospective follow-up study of a Swedish cohort, 1106
- Radiographic progression, slow in a 15-year follow-up of patients with undifferentiated spondylarthritis, (A1) 157
- Radiographs, skyline are better than lateral for detecting progression of patellofemoral osteoarthritis, (A1) 98
- Radiography
clinical improvement and radiological deterioration in rheumatoid arthritis: evidence that the pathogenesis of synovial inflammation and articular erosion may differ, 1263
plain and magnetic resonance imaging, comparative study in the assessment of sacroiliitis in early spondylarthritis, (A1) 117
quantitative, microfoal techniques: measurement of cancellous bone organization, (S3) 19
- Radiological and clinical progression, differences in rheumatoid arthritis, (A1) 199
- Radiological erosions, simple algorithm to predict development in rheumatoid arthritis, (A1) 119
- Radiology in ankylosing spondylitis: comparison of two scoring systems, (A2) 23
- Radiosynovectomy efficacy evaluation of the knee joint with gadolinium-enhanced magnetic nuclear resonance, (A1) 224
- RAFFERTY D *see* Hamilton J
- RAFI S *see* Mahmud T
- RAFTER GW. Pregnancy, schizophrenia and rheumatoid arthritis, (L) 603
- RAGHUPATHY R *see* Malaviya AN
- RAHMAN MA *see* Ravirajan CT
- RAJAN KT *see* Byrne PAC
- RAJTA R *see* Makula E
- RALSTON SH *see* England AJ; Grabowski PS
- RAMAGE W *see* Yanni G
- RAMESAR R *see* Beighton PH
- RAMOS J *see* Mulero J
- RAMOS M *see* Garcia-Carrasco M
- RAMOS P *see* Ames PRJ; Brito M; Cantabrana A
- RAMOS P, GANGI M, BAREN A, FILIPPA D, ELKON KB. Histological analysis of murine CD95/FAS/APO-1, (A1) 39
- RANDS A *see* Harvey GR
- RANKIN ECC *see* Choy EHS
- RAOOF AL-SHAYE A *see* Malaviya AN
- RAPSON N *see* Isaacs JD

- RASHID T see Tiwana H
- RASKER JJ see Brus HLM; Riemsma RP; van den Hoogen FHJ; van der Wardt AM
- RASMUSSEN N see Luqmani RA
- RASPE A see RASPE H
- RASPE H, MATTHIS CH, RASPE A. Back pain in Great Britain and Germany, (A1) 142
- RASPE H see also Kohlman T
- RASPE HH see Mau W
- RAVAUD P, AULELEY G-R, CHASTANG C, ROUSSELIN B, PAOLOZZI L, AMOR B, DOUGADOS M. Knee joint space width measurement: an experimental study of the influence of radiographic procedure and joint positioning, 761
- RAVENTOS A see Olivé A
- RAVIRAJAN CT, ANILKUMAR TV, SARRAF CE, ALISON MR, ISENBERG DA. Analysis of apoptosis in lymphoid organs and lupus disease in murine systemic lupus erythematosus, (A1) 40
- RAVIRAJAN CT, RAHMAN MA, EHRENSTEIN MR, LATCHMAN DS, ISENBERG DA. Sequence analysis and pathogenic properties of a human IgG anti-DNA antibody, (A1) 111
- Raynaud's phenomenon
- economic nailfold capillaroscopy study: follow-up and evaluation as prognostic value, (A1) 49
 - primary
 - non-invasive assessment of vascular reactivity in forearm skin, 1281 and systemic sclerosis, increased nailfold capillary dimensions, 1127
 - primary suspected, prevalence of underlying connective tissue diseases, (A2) 26
 - primary and systemic sclerosis
 - digital and microvascular blood flow responses, (A1) 42
 - digital neurovascular mechanisms as assessed by iontophoresis and laser Doppler techniques, (A1) 48
 - and scleroderma, pilot study of low molecular weight heparin treatment, (A1) 43
 - thermal stress testing and video nail-fold capillaroscopy in children, (A2) 26
- Raynaud's phenomenon in rheumatoid arthritis, 752
- Reactive arthritis and persistent infection, *Chlamydia pneumoniae*, (L) 1189
- Reactive oxygen species generation, progressive loss in neutrophils undergoing apoptosis: an antiinflammatory effect in synovitis, (A1) 61
- REANEY L see Coombes G
- Red cell complement receptor 1 deficiency in systemic lupus erythematosus in southern Chinese: correlation with disease activity, (A1) 26
- REECE R see Devlin J
- REECE RJ, VEALE DJ, EMERY P. Audit of needle arthroscopy service, (A1) 127
- REES DHE see Dalziel M
- REES J see Hurley MV
- REES JIS see Byrne PAC
- REEVE J see O'Neill TW
- Referral
- do GPs refer to rheumatologists on the basis of symptoms or as a result of tests?, (A2) 30
 - early and outcome in rheumatoid arthritis, (A1) 229
- Referral service, acute rheumatology, clinical need for, 389
- REGAN M see Ledingham J
- REGINATO A see Hardwick L
- REGINSTER JY, DISTEL M, BLUHMKE E. Double-blind, three-week study to compare the efficacy and safety of meloxicam 7.5 mg and meloxicam 15 mg in patients with rheumatoid arthritis, (S1) 17
- Rehabilitation, deficient during the course of early RA, (A1) 228
- REICHLIN M see Garton MJ
- REICHLIN MW see Garton MJ
- REID AJC, SCOTT DGI, WATTS RA, HARRISON BDW, WATKIN SW, MCCANN BG. Churg Strauss syndrome in a district general hospital, (A1) 53
- REID DM, COOPER C, KIRWAN JR. Effects of corticosteroids on bone mass over 2 years when used in management of early rheumatoid arthritis, (A1) 210
- REID DM, ENGLAND AJ. Bone mass measurements, peripheral—is there any clinical value in rheumatoid arthritis?, (E) 109
- REID DM see also England AJ; Martin JC; Murray R; O'Neill TW
- REID R see Murray R
- REINO S see Graña J
- REISS G see Drees-Werling U
- Reiter's syndrome, successful treatment in a patient with AIDS with methotrexate and corticosteroids, (L) 295
- Remission
- natural in inflammatory polyarthritis: issues of definition and prediction, 1096
 - predicting and deciding on in rheumatoid arthritis, (E) 1039
 - Remission in a prospective study of patients with rheumatoid arthritis. American Rheumatism Association preliminary remission criteria in relation to disease activity score, 1101
 - Removal of patients from rheumatology waiting list, (A1) 129
 - Renal disease analysis and its involvement in mortality in patients with systemic lupus erythematosus, (A1) 37
 - Renal tubular acidosis and thyrotoxicosis presenting as hypokalaemic paralysis, 289
 - Renal tubular dysfunction is associated with endothelial damage in systemic sclerosis, (A1) 44
 - RENOUX M see Hilliquin P
 - RESINES C see De La Mata J
 - Response index for monitoring efficacy of DMARDs in rheumatoid arthritis, (A2) 21
 - Retinoic acid, TGF β and IL-1 α effects on homeostasis of aggrecan and biglycan from articular cartilage, (A1) 7
 - Retroviral envelope protein, autoimmunisation to during pregnancy—an autoantigen in congenital heart block, (A1) 112
 - Retrovirus-3, human endogenous, a highly conserved immunoregulatory sequence?, (A1) 17
 - REUNANEN A see Heliövaara M
 - REVERTER JC see Navarro M
 - RGUEZ-LOZANO B see Hdez-Berian JA
 - Rheumatic disease patients, prone to Sjögren's syndrome and/or lymphoma, mount an antibody response to BHRF1, the Epstein-Barr viral homologue of BCL-2, 1075
 - Rheumatic diseases, general public's perceptions, (A1) 126
 - Rheumatic disorders of the eye and the various structures involved, 258
 - Rheumatic fever, continued low level occurrence in an Irish population, (A1) 164
 - Rheumatic manifestations in liver transplant recipients, (A1) 87
 - Rheumatoid arthritis
 - 24hr activity in: comparison with HAQ, (A1) 233
 - adding low-dose cyclosporin A to parenteral gold therapy: a double-blind placebo-controlled study, 1142
 - β_2 -adrenergic receptors on synovial fluid lymphocytes and peripheral blood lymphocyte subsets, (A1) 65
 - agalactosyl IgG [Gal(o)]—an analysis of its clinical utility in the long-term follow-up, 1063
 - age effects on the efficacy and tolerance of methotrexate, 453
 - American College of Rheumatology core set and SF-36—physical and social functioning measures of the SF-36 are strongly associated with the Health Assessment Questionnaire, (A1) 195
 - ANCA in, (L) 1328
 - angiogenesis: a critical process in the pathogenesis—a role for VEGF, (E) 917
 - animal models, (E) 1041
 - anti-ganglioside antibodies in patients complicated by peripheral neuropathy, 725
 - anti-keratin antibodies, disease activity and radiological damage, (A2) 14
 - antibodies to four Gram-negative bacteria which share sequences with the rheumatoid arthritis susceptibility motif, 592
 - antibodies to synovial nerve antigens reflect severity, (A1) 171
 - antineutrophil cytoplasmic antibodies, 38
 - are we serving female patients less well than males? Baseline characteristics of patients enrolled into sulphasalazine studies, (A1) 213
 - arthroscopic synovectomy in knee joint synovitis: long-term outcome, 463
 - associated glycosylation changes may not be idiosyncratic to IgG, (A1) 174
 - association of extended TNF-HLA haplotypes in males and females, (A1) 189
 - association of HLA with toxicity in patients receiving gold sodium thiomalate in a Spanish population, (A1) 211
 - association of the shared epitope with radiological severity, (A1) 187
 - atlantoaxial subluxation, stiff collar for treatment, 771
 - atypical synovial cells are involved in collagen type I mRNA-expression, (A1) 178
 - audit of shared care between hospital and general practice in management, (L) 398
 - autoreactive T cells recognize a new 68-k autoantigen, (A1) 107
 - bone mass measurement and bone metabolism: a review, 309
 - and bronchiectasis, defective antibody production, (A1) 201
 - CAMPATH-1H—an intravenous dose-ranging study, 231
 - cause of death at autopsy in 22 patients, (A2) 22
 - causes of cachexia, (A1) 200
 - CD1 expression in skin and synovial membrane, (A1) 179
 - CD8+ T cells and their T cell receptor repertoires, (A1) 183
 - CD69 on synovial T cells correlates with disease activity, (L) 397
 - central nervous system side effects with low dose methotrexate and mycristin therapies, (A1) 216
 - changes in markers of bone metabolism, (A2) 3

- clinical associations of dual-energy X-ray absorptiometry measurement of hand bone mass, 1256
- clinical features of HRCT diagnosed interstitial lung disease, (A2) 16
- clinical improvement and radiological deterioration: evidence that the pathogenesis of synovial inflammation and articular erosion may differ, 1263
- combination therapy with cyclosporin, (S2) 19
- comparison of disease status in patients attending and not attending a specialist clinic, 1169
- compound heterozygosity for the shared epitope and the risk and severity of rheumatoid arthritis, (A1) 187
- construct validity of EuroQol in 240 patients stratified by functional class, (A2) 18
- correlates of functional disability: a cross-sectional study of 706 patients in four European countries, 746
- corticosteroids in management, (E) 2
- cost, 781
- detection of active by ^{99m}Tc human immunoglobulin (IgG), (A2) 15
- determination of cytokine profiles in RA synovial tissue using reverse transcriptase PCR, (A1) 6
- development of foot deformities, (A1) 198
- development of radiographic damage during the first 5-6 yr. Prospective follow-up study of a Swedish cohort, 1106
- development and validation of response criteria: steps towards an international consensus on prognostic markers, (S2) 4
- diagnostic value of anti-RA33 antibody, antikeratin antibody, antiperinuclear factor and antinuclear antibody: comparison with rheumatoid factor, 620
- differences between the clinical and radiological progression, (A1) 199
- discrepancies in 5 year mortality during second line therapy—cultural or procedural?, (A1) 234
- disease expression and HLA association: comparison of Egypt and the UK, (A1) 185
- dissociation of immune response to the degraded and nondegraded (native) Sa antigens, (A1) 18
- does exposure to immunosuppressive therapy increase the 10 year malignancy and mortality risks? A matched cohort study, (A1) 227, 738
- double blind study assessing safety and efficacy of novel IL-1 antagonist IX207-887, (A1) 214
- double-blind, three-week study to compare the efficacy and safety of meloxicam 7.5 mg and meloxicam 15 mg, (S1) 17
- dysfunctional allele of the mannose binding protein gene, (A1) 189
- early
- 50% of longterm radiological progression can be predicted in the first year, (A1) 206
 - ARA 1987 criteria do not perform well, (A2) 18
 - deficient rehabilitation during the course of, (A1) 228
 - joint destruction recommences after glucocorticoids are withdrawn, (A2) 22
 - pulmonary abnormalities, (A2) 16
 - purine enzyme activities, (A1) 192
 - synovitis is associated with future radiological erosions in individual joints of the hands and feet, (A1) 229
- early referral and outcome, (A1) 229
- effect of 6 months treatment on serum pro-stromelysin and procollagenase levels, (A1) 120
- effect of corticosteroids on bone markers, (A1) 120
- effect of second line anti-rheumatic therapy on gait characteristics, (A1) 212
- effects of bone disease duration appendicular bone mineral measurements, (A1) 86
- effects of corticosteroids on bone mass over 2 years when used in management, (A1) 210
- effects of oral clodronate on disease activity and bone mineral density, (A2) 21
- effects of total knee replacements in patients with rheumatoid versus osteoarthritis, (A2) 32
- elderly onset and adult, comparison with CD8 lymphocyte subsets in active polymyalgia rheumatica, 642
- employment attrition in a community based inception cohort, (A1) 235
- endogenous opioid tone, 436
- engineered human anti-tumour necrosis factor alpha antibody reduces serum interleukin-6 and urine bone markers but has no effect on soluble CD4 and soluble interleukin-2 receptor, (A1) 214
- eruption of vasculitis during treatment with leflunomide, (A1) 226
- evaluation of a Dutch version of the AIMS2, 755
- evidence for a central component to pain, (A1) 125
- expression of bcl-2, 611
- expression of cyclooxygenase-2 in human and an animal model, 711
- expression of the Lewis X trisaccharide on serum IgG, (A1) 175
- expression of the multidrug resistance glycoprotein 170 in peripheral blood lymphocytes, 430
- factors affecting ambulation, (A2) 20
- Fc γ RIIA polymorphism, (A1) 28
- fore-arm bone mineral density in non-steroid treated is reduced in patients on long-term methotrexate compared with other disease-modifying drugs, (A1) 209
- frequency of HLA class II antigens amongst Pakistani patients, (A1) 185
- gadolinium-enhanced magnetic resonance imaging correlates with synovial vascularity, (A1) 204
- gadolinium-enhanced magnetic resonance imaging of the finger joints, (A1) 118
- β 1,4-galactosyltransferase isoenzyme changes in serum, (A1) 174
- glycosylation changes of IgG during fasting, 117
- hand bone mineral density, cross-sectional area of forearm musculature and grip strength, (A1) 205
- hand MRI changes, (A1) 118
- health economics as an aspect of health outcome: basic principles and application, (L) 494
- and heparin, (L) 808
- high-resolution magnetic-resonance imaging of erosive changes in the metacarpophalangeal joint, (A1) 124
- HLA class I antigens in serum and synovial fluid, (A1) 176
- HLA class II antigens and synovial fluid cytology, (A2) 12
- hormonal associations due to HLA DR4, (A1) 119
- hyper-IgM syndrome associated with: report of RA in a patient with primary impaired CD40 pathway, 282
- IL-12 expression and quantitation by competitive PCR in the synovium, (A1) 180
- imaging inflammation: comparisons between ^{111}In -anti-E-selectin MAB 1.2B6 and ^{99m}Tc - and ^{111}In -immunoglobulin, (A1) 203
- importance of cervical narrow disc spaces, (A1) 206
- increased catalase activity in synovial fluid of patients compared to osteoarthritis patients, (A1) 70
- influence of formal education on compliance and disease activity in early RA patients starting sulphasalazine, (A1) 216
- informal care for patients, (A1) 127
- interference by IgM rheumatoid factor on ELISA for antibodies to mycobacterial 65 kDa heat shock protein, (L) 300
- interleukin-15 expression, (A1) 180
- intracellular calcium release in T-lymphocytes, (A1) 11
- intramuscular gold therapy in 440 rheumatoid patients—results over 5 years in relation to level of social deprivation at outset, (A1) 213
- is the control of disease progression within our grasp? review of the GRISAR study, (A2) 8
- is homeopathy effective, (A1) 227
- is it becoming a milder disease? or are we starting second-line therapy in patients with milder disease?, 1305
- is there any clinical value in peripheral bone mass measurements?, (E) 109
- joint assessment, (S2) 14
- keratoconjunctivitis sicca in: prevalence in an outpatient population and role of Opticare eye drop dispenser, (A2) 20
- knee laxity, 560
- linkage analysis studies of candidate genes on affected sib pair RA families, (A1) 190
- linkage and association studies of the natural resistance associated macrophage protein locus, (A1) 188
- long-term study to evaluate safety and efficacy of meloxicam therapy, (S1) 29
- low prevalence of hay fever, (A1) 197
- lower gastrointestinal and renal side effects of the COX-2 inhibitor meloxicam, compared to naproxen in rheumatoid arthritis patients, (A1) 223
- mannose-binding protein polymorphism, (A1) 188
- mapping and characterisation of the genes involved, (A2) 10
- matched cross-sectional study of HLA DR types, rheumatoid factor and X-ray damage in North Indians and Pakistanis compared with Northern European patients, (A2) 11
- measurement of morning stiffness in clinical trials, (A1) 197
- measuring disease activity and outcome using the Nottingham Health Profile, (A1) 231
- measuring outcome—which measures are suitable for routine clinical use, (L) 301
- methotrexate in: toxicity issues, (E) 403
- model, use of lymphoid cells capable of transferring disease as carriers of immunomodulators, (A1) 182
- mortality in extended pedigree multicase RA families, (A2) 11
- MRI evaluation of the knee, (S3) 35
- neurological signs in patients receiving gold, (L) 803
- neuropeptides stimulate *in vitro* cytokine release in human leucocytes, (A1) 8
- newly diagnosed, anterior pituitary function, 350
- objective measurement of patient activity before and after steroid injection of the knee, (A1) 208

- occurrence of pulmonary complications during methotrexate therapy, 441
- and osteoarthritis
clinical and laboratory assessments, (S3) 6
difference in expression of the plasminogen activation system in synovial tissue, 416
- outcome measures in: relationship with disease activity and anatomic damage, (A1) 231
- parathyroid hormone-related peptide in synovial fluid and disease activity, 1056
- parvovirus B19, (L) 494
- patient-centred approach to evaluation and treatment: the development of a clinical tool to measure patient-perceived handicap, 921
- perception by patients of the risk benefit ratio of drug treatment, (A1) 212
- peripheral beta-endorphin in: a correlation with the disease activity, (A1) 193
- plasma fatty acid pattern, (A1) 191
- predicting and deciding on remission, (E) 1039
- prediction of outcome in 537 patients with newly diagnosed examining the effect of age at onset and gender, (A1) 233
- prediction of permanent work disability in a follow-up study of early rheumatoid arthritis: results of a tree structured analysis using RECPAM, 652
- pregnancy and schizophrenia, (L) 603
- prevalence of secondary Sjögren's syndrome in multicase RA families, (A2) 10
- prevalence of self-reported depression, 879
- prognostic value of contrast enhanced Gd-DTPA MRI for development of bone erosive changes, (S3) 27
- psychological and psychopathological evaluation of patients, (A1) 47
- pulmonary function during treatment with low-dose methotrexate: a longitudinal study, 446
- quality of life: EuroQol is as responsive as ACR disease activity core measures to self-reported change in RA, (A2) 17
- quantified neurological examination with emphasis on motor and sensory functions, 1116
- randomised controlled study of information leaflets, (A2) 32
- randomized clinical trial of in-patient multidisciplinary treatment versus routine out-patient care, 475
- randomized double-blind controlled trial of sulphasalazine combined with pulses of methylprednisolone or placebo in treatment, 150
- randomized trial of testosterone therapy in males, 568
- Raynaud's phenomenon, 752
- reduced influence of catecholamines on proliferation of lymphocytes, (A1) 65
- relationships between clinical measures of disease activity and biochemical markers, (A1) 230
- relationships of HLA-DRB1 genotypes to susceptibility and severity, (A1) 186
- relevance of daytime and nocturnal pain measurements, (A1) 193
- reliability of simple parameters in the assessment of clinical activity, (A1) 232
- remission in a prospective study of patients. American Rheumatism Association preliminary remission criteria in relation to disease activity score, 1101
- response index for monitoring efficacy of DMARDs, (A2) 21
- rheumatoid factor epitope profiles, (A1) 173
- rheumatoid factor isotypes, an association with radiological progression, (A1) 207
- rheumatoid factor isotypes and radiological progression, (A2) 14
- risk and serum cholesterol in a cohort of 52 800 men and women, 255
- role of human immunoglobulin-Tc99 scan in the assessment of joint inflammation, (A1) 203
- role of TNF α as an inhibitor of erythropoiesis: insights from TNF α blockade *in vivo*, (A1) 217
- secondary amyloidosis in patients: diagnostic and prognostic value of gastroduodenal biopsy, 44
- sensitivity of different analytical methods to detect disease susceptibility genes in RA sibling pair families, (A1) 190
- serum IgA immunoglobulins and sacroiliitis, (A1) 201
- serum markers of articular cartilage and synovial tissue metabolism in patients treated with prednisolone, (A1) 238
- serum sex hormones in pregnancy: comparison between lupus, rheumatoid and control patients, (A1) 35
- serum tenascin levels, (A1) 176
- severe, new perspectives on controlling progression—summary of the potential role of cyclosporin, (S2) 1
- shared epitope prevalence in patients with psoriatic arthritis, (A1) 159
- sialylation of serum immunoglobulin G in relation to galactosylation and rheumatoid factor expression, (A1) 175
- simple algorithm to predict the development of radiological erosions, (A1) 119
- six-month double-blind trial to compare the efficacy and safety of meloxicam 7.5 mg and naproxen 750 mg, (S1) 22
- social deprivation profile in patients commencing sulphasalazine therapy 1990–1993, (A1) 226
- socio-economic factors: effect on functional outcome at three years in 516 patients, (A1) 234
- sodium aurothiomalate and sulphasalazine combined in patients who showed a partial response to sulphasalazine alone, (A1) 224
- somatic mutation and gene fingerprinting of immunoglobulin lambda light chains in the synovium, (A1) 172
- study of foot problems of a consecutive group of patients, (A1) 198
- study to assess the tolerability and safety of meloxicam in patients who have mild renal impairment, (A1) 223
- sulphasalazine-induced systemic lupus erythematosus, (L) 104
- susceptibility of peripheral blood mononuclear cells to peroxide-induced DNA damage, (A1) 184
- synovial fluid interleukin-8 and neutrophil function, 1244
- tenidap versus diclofenac, (L) 602
- tenosynovitis of the hand: major determinant of functional ability, (A1) 202
- Th1 and Th2 cytokines, (A1) 179
- three novel polymorphisms in the regulatory region of the corticotropin releasing hormone gene, (A1) 107
- tolerability of multiple administration of intramuscular meloxicam: comparison with intramuscular piroxicam, (S1) 51
- treatment—does it affect society's cost for the disease?, 791
- treatment with antibody to TNF α decreases expression and shedding of E-selectin, (A1) 218
- tyrosine phosphorylation in neutrophils from synovial fluid, 846
- unusual complications of cervical spine surgery for cervical myelopathy, 682
- valgus heel deformity and plantar pressure distribution of the forefoot, (A1) 199
- validity testing of the MOS SF36 generic health profile, (A2) 17
- value of high mineral content water in the short-term management, (A1) 228
- video image analysis of hands: development of an anatomic index as a potential outcome measure, 1274
- walking time may be a very useful measure of clinical laboratory and radiological status, (A1) 195
- weight loss and body composition, (A1) 200
- why not use OSRA? Comparison of Overall Status in Rheumatoid Arthritis with American College of Rheumatology core set and other validated indices of disease activity, (A1) 196
- Rheumatoid disease and osteoarthritis, comparison of the clinical features of hospital out-patients in Pakistan and England, 994
- Rheumatoid factor
epitope profiles in rheumatoid arthritis, (A1) 173
HLA DR type and X-ray damage, matched cross-sectional study in North Indians and Pakistanis compared with Northern European RA patients, (A2) 11
isotype an association with radiological progression in patients with RA, (A1) 207
isotypes and radiological progression in rheumatoid arthritis, (A2) 14
using the Pepsan technique to map binding to IgG, (A1) 173
- Rheumatoid T cell functional deficiency can be reversed with trispecific antibody stimulation, (A1) 13
- Rheumatological information provision, (E) 503
- Rheumatology
academic, consequences of the Calman Report, (E) 608
clinics, what do general practitioners expect, (E) 920
and dermatology: sister specialties in Italy and Malta, 1312
and immunology in Bergen—from basic science to clinical application, 1159
initiative to reduce orthopaedic waiting lists, (A1) 129
in Israel, 778
manpower and practice, changing patterns in the UK in the 1990s, 483
and medical education in Great Britain, 269
nurse practitioner role in primary care: an experiment in the further education of the practice nurse, 581
outpatient workload increases inexorably, (A2) 29
services, survey of general practice requirements in Leicestershire, (A2) 29
trainees, survey of attitudes and current practices regarding among UK rheumatologists, (A1) 131
training, (L) 1190
undergraduate education, (E) 110, (E) 203
- RHODES CJ *see* Naughton DP
- RIBERA A *see* Lience E
- Ribonucleoproteins, prevalence and clinical significance of antibodies to in systemic lupus erythematosus in Malaysia, 129
- Ribosomal P and Lba5 autoantibodies in NP-SLE, (A1) 111
- RICHARD M *see* Hellio le Graverand MP

- RICHARDS AJ. Audit of shared care between hospital and general practice in the management of rheumatoid arthritis, (L) 398
- RICHARDS PJ, AMOS N, WILLIAMS BD. Comparison of two bisphosphonates on TNF α production by PBMCs, (A1) 4
- RICHARDSON K see Gordon C
- RICHMOND R, MCCRORIE ER, SALTER DM, DUNNE E, NUKI G. Serum tenascin levels in patients with rheumatoid and psoriatic arthritis, (A1) 176
- RIDGEWAY J see Haworth S
- RIEDL M see Templ E
- RIELEY F see Chaloner C; Teh LS
- RIEMEKASTEN G, MARELL J, TREBELIAHR G, SCHNEIDER- MERGENER J, HENKLEIN P, BURMESTER GR, THIELE B, HIEPE F. Synthesis of a synthetic fragment of SmD-polypeptide reveals a probable conformational epitope bound by lupus sera with a high sensitivity and specificity, (A1) 20
- RIEMSA RP, KLEIN G, TAAL E, RASKER JJ, WIEGMAN O. Informal care for patients with rheumatoid arthritis, (A1) 127
- RIEMSA RP, TAAL E, RASKER JJ, HOUTMAN PM, VAN PAASSEN HC, WIEGMAN O. Evaluation of a Dutch version of the AIMS2 for patients with rheumatoid arthritis, 755
- RIGBY S, COOKE S, WEERASINGHE D, PRICE EJ, VENABLES PJW. tax gene of HTLV-1 is not present in the salivary glands of patients with idiopathic Sjögren's syndrome, (A1) 16
- RIGON C see Fiocco U
- RING F see Minaur NJ
- RISCHMUELLER M, DAVIES RP, SMITH MD. Three year follow-up of giant cell arteritis presenting with a chronic cough and upper limb ischaemic symptoms, 800
- Risk factors
of hyperuricaemia in Catalonia-II survey, (A1) 146
for incident low back pain in nurses, (A1) 123
for worse disease activity and function in ankylosing spondylitis, (A2) 24
- RIVERA J, GARCIA-MONFORTE A. Presence of rheumatic diseases in chronic infection by hepatitis C virus, (A1) 168
- RNA polymerases, antibodies to in systemic sclerosis: clinical and serological associations, (A1) 46
- ROBERTS R see Smith G
- ROBERTS TPL see Demars F
- ROBERTSON DAF see McFarlane XA
- ROCAS A see Brito M; Cantabrana A
- RODRIGUEZ R, PLAZA MA, DE MIGUEL E, PASCUAL C, ATERO F, HERNANZ A. Decreased plasma amino acid concentrations in fibromyalgia patients, (A1) 137
- RODRIGUEZ-MAHOU M see Abad MA; Montegudo I
- RODRIGUEZ-MORENO J see Gómez-Vaquero C
- RODRIGUEZ-MORENO J see Narváez J
- RODRIGUEZ-VALVERDE V see Blanco R; González-Gay MA; Sánchez-Andrada S
- RODRIGUEZ-VALVERDE V, BLANCO R, DEL NAL N, ERAUQUIN C, PINILLOS V, MARTINEZ-TABOADA VM. Study of 112 patients with septic arthritis. Clinical subgroups and predisposing factors, (A1) 164
- RODRIGUEZ-VALVERDE V see also Belzunegui J
- ROES L see Ames PRJ
- ROGERS J see Sharma S
- ROHAN D see Montague S
- ROIG-ESCOFET D see Fiter J; Gomez-Vaquero C; Gómez-Vaquero C; Moragues-Pastor C; Narváez J; Ros S
- ROITI I see Bodman-Smith K
- ROIVAINEN A, SÖDERSTRÖM K-O, PIRILÄ L, ARO H, KORTEKANGAS P, MERILAHTI-PALO R, YLI-JAMA T, TOIVANEN A, TOIVANEN P. Oncoprotein expression in human synovial tissue: an immunohistochemical study of different types of arthritis, 933
- ROIVAINEN A see also Isomäki P
- ROJKOVICH B, GIBSON T. Relevance of daytime and nocturnal pain measurements in rheumatoid arthritis, (A1) 193
- ROLDAN R see Collantes E
- ROLDAN R see González J
- Rolipram, amelioration of established collagen-induced arthritis following treatment, (A1) 75
- ROMAGNOLI M see Patoia L
- ROMÁN JA, PUIG N, PEREZ S, ALCAÑIZ C, PASCUAL E. Association of HLA with toxicity in patients with rheumatoid arthritis receiving gold sodium thiomalate in a Spanish population, (A1) 211
- ROMAS E see Hill C
- ROMEY J see Olivé A
- RONDAY HK, SMITS HH, VAN MUIJEN GNP, PRUSZCZYNSKI MSM, DOLHAIN RJEM, VAN LANGELAAN EJ, BREEDVELD FC, VERHEIJEN JH. Difference in expression of the plasminogen activation system in synovial tissue of patients with rheumatoid arthritis and osteoarthritis, 416
- ROOK GAW see Leirisalo-Repo M
- ROS S, JUANOLA X, PUJOL M, CASTELL M, ROIG-ESCOFET D. Effectiveness and response predictive factors of Y⁹⁰ synoviorthesis, (A1) 208
- ROS S see also Fiter J; Gomez-Vaquero C; Gómez-Vaquero C
- ROSS EL, D'CRUZ D, GAFFNEY K, MORROW JWJ. Restricted TCR expression on synovial fluid CD4 and CD8 T cells in psoriatic arthritis, (A1) 160
- ROSS S, WILLIAMS RO, MASON LJ, MARINOVA L, MAURI C, FELDMANN M, MAINI RN. Amelioration of established collagen-induced arthritis following treatment with rolipram, (A1) 75
- ROSSER S see Anderson ME
- ROSSETTI A see Marchesoni A
- ROSSI A see Ferri C
- ROTES-QUEROL J. Clinical manifestations of diffuse idiopathic skeletal hyperostosis (DISH), (E) 1193
- ROTH W see Türk D
- ROUSSELIN B see Ravaud P
- ROUSSOU E, CHOWDHRY KM, KENNEDY LG, CALIN A. Ankylosing spondylitis: ethnic differences, (A1) 149
- ROUX E see Weber M
- ROWAN AD see Clark IM
- ROWLEY DI see McNicholas MJ
- ROY S see Yeap SS
- ROYO SERRANO M see Pina Salvador JM
- ROZMAN B see Jevtic V
- RUBALTELLI L see Fiocco U
- RUDGE SR, CAUGHEY D, MARKS R, KERR A. Prevalence of chronic anterior uveitis in New Zealand children with juvenile chronic arthritis, (A2) 25
- RUDWALEIT M, TIKLY M, KHAMASHTA M, GIBSON K, KLINKE J, HUGHES GRV, WORDSWORTH P. TNF α promoter polymorphisms show inter-ethnic differences in association with SLE, (A1) 28
- RUÍZ-IRASTORZA G, LIMA F, ALVES J, KHAMASHTA MA, SIMPSON J, HUGHES GRV, BUCHANAN NMM. Increased rate of lupus flare during pregnancy and the puerperium: a prospective study of 78 pregnancies, 133
- RUÍZ-IRASTORZA G. Polyarteritis nodosa presenting as ischaemic colitis, (L) 1333
- RUÍZ-LUCEA E see Calabozo M; Perez-Ruiz F
- RUSSELL G see Scott LJ
- RUSSELL K see Öcal L
- RUTA D see Hurst NP
- RYAN P see Mishra MB
- RYAN PJ see Dolan AL; Hamilton I
- RYAN SJ, DAWES PT, MAYER B. Does inflammatory arthritis affect sexuality?, (A2) 19
- RYDGREN L, FREDRIKSON E. Development of foot deformities in early rheumatoid arthritis, (A1) 198
- S-antigen, retinal, antibodies against in patients with juvenile chronic arthritis-associated uveitis, (L) 101
- Sa antigens, dissociation of immune response to degraded and nondegraded (native) in rheumatoid arthritis and systemic lupus erythematosus, (A1) 18
- SAADALLA-NAZHAT RA see Nazhat NB
- SABATE M see Moragues-Pastor C
- Sacroiliitis
comparative study of magnetic resonance imaging and plain radiography in assessment in early spondylarthritis, (A1) 117
serum IgA immunoglobulins and rheumatoid arthritis, (A1) 201
use of multi-bacterial nested PCR to screen synovial fluid and membrane samples, (A1) 113
- SAEED IT, JAWAD ASM. Cause of death at autopsy in 22 patients with rheumatoid arthritis, (A2) 22
- SAEZ F see Calabozo M
- ŞAHİN (ÖZGÜN) Ş, LAWRENCE R, DİRESKENELİ H, HAMURYUDAN V, YAZICI H, AKOĞLU T. Monocyte activity in Behçet's disease, 424
- SAHINOGLU T, BHATT B, HUGHES FJ, BLAKE DR, STEVENS CR. Osteoblastic bone nodules and bone morphogenetic proteins are induced bone growth stimulating pulsed electromagnetic fields, (A1) 115
- SAHINOGLU T see also Ali NN; Jawed S
- ST. CLAIR EW see Isaacs JD
- SAITOH N see Matsuda J
- SAITOH Y see Furukawa M
- SAKAMOTO A see Furukawa M
- SAKKAS LI, JOHANSON NA, BIRKHOLDER, MITRA A, SALGAME P, PLATSOU-CAS CD. IL-12 expression and quantitation by competitive PCR in the synovium of patients with rheumatoid arthritis and osteoarthritis, (A1) 180
- SALAFFI F see Manganelli P
- SALAS E see Medrano C
- SALGAME P see Sakkas LI
- SALIH AM, AHMED S, NIXON NB, DAWES PT, MATTEY DL. Relationship between soluble adhesion molecules and anti-endothelial cell antibodies in vasculitis, (A1) 52
- SALIH AM, NIXON NB, DAWES PT, MATTEY DL. Antibodies to synovial nerve antigens reflect severity of rheumatoid arthritis, (A1) 171

- Endothelial damage is an important mechanism in the pathogenesis of rheumatoid neuropathy, (A1) 170
- SALIH AM, NIXON NB, GAGAN RM, HEATH P, HAWKINS CP, DAWES PT, MATTEY DL. Anti-ganglioside antibodies in patients with rheumatoid arthritis complicated by peripheral neuropathy, 725
- SALIH AM, NIXON NB, HAWKINS CP, HEATH P, DAWES PT, MATTEY DL. Antibodies to nerve antigens in rheumatoid peripheral neuropathy, (A1) 171
- Sera from patients with rheumatoid peripheral neuropathy recognise a common antigen in neuroblastoma cells, (A1) 170
- SALISBURY C *see* Sharif M
- SALMINEN S *see* Malin M
- SALMON M *see* Gordon C; Griffiths B
- Salmonella infection in Saudi patients with systemic lupus erythematosus, (A1) 35
- Salmonella-triggered familial reactive arthritis, (L) 908
- SALMORAL A *see* Gonzalez-Dominguez J
- SALTER D *see* Bavington C; Jobanputra P
- SALTER DM *see* Richmond R
- SALVARANI C *see* Boiardi L
- SAMANTA A *see* Wright J
- SAME restores the changes in proliferation and in the synthesis of fibronectin and proteoglycans induced by TNF α on cultured synovial cells, (A1) 63
- SAMUELS A *see* Koh WH
- SANCHEZ J, BRICENO F, JIMENEZ MD, PÉREZ J, MANZANEQUE L, MARENCO JL. Stroke in patients with systemic lupus erythematosus, (A1) 33
- SANCHEZ P *see* Collantes E
- SANCHEZ-ANDRADA S, FERNÁNDEZ-ECHEVARRÍA MA, CORRALES A, PEÑA JL, RODRIGUEZ-VALVERDE V, CALVO-ALÉN J. Outcome measures in rheumatoid arthritis: relationship with disease activity and anatomic damage, (A1) 231
- SANCHEZ-ANDRADA S *see also* Calvo-Alén J
- SANCHEZ-ANDRADE A *see* Blanco R
- SANCHEZ-NISTAL MA *see* Fernández-Dapica P
- SANCHEZ-PERNAUTE O, PALACIOS I, LÓPEZ-ARMADA MJ, HERNÁNDEZ P, MIGUELEZ R, NAVARRO F, EGIDO J, HERRERO-BEAUMONT G. Tenidap decreases synovial growth (hyperplasia and increased extra-cellular matrix production) and cell infiltration in rabbits with chronic arthritis, (A1) 72
- SÁNCHEZ-PERNAUTE O *see also* Gutierrez S; Palacios I
- SANCHIS J *see* Giralt P
- SANDERS P *see* Blann AD
- SANDERS PA, TEH LS, HILLARBY C, DENTON J, MOUSSA KMK, GRENNAN DM. HLA class II antigens and synovial fluid cytology in rheumatoid arthritis, (A2) 12
- SANSOM DM *see* Balsa A; Patel YI
- SANTIAGO J *see* González-Gay MA
- SANTUCCI L *see* Patoia L
- SANY J *see* Bologna C; Jorgensen C
- SAPHO syndrome in an eleven year old girl, (A2) 36
- SARAUX A, ALLAIN J, GUEDES C, BARON D, YOUNOU P, LE GOFF P. Raynaud's phenomenon in rheumatoid arthritis, 752
- SARAUX A. Antiperinuclear factor in spondylarthropathies, (L) 805
- SARRAF CE *see* Ravirajan CT
- SATTARINIA A *see* Patoia L
- Scalp necrosis in giant cell arteritis, (A2) 36
- SCARPA R, LUBRANO E, COLAO A, MARZULLO P, FERONE D, FERRUCCI MG, DE BRASI D, MEROLA B, LOMBARDI G, ORIENTE P. Articular complaints in acromegalic patients: a preliminary report on clinical and radiological features, (A1) 94
- SCARPA R *see also* Lubrano E
- SCHADE C, SPECKER CH, SCHNEIDER M, LAKOMEK H-J, SCHWOCHAU M. Autoantibodies in ankylosing spondylitis: molecular analysis of the antigen, (A1) 152
- SCHAEVERBEKE T *see* Sibilia J
- SCHAIKE H-G. Why does an inflammation in the joint hurt?, (E) 405
- SCHATTENKIRCHNER M *see* Wojtulewski JA
- SCHEVEN BAA *see* van der Veen MH
- Schizophrenia, pregnancy and rheumatoid arthritis, (L) 603
- SCHLESINGER BC *see* Gordon C
- SCHMIDT RE *see* Hoffmeyer F
- SCHNEIDER EM *see* Specker Ch
- SCHNEIDER M *see* Ostendorf B; Schade C; Specker Ch; Wedekind F
- SCHNEIDER-MERGENER J *see* Riemekasten G
- SCHNITZER TJ *see* Isaacs JD
- SCHOFIELD JB *see* Chapman PT
- SCHOLLUM J *see* Macfarlane GJ
- School Christmas shows...look out the backache is behind you, (L) 1336
- SCHREY G *see* Boonen A
- SCHWARZ IM, HILLS BA. Synovial surfactant: lamellar bodies in type B synoviocytes and proteolipid in synovial fluid and the articular lining, 821
- SCHWOCHAU M *see* Schade C; Specker Ch
- Sciatica, developing a care path from primary through secondary to tertiary care, (A1) 143
- Scleroderma
- co-existent with coeliac disease, (L) 807
- and control, differential modulation of fibroblast proliferation and collagen biosynthesis by endothelin, (A1) 47
- DNA allelic alterations within VNTR loci, 1216
- endothelial cell derived interleukin 1 and basic fibroblast growth factor independently modulate fibroblast properties, (A1) 41
- intermittent iloprost infusion therapy of pulmonary hypertension—a pilot study, (L) 1187
- left ventricular function, 983
- localized, cardiac abnormalities, (L) 703
- massive pericardial effusion, (L) 807
- newly diagnosed, predictors of mortality in a cohort of patients, (A1) 122
- and Raynaud's phenomenon, pilot study of low molecular weight heparin treatment, (A1) 43
- renal crisis: association with anti RNA polymerase, improved outcome with iloprost, (A1) 41
- survival following the onset: results from a retrospective inception cohort study of the UK patient population, 1122
- telomere reduction in: a possible cause for chromosomal instability, 732
- Scleroderma-polymyositis overlap syndrome, antinuclear and antinuclear antibodies, (L) 1326
- Scleromyositis, failure of first-line therapy with intravenous immunoglobulin in a child, (L) 1029
- Sclerosis, subchondral and fractal signature analysis in hip osteoarthritis, (A1) 103
- Scoring system for assessing synovial membrane abnormalities at arthroscopy in knee osteoarthritis, (S3) 15
- SCOTT D *see* Houssien D
- SCOTT DGI. Report of Special Interest Group held at the British Society of Rheumatology meeting in Glasgow, April 1995, 394
- SCOTT DGI *see also* Barrett EM; Carruthers DM; Jolliffe VA; Reid AJC; Symmons DPM
- SCOTT DL, HOUSSEIN DA
- Clinical and laboratory assessments in rheumatoid arthritis and osteoarthritis, (S3) 6
- Joint assessment in rheumatoid arthritis, (S2) 14
- SCOTT DL *see also* Fenn GC; Fisher P; Houssein DA; Houssien DA; Mahmud T; Preedy VR; Smith EC
- SCOTT LJ, NIXON NB, RUSSELL G, DAWES PT, MATTEY DL. Normal urine provides an environment that augments complex formation between IgA and α 1-antitrypsin *in vitro*, (A2) 13
- Scurvy, osteoporosis and megaloblastic anaemia due to alleged food intolerance, (L) 701
- SEGOVIA C *see* Brito M; Cantabrana A
- SEGURA GARCÍA C *see* Hdez-Berriain JA
- SEIBOLD JR, McCLOSKEY DA, CLEMENTS PJ, FURST DE. Total skin score as an independent outcome variable in systemic sclerosis—survey of biases of clinical experts, (A1) 45
- SEIDL G *see* Kainberger F
- SEIFERT B *see* Stoll T
- Self-management community based arthritis courses for older people, (A2) 31
- SEMPERE JM *see* Navarro FJ
- SENÉCAL J-L *see* Uthman I
- SEPE A *see* Jevtic V
- Septic arthritis, study of 112 patients. Clinical subgroups and predisposing factors, (A1) 164
- Septic bursitis: predisposing factors, clinical and therapeutical features, (A1) 165
- Septicaemia, elevated levels of corticotrophin-releasing factor binding protein in the blood of patients and the presence of novel ligands in synovial fluid, 120
- SEREBOUR F *see* Malaviya AN
- Serine proteinase inhibitor antileukoproteinase specifically accumulates in normal but not in arthritic cartilage, (A1) 68
- Serotonin, endogenous, role in adjuvant-induced arthritis in the rat, 112
- SERRANO L *see* Fernández-Dapica P
- SERRE C *see* Houssien D
- Sex hormones in pregnancy: comparison between lupus, rheumatoid and control patients, (A1) 35
- Sexuality, does inflammatory arthritis affect?, (A2) 19
- SF-20 and SF-36 comparison in the assessment of quality of life in SLE, (A1) 31
- SHAFFER JL *see* Teh LS
- SHAMES DM *see* Demars F
- SHANKMAN S *see* Markov G
- Shared care, audit of between hospital and general practice in the management of rheumatoid arthritis, (L) 398
- Shared epitope
- association with radiological severity of rheumatoid arthritis, (A1) 187

- compound heterozygosity and the risk and severity of rheumatoid arthritis, (A1) 187
- SHARIF M, ELSON CJE, DIEPPE PA, KIRWAN JR. C-reactive protein is elevated in osteoarthritis, (A1) 99
- SHARIF M, OSBORNE D, MEADOWS K, WOODHOUSE S, COLVIN E, SHEPSTONE L, DIEPPE PA. Chondroitin sulphate isomers and proteoglycan markers in synovial fluids from healthy volunteers and patients with arthritis, (A1) 239
- SHARIF M, OSBORNE DJ, MEADOWS K, WOODHOUSE SM, COLVIN EM, SHEPSTONE L, DIEPPE PA. Relevance of chondroitin and keratan sulphate markers in normal and arthritic synovial fluid, 951
- SHARIF M, SALISBURY C, KIRWAN JR. Serum markers of articular cartilage and synovial tissue metabolism in patients treated with prednisolone for early rheumatoid arthritis, (A1) 238
- SHARIF M *see also* Cooper AM; Taylor DJ
- SHARMA A, MENON S, ISENBERG DA. Identification of idiotypes on human monoclonal IgG anti-cardiolipin antibodies and in the sera of patients with autoimmune rheumatic diseases, (A1) 22
- SHARMA B *see* Hamilton I
- SHARMA S, ROGERS J, WATT I, BUCKLAND-WRIGHT C. Subchondral sclerosis and fractal signature analysis in hip osteoarthritis, (A1) 103
- SHARMA SD, SMITH EM, HAZLEMAN BL. Thermographic changes in patients with work-related chronic upper limb pain, (A1) 139
- SHARP CA *see* Martin JC
- SHARPE DN *see* Armstrong GP
- SHATFORD J *see* Hardwick LJ
- SHAWIS TN. Pancreatic function in systemic sclerosis, (L) 298
- SHEARING-LYNNE C, BEXENDINE S, THOMPSON PW. Evaluation of disease and drug monitoring in inflammatory arthritis, (A1) 230
- SHEEHAN NJ. Co-existent coeliac disease and scleroderma, (L) 807
- SHELDON P, KADIOGLU A. Is the corticosteroid mediated change in lymphocyte adhesion to mucosal endothelium by mononuclear cells from rheumatoid arthritis patients due to a quantitative change in lymphocyte subsets?, (A1) 14
- SHELDON P, WRIGHT J. Reduced hip and wrist fracture incidence of an Asian population in Leicester, (A2) 3
- SHELDON P *see also* KADIOGLU A
- SHEPSTONE L *see* Sharif M
- SHERMAN T *see* Pugh MT
- SHIMADA S *see* Wada M
- SHIMIZU K *see* Nishikai M
- SHINGU M *see* Fujikawa Y
- SHIPLEY M *see* Symmons DPM
- SHIROKY JB *see* Newkirk MM
- SHIWEEN X, DENTON CP, DASHWOOD MR, ABRAHAM DJ, BLACK CM. Differential modulation of scleroderma and control fibroblast proliferation and collagen biosynthesis by endothelin, (A1) 47
- SHIWEEN X *see also* Denton CP
- SHOENFELD Y *see* Ehrenfeld M
- SHORTALL E, ISENBERG DA, NEWMAN SP
Disease activity vs illness perception as predictors of health status and psychological well-being in patients with systemic lupus erythematosus: longitudinal analysis, (A1) 34
Influence of coping strategy on outcome in systemic lupus erythematosus, (A1) 33
- Shoulder
painful: can consultants agree?, 1172
painful rheumatic, comparison of glenohumeral injections of triamcinolone hexacetonide, (A1) 132
- Shoulder pain, self-reported
frequency of restricted range of movement in individuals: results from a population-based survey, 1137
frequency of restricted range of movement results from a population based survey, (A1) 132
- SHUSTIK C *see* Newkirk MM
- Sialylation of serum immunoglobulin G in rheumatoid arthritis in relation to galactosylation and rheumatoid factor expression, (A1) 175
- SIBANDA EN, BHAYANI H, SOUTHWOOD T, GASTON JSH. Mapping the GroEL epitope recognised by a synovial fluid T cell clone from a boy with juvenile chronic arthritis, (A1) 162
- SIBILIA J, DURANDY A, SCHAEVERBEKE T, FERMAND JP. Hyper-IgM syndrome associated with rheumatoid arthritis: report of RA in a patient with primary impaired CD40 pathway, 282
- SIEPER J *see* Braun J; Wilkinson NZ
- SIGAL E *see* Kang RY
- Signalling, PKC mediated cellular activation, gold sodium thiomalate does not inhibit through CD44 and LFA-1, (A1) 14
- Silencer, cis-acting, DNA sequence and characterization: murine p75 TNF receptor promoter region, (A1) 9
- Silica glass fragments from steroid containing ampoules augment the intra-articular inflammatory response, (A1) 121
- SILMAN A *see* Brennan P; Bryan C; Hajeer A; John S; John SL; Marlow A
- SILMAN AJ *see* Bryan C; Harrison BJ; Macfarlane GJ; Nahit ES; Newman J; O'NEILL TW; O'Neill TW; Pope DP; Pope DR; Symmons DPM
- SILVA H *see* De La Iglesia JL
- SILVERSTONE E *see* Martin JC
- SIM B *see* Guerin J
- SIMÉON CP. Systemic sclerosis in men: clinical and immunological differences, (L) 910
- SIMON CH, DIJKMANS BAC, HAZES JMW, VANDENBROUCKE JP, BREEDVELD FC. Monitoring DMARDs in daily rheumatological practice: are laboratory tests of any use?, (A1) 217
- SIMPSON J *see* Ruiz-Irastorza G
- SINCLAIR D, COOMBS EJ, HULL RG. Do GPs refer to rheumatologists on the basis of symptoms or as a result of tests?, (A2) 30
- SINCLAIR H *see* Bodman-Smith K
- SINNEMA G *see* van der Net J
- SIPEK A *see* Jevtic V
- SIRAG SM *see* El-Shal EW
- SITUNAYAKE RD *see* Griffiths B; Pugh MT
- Sjögren's syndrome
idiopathic, tax gene of HTLV-1 is not present in the salivary glands, (A1) 16
liver involvement and hepatitis C virus infection, (A1) 15
a new model, T-cell repertoire of infiltrating T cells in lacrimal glands, salivary glands and kidneys from alymphoplasia (*aly*) mutant mice, 1223
prevalence in multicase RA families, (A2) 10
primary
complicated by cryoglobulinaemic glomerulonephritis, pericardial and pleural effusions, 72
parotid gland ultrasonography as a diagnostic tool, 972
and patients HCV positive with sicca syndrome, immunological features, (A1) 15
T cell clones from a subset of patients produce high levels of IL-10, (A1) 16
revisited: autoimmune epithelitis, (E) 204
seronegative, manifested as a subset of chronic fatigue syndrome, 471
Skin score, total as an independent outcome variable in systemic sclerosis—survey of biases of clinical experts, (A1) 45
- SLICC/ACR damage index
association with measures of disease activity and health status in patients with systemic lupus erythematosus, (A1) 29
is valid, and renal and pulmonary organ scores are predictors of severe outcome in patients with systemic lupus erythematosus, 248
- SLIEMAN C *see* Lioté F
- SLINGSBY JH, NORSWORTHY PJ, PEARCE GJ, VAISHNAV AK, DAVIES KA, ISSLER H, MORLEY BJ, WALFORD MJ. Molecular basis of complete C1Q deficiency in three patients, (A1) 27
- SLINGSBY JH *see also* Hogarth MB
- SmD-polypeptide, synthesis of a synthetic fragment reveals a probable conformational epitope bound by lupus sera with a high sensitivity and specificity, (A1) 20
- SMEDLEY J, EGGER P, COOPER C, COGGON D. Risk factors for incident low back pain in nurses, (A1) 123
- SMEEDSTAD LM, MOUM T, GUILLEMIN F, KVIEEN TK, FINCH MB, SUURMEIJER TPBM, VAN DEN HEUVEL WJA. Correlates of functional disability in early rheumatoid arthritis: a cross-sectional study of 706 patients in four European countries, 746
- SMITH C *see* Jayanetti S
- SMITH CP *see* Elliott M
- SMITH EC, BERRY H, SCOTT DL. Clinical need for an acute rheumatology referral service, 389
- SMITH EM *see* Sharma SD
- SMITH G, ROBERTS R, HALL C, NUKI G. Reversible ovulatory failure associated with development of luteinized unruptured follicles in women with inflammatory arthritis taking non-steroidal anti-inflammatory drugs, 458
- SMITH H-J. Contrast-enhanced MRI of rheumatic joint disease, (S3) 44
- SMITH JL *see* Wong WM
- SMITH MD *see* Rischmueller M
- SMITH RW, HOGG J, CAWLEY MID, JARVIS L. Comparative study of magnetic resonance imaging and plain radiography in the assessment of sacroiliitis in early spondylarthritis, (A1) 117
- SMITH SR *see* Lingard EA
- SMITS HH *see* Ronday HK
- SMOLEN JS *see* Studnicka-Benke A
- SMYTH AE, BRUCE IN, BELL AL. Prevalence of underlying connective tissue diseases in patients with suspected primary Raynaud's phenomenon, (A2) 26
- SMYTH LJC *see* Snowden N
- SNOWDEN N, BOOTH J, MORAN A, HAENEY MR, SWINSON DR. Defective antibody production in patients with rheumatoid arthritis and bronchiectasis, (A1) 201

- SNOWDEN N, SMYTH LJC, HAJEER A, OLLIER WER. Fc γ RIIA polymorphism in SLE and RA, (A1) 28
- SNOWDEN N *see also* Stanworth SJ
- So A *see* Touray M
- So AKL *see* Loizou S
- Social deprivation
- intramuscular gold therapy in 440 rheumatoid patients—results over 5 years in relation to level at outset, (A1) 213
 - profile in patients commencing sulphasalazine therapy for rheumatoid arthritis 1990–1993, (A1) 226
- Socio-economic factors in rheumatoid arthritis: effect on functional outcome at three years in 516 patients, (A1) 234
- SÖDERSTRÖM K-O *see* Isomäki P; Roivainen A
- Sodium aurothiomalate and sulphasalazine combined in patients with rheumatoid arthritis who showed a partial response to sulphasalazine alone, (A1) 224
- SOLAN SK *see* Lingard EA
- SOLOMON C *see* Highton J
- Somatotroph axis in the fibromyalgia syndrome, (A1) 139
- SOPWITH M *see* Choy EHS
- SORIGUER F *see* Collantes E
- SOUTHWOOD T *see* Hackett J; Sibanda EN
- SOUTHWOOD TR *see* Pugh MT
- Spanish Sond. Study Group *see* Císal A
- SPECHT T *see* Baerwald CGO
- Special Interests Group, report, held at the British Society of Rheumatology meeting in Glasgow, April 1995, 394
- Specialist training into the new millenium: the price of conformity with Europe, (E) 607
- SPECKER Ch, BLÄS St, SCHWOCHAU M, SCHNEIDER EM, SCHNEIDER M. Autoreactive T cells of RA patients recognize a new 68-k autoantigen, (A1) 107
- SPECKER Ch *see also* Ostendorf B; Schade C
- SPECT scan value in identifying back pain likely to benefit from facet joint injection, 1269
- SPECTOR T *see* Cooper C
- SPECTOR TD, CICUTTINI F, BAKER J, LOUGHLIN JA, HART DJ. Genetic influences on osteoarthritis: a study of twins, (A1) 116
- SPECTOR TD *see also* Arden NK; Etherington J; Griffiths GO; Hall GM; Keen RW; Masud T; Nandra D; O'Neill TW
- SPEED CA, CAMPBELL R, FORDHAM JN, UNSWORTH A, HASLOCK I. Hand bone mineral density, cross-sectional area of forearm musculature and grip strength in RA, (A1) 205
- SPEED CA, FORDHAM JN. Central nervous system side effects with low dose methotrexate and mycristin therapies in rheumatoid arthritis, (A1) 216
- SPEED CA, FORDHAM JN, CUNNINGHAM JL. Simultaneous bilateral tibial stress fractures in a 15-year-old milkman—a case report, 905
- SPEED CA, HESLOP P, WALKER DJ. Factors affecting ambulation in rheumatoid arthritis, (A2) 20
- Spinal pseudoarthrosis complicating ankylosing spondylitis: a report of two patients, 1319
- Spine position, how much does it affect bone mineral density values?, (A1) 82
- Spondarthritis, undifferentiated, slow radiographic progression in a 15-year follow-up of patients, (A1) 157
- Spondiloepiphyseal dysplasia associated with familial CPPD deposition disease: study of a pedigree, (A1) 147
- Spondylarthritis
- comparative study of magnetic resonance imaging and plain radiography in the assessment of sacroiliitis, (A1) 117
 - in the Gambia—evidence for a non-HLA B27 protective effect, (A1) 113
- Spondylarthropathies
- antiperinuclear factor, (L) 804, (L) 805
 - are antineutrophil cytoplasmic antibodies associated?, (L) 1032
 - assessment of the efficacy of sacroiliac corticosteroid injections: a double-blind study, 767
 - assessment of performance of ESSG criteria according to diagnostic age, (A1) 150
 - combination therapy of anti-CD4 and anti-IL6 monoclonal antibodies, (L) 1331
 - early oligosymptomatic HLA-B27 typing in the diagnosis, (C) 192
 - influence of non-steroidal antiinflammatory drugs on the evolution of abdominal scintigraphy, (A1) 154
 - their distribution by diagnostic criteria in three countries, (A1) 150
 - undifferentiated seronegative in females, (L) 395
- Spondylitis, severe, agalactosyl IgG is a marker, (A1) 153
- Spondylodiscitis
- cervical, in a patient with ankylosing spondylitis, (L) 1034
 - epidemiology in ankylosing spondylitis—a controlled study, 660
 - infectious: a revision of 98 cases, (A1) 166
 - septic, induced by pyogenic germs in patients without addiction to intravenous drugs, (A1) 165
- SPRONK PE, BOOTSMA H, HORST G, HUITEMA MG, LIMBURG PC, COHEN TERVAERT JW, KALLENBERG CGM. Antineutrophil cytoplasmic antibodies in systemic lupus erythematosus, 625
- STAGG AJ, HUGHES RA, KEAT ACS, ELSLEY WJ, KNIGHT SC. Antigen-presenting cells but not lymphocytes in the joint may indicate the cause of reactive arthritis, 1082
- Stanford Health Assessment Questionnaire, are we making the most of it, 574
- STANWORTH SJ, DONN RP, DAVIES EJ, HASSALL A, DAWES P, SNOWDEN N, OLLIER WER. Mannose-binding protein polymorphism and rheumatoid arthritis, (A1) 188
- STATHOPOULU E, BAILDAM EM, HOLT PJL. Gower's sign in children with juvenile chronic arthritis, (A1) 79
- STEAD H *see* Walker DJ
- STEEL SA *see* Doherty SM
- STEI P, KRUSS B, WIEGLEB J, TRACH V. Local tissue tolerability of meloxicam, a new NSAID: indications for parenteral, dermal and mucosal administration, (S1) 44
- STEINER A *see* Jones SD
- STEINER G *see* Studnicka-Benke A
- STEPHENS C *see* Artlett CM
- Stereological analysis of the synovial membrane in rheumatic disorders, (A1) 177
- STERN M *see* Lioté F
- Steroids
- low dose treatment, effect on serum stromelysin and collagenase levels in RA patients followed for 3 years, (A1) 210
 - use in patients undergoing lower limb arthroplasty, (A1) 209
- STEUER A, AMES PRJ, DENMAN AM. Neurological involvement in Behçet's syndrome in the UK, (A2) 28
- STEVEN MM, LOWRY S. Undergraduate education in rheumatology, (E) 110
- STEVENS CO *see* Artlett CM
- STEVENS CR *see* Ali NN; Jawed S; Sahinoglu T
- STEVENSON A *see* Over KE
- STEVENSON JM *see* Mannion AF
- STEWART A *see* Martin JC
- STEWART I *see* Haworth S
- STEWART P *see* Akil M
- STEWART WR *see* Plant MJ
- STICHTENOTH DO, WAGNER B, TSIKAS D, GUTZKI F-M, DEGNER FL, FRÖLICH JC. Meloxicam does not inhibit platelet aggregation and renal prostaglandin E₂ synthesis in healthy volunteers, (A1) 220
- STICHTENOTH DO *see also* Frölich JC
- Stiff collar for the treatment of rheumatoid atlantoaxial subluxation, 771
- Stiffness, morning, measurement in rheumatoid arthritis clinical trials, (A1) 197
- STOCKER CJ *see* Chapman PT
- STOFFEL MP, CSERNOK E, WEINERTH J, GROSS WL. Immunodiagnostic importance of the BPI-ANCA system, (A1) 50
- Stoke Ankylosing Spondylitis Spine Score (SASSS), radiological outcome in ankylosing spondylitis, 373
- STOLK J, BOERBOOMS A, DE ABREU R, DE KONING D, KERSTENS P, DE GRAAF R, VAN DE PUTTE L. Purine enzyme activities in early rheumatoid arthritis, (A1) 192
- STOLL T, SEIFERT B, ISENBERG DA. SLICC/ACR damage index is valid, and renal and pulmonary organ scores are predictors of severe outcome in patients with systemic lupus erythematosus, 248
- STOLL T, STUCKI G, MALIK J, PYKE S, ISENBERG DA. Association of the SLICC/ACR damage index with measures of disease activity and health status in patients with systemic lupus erythematosus, (A1) 29
- STOLL T, STUCKI G, MALIK J, PYKE S, ISENBERG DA. Further validity of the BILAG disease activity index in patients with systemic lupus erythematosus, (A1) 29
- STOLL T *see also* Gordon C
- STOLTENBERG M *see* Østergaard M
- STONE M *see* Patel S
- STRANGE R *see* Hajeer A
- STRATTON R, BURNS A, BUNN C, BLACK C. Scleroderma renal crisis: association with anti RNA polymerase, improved outcome with iloprost, (A1) 41
- STRATTON R *see also* Denton CP; Dolan AL
- Stroke in patients with systemic lupus erythematosus, (A1) 33
- Stromelysin and collagenase, effect of low dose steroid treatment on levels in RA patients followed for 3 years, (A1) 210
- STUBBINGS A *see* Hurst NP
- STUCKI G. Predicting and deciding on remission in rheumatoid arthritis, (E) 1039
- STUCKI G *see also* Stoll T
- STUDNICKA-BENKE A, STEINER G, PETERA P, SMOLEN JS. Tumour necrosis factor alpha and its soluble receptors parallel clinical disease and autoimmune activity in systemic lupus erythematosus, 1067
- STURROCK RD *see* Crilly A; Houston T; Lim KL; McInnes IB; Morris AJ
- Subarachnoid haemorrhage secondary to ruptured cerebral aneurysm in a patient with osteogenesis imperfecta, (L) 1332

- SUBIACO S *see* Manganelli P
- SUGARS K *see* Mason JC
- Sulphasalazine
- and azathioprine combination therapy, (L) 493
 - combined with pulses of methylprednisolone or placebo, randomized double-blind controlled trial in the treatment of rheumatoid arthritis, 150
 - influence of formal education on compliance and disease activity in early RA patients starting treatment, (A1) 216
 - response to and antibody levels to *Clostridium perfringens*, (L) 1327
 - and sodium aurothiomalate in patients with rheumatoid arthritis who showed a partial response to sulphasalazine alone, (A1) 224
 - therapy, social deprivation profile in patients commencing therapy for rheumatoid arthritis 1990-1993, (A1) 226
- Sulphasalazine-induced systemic lupus erythematosus in a patient with rheumatoid arthritis, (L) 104
- SULTANA S *see* Gibson T
- SUMAR N *see* Bodman-Smith K; Kjeldsen-Kragh J
- SUMIDA T *see* Furukawa M
- SUMMERS CA, CLARK IM, HAZLEMAN BL, CAWSTON TE. Oncostatin M is raised in rheumatoid synovial fluids and interacts with IL-1 to promote cartilage collagen destruction *in vitro*, (A1) 109
- Surfactant, synovial: lamellar bodies in type B synoviocytes and proteolipid in synovial fluid and the articular lining, 821
- Survival following the onset of scleroderma: results from a retrospective inception cohort study of the UK patient population, 1122
- SUURMEIJER TPBM *see* Smedstad LM
- SUZUKI H, HICKLING P, LYONS CBA. Case of primary Sjögren's syndrome, complicated by cryoglobulinaemic glomerulonephritis, pericardial and pleural effusions, 72
- SWAAK AJG. Tenidap versus diclofenac in rheumatoid arthritis, (L) 602
- SWAAK AJG *see also* van den Hoogen FHJ
- SWANNELL AJ. Specialist training into the new millennium: the price of conformity with Europe, (E) 607
- SWINSON DR *see* Snowden N
- SYED A *see* Gibson T
- SYMMONS D *see* Brennan P; Jones M
- SYMMONS DPM, BANKHEAD CR, GRIFFITHS I, SHIPLEY M. Changing patterns of rheumatology manpower and practice in the UK in the 1990s, 483
- SYMMONS DPM, BRENNAN P, HARRISON BJ, BARRETT EM, CHAKRAVARTY K, SCOTT DGI, SILMAN AJ. Simple algorithm to predict the development of radiological erosions in patients with early rheumatoid arthritis, (A1) 119
- SYMMONS DPM *see also* Barrett EM; Carruthers DM; Harrison BJ; Jones MA
- SYMONS MCR *see* Naughton DP
- Symptom impact patterns in acute and chronic knee pain, (A1) 133
- Synergistic interaction between morphine and indomethacin attenuates pain sensation and tissue destruction in adjuvant arthritis, (A1) 75
- Synovectomy, arthroscopic in rheumatoid and psoriatic knee joint synovitis: long-term outcome, 463
- Synovial angiotensin converting enzyme in human arthritis, (A1) 66
- Synovial chondromatosis: prevalence in familial chondrocalcinosis, (A1) 147
- Synovial fibroblasts, facilitated expression of factors involved in B lymphocyte maturation and survival, (A1) 110
- Synovial fluid
- from knees with secondary osteoarthritis has much higher nitrite concentrations than serum, suggesting local production of nitric oxide, (A1) 105
 - interleukin-8 and neutrophil function in rheumatoid arthritis and seronegative polyarthritis, 1244
 - normal and arthritic, chondroitin and keratan sulphate markers, 951
 - routine culture: is it necessary? Lessons from an audit, (A2) 34
 - and serum antibodies against Chlamydia in different forms of arthritis: intra-articular IgA production in *Chlamydia* sexually acquired reactive arthritis, 548
- Synovial infiltrate nature in the synovial membrane in reactive arthritis, (A1) 163
- Synovial inflammation, quantitative assessment by dynamic gadolinium-enhanced magnetic resonance imaging. A study of the effect of intra-articular methylprednisolone on the rate of early synovial enhancement, 50
- Synovial membrane
- abnormalities, proposed scoring system for assessing at arthroscopy in knee osteoarthritis, (S3) 15
 - quantitative assessment in the rheumatoid wrist: an easily obtained MRI score reflects the synovial volume, 965
 - rheumatoid, increased expression of interferon (IFN)-gamma together with IFN-gamma receptor compared with synovium of patients with osteoarthritis, 24
 - stereological analysis in rheumatic disorders, (A1) 177
- Synovial nerve antigens, antibodies to reflect severity of rheumatoid arthritis, (A1) 171
- Synovial surfactant: lamellar bodies in type B synoviocytes and proteolipid in synovial fluid and the articular lining, 821
- Synovial T cells, CD69 in rheumatoid arthritis correlates with disease activity, (L) 397
- Synovial vascular permeability, mapping abnormal in temporomandibular joint arthritis in the rabbit using MRI, (S3) 24
- Synoviorthesis, Y^{90} , effectiveness and response predictive factors, (A1) 208
- Synovitis
- chronic, role of adhesion mechanisms in the pathogenesis, (A1) 125, 1198
 - clinical, arthroscopic and histological features in SLE, (A1) 36
 - crystal, serial measurement of joint stiffness following intra-articular steroid: an early increase may indicate sub-clinical, (A1) 211
 - inflammatory, NADH oxidase activity measured by chemiluminescence, (A1) 69
 - is associated with future radiological erosions in individual joints of the hands and feet of patients with early RA, (A1) 229
 - progressive loss of reactive oxygen species generation in neutrophils undergoing apoptosis: an anti-inflammatory effect, (A1) 61
- Synovium
- and cartilage
 - MRI in the assessment, (S3) 3
 - pathophysiology, (S3) 11
 - inflammatory, intravenous MRI contrast enhancement: a dose study, (S3) 31
 - rheumatoid
 - bone resorption by tartrate-resistant acid phosphatase-positive multinuclear cells, 213
 - clonally expanded plasma cells, (A1) 172
 - differential expression of the costimulatory molecules B7.1 (CD80) and B7.2 (CD86), 33
 - immunolocalisation of inducible nitric oxide synthase, (A1) 177
- Systemic lupus erythematosus
- active, is associated with the recruitment of naive/resting T cells, 226
 - analysis of renal disease and its involvement in mortality, (A1) 37
 - antibodies to C1q(CLR) are markers for severe disease, (A1) 21
 - anticardiolipin antibodies and renal disease, (A1) 20
 - antineutrophil cytoplasmic antibodies, 625
 - antiphospholipid antibodies from patients infected with human B19 parvovirus have similar specificity and co-factor dependence, (A1) 22
 - and antiphospholipid syndrome, obstetric outcome: the St Thomas' Hospital Lupus in Pregnancy clinic experience, (A1) 123
 - association of the SLICC/ACR damage index with measures of disease activity and health status, (A1) 29
 - causes of death in two hundred patients, (A1) 37
 - clinical, arthroscopic and histological features of synovitis, (A1) 36
 - clinical significance of non-precipitating anti-nRNP antibodies, (A1) 18
 - comparison of disease activity of ethnic groups belong to 2 university hospitals, (A1) 34
 - comparison of the SF-20 and SF-36 in the assessment of quality of life, (A1) 31
 - complement and immune complexes, 5
 - cyclosporin A in the treatment: results of an open clinical study, 669
 - diagnostic relevance of anti-Sm antibodies detected by ELISA, (A1) 19
 - disease activity vs illness perception as predictors of health status and psychological well-being: longitudinal analysis, (A1) 34
 - dissociation of immune response to the degraded and nondegraded (native) Sa antigens, (A1) 18
 - FcgammaRIIA polymorphism, (A1) 28
 - fibrinogen in: an atherothrombotic link?, (A1) 122
 - further characterization of anti-endothelial cell antibodies by controlled immunoblotting, 1231
 - further validity of the BILAG disease activity index, (A1) 29
 - genetic analysis of TAP2 in patients from two ethnic groups, 529
 - HLA-DRB1 and complement C4 null alleles in a Spanish population, (A1) 26
 - human immunodeficiency virus infection, (A2) 35
 - IgG FcgammaRIIA polymorphism: genotypic analysis in Caucasian, Chinese and Afro-Caribbean patients, (A1) 27
 - influence of coping strategy on outcome, (A1) 33
 - integrin-induced T-cell signaling: a novel tool for disease monitoring, (A1) 38
 - multiple cryptococcal brain abscesses, (L) 1334
 - murine, analysis of apoptosis in lymphoid organs and lupus disease, (A1) 40
 - neurological manifestations in Saudi patients, (A1) 32
 - neuropsychiatric
 - Lba5 and ribosomal P autoantibodies, (A1) 111
 - pulse cyclophosphamide in the treatment, (A1) 32
 - ovulation-induction therapy risks, 1184

- peripheral lymphocyte apoptosis and bcl-2 expression: correlation with disease activity, (A1) 39
- polymorphism of the extracellular domain of the human Fas gene, (A1) 25
- predominance of IgM anti-U1RNP antibodies, 534
- prevalence and clinical significance of antibodies to ribonucleoproteins in Malaysia, 129
- previously undiagnosed in acute psychiatric hospital admissions, (A1) 36
- quality of life is influenced by disease activity but is impaired in all SLE patients, (A1) 31
- red cell complement receptor 1 deficiency in southern Chinese: correlation with disease activity, (A1) 26
- salmonella infection in Saudi patients, (A1) 35
- serum sex hormones in pregnancy: comparison between lupus, rheumatoid and control patients, (A1) 35
- SLICC/ACR damage index is valid, and renal and pulmonary organ scores are predictors of severe outcome, 248
- stroke in, (A1) 33
- sulphasalazine-induced in a patient with rheumatoid arthritis, (L) 104
- thyroid autoimmunity in: the clinical significance of a fluctuating course, (L) 195
- TNF A2 B3 and D2 microsatellite allele frequencies are increased and correlate with clinical outcome, (A1) 25
- TNFA promoter polymorphisms show interethnic differences, (A1) 28
- trabecular and cortical bone loss, 244
- tumour necrosis factor alpha and its soluble receptors parallel clinical disease and autoimmune activity, 1067
- ultraviolet irradiation: friend or foe?, 1002
- Systemic mast cell disease, severe osteoporosis due to: successful treatment with interferon alpha-2b, 898
- Systemic sclerosis
- antibodies to RNA polymerases: clinical and serological associations, (A1) 46
 - ascorbic acid absorption, (A1) 43
 - comparison of methotrexate with placebo in treatment: a 24 week randomized double-blind trial, followed by a 24 week observation trial, 364
 - HLA-DQB1 polymorphism and antibodies to RNA polymerases, (A1) 45
 - intermittent pulse prednisolone and cyclophosphamide for the treatment of interstitial lung disease, (A2) 27
 - limited cutaneous, contemporaneous activation of endothelial responses and cellular immunity, (A2) 27
 - longitudinal study of anti-endothelial cell antibody activity, (A1) 46
 - in men: clinical and immunological differences, (L) 910
 - non-invasive assessment of vascular reactivity in forearm skin, 1281
 - pancreatic function in, (L) 298
 - and primary Raynaud's phenomenon
 - digital and microvascular blood flow responses, (A1) 42
 - digital neurovascular mechanisms as assessed by iontophoresis and laser Doppler techniques, (A1) 48
 - increased nailfold capillary dimensions, 1127
 - psychological and psychopathological evaluation of patients, (A1) 47
 - reduced levels of soluble L-selectin, (A1) 42
 - renal tubular dysfunction is associated with endothelial damage, (A1) 44
 - risk factors for progression of oesophageal dysmotility, (A1) 44
 - total skin score as an independent outcome variable—survey of biases of clinical experts, (A1) 45
- Systemic vasculitis, intravenous immunoglobulin as sole therapy, 1150
- SZETO CC, CHOW CC, LI KY, KO TC, YEUNG VTF, COCKRAM CS. Thyrotoxicosis and renal tubular acidosis presenting as hypokalaemic paralysis, 289
- SZYMOWIAK CH see Johnston TW
- T cell clones
- from a subset of patients with primary Sjögren's syndrome produce high levels of IL-10, (A1) 16
 - synovial fluid derived respond to *Chlamydia trachomatis* protein antigens, (A1) 114
- T cells
- autoreactive of RA patients recognize a new 68-k autoantigen, (A1) 107
 - integrin-induced signaling in systemic lupus erythematosus: a novel tool for disease monitoring, (A1) 38
 - naive/resting, active systemic lupus erythematosus is associated with recruitment, 226
- T lymphocyte
- expansions in blood and synovial fluid in reactive arthritis, (A1) 183
 - immunophenotyping in polymyositis and dermatomyositis, 839
- T-cell receptor variable gene segment TCRBV12S4 null allele frequency in UK juvenile chronic arthritis, (A1) 80
- T-cell repertoire of infiltrating T cells in lachrymal glands, salivary glands and kidneys from alymphoplasia (*ALY*) mutant mice: a new model for Sjögren's syndrome, 1223
- TAAL E see Brus HLM; Riemsma RP; van der Wardt AM
- TABUENCA MJ see Isasi C; Perez A
- TADA S see Kobayashi H
- TAIT D see Loizou S
- TAK P see Paleolog E
- TAK PP see Dolhain RJEM
- TAKASUGI K see Kobayashi H
- TALAMO J, WILKINSON P, COX N, DAVIES P, DEVLIN J, DIXEY J, EMERY P, GALLIVAN S, GOUGH A, JAMES D, PROUSE P, WILLIAMS P, WINFIELD J, YOUNG A. Socio-economic factors in rheumatoid arthritis: effect on functional outcome at three years in 516 patients, (A1) 234
- TALBOT H see Luqmani RA
- TAN PLJ see Armstrong GP
- TANI M see Nishikai M
- TANI Y see Tiwana H
- TANKURT E. Resolving familial Mediterranean fever attacks with interferon alpha, (L) 1188
- TANQUERAY AB see Ismail AA
- TAP2 genetic analysis in systemic lupus erythematosus patients from two ethnic groups, 529
- TAP genes, MHC-encoded in rheumatic diseases, (E) 501
- TARASEVICIUS T see Malaviya AN
- Tartrate-resistant acid phosphatase-positive multinuclear cells, bone resorption by isolated from rheumatoid synovium, 213
- TAVERNIER C see Maillefert JF
- tax gene of HTLV-1 is not present in the salivary glands of patients with idiopathic Sjögren's syndrome, (A1) 16
- TAYLOR A see Kennedy LG
- TAYLOR AL, BALAKRISHNAN C, CALIN A. Risk factors for worse disease activity and function in ankylosing spondylitis, (A2) 24
- TAYLOR AL see also Haigh RC
- TAYLOR DJ, FARAGHER EB, SHARIF M, KIRWAN J. Effect of low dose steroid treatment on serum stromelysin and collagenase levels in RA patients followed for 3 years, (A1) 210
- TAYLOR DJ see also Cheung NT
- TAYLOR H see Mishra MB
- TAYLOR HG see Avers HL
- TAYLOR JC, DOYLE DV, THOMPSON AV, MULCAHY B, WILTSHIRE P
- Compston DEXA scanning guidelines may underestimate community needs, (A1) 84
 - Low bone density in patients falling outside Compston guidelines for DEXA scanning, (A1) 83
- TAYLOR P see Paleolog E
- TAYLOR PC, CHU CQ, PLATER-ZYBERK C, MAINI RN. Transfer of type II collagen-induced arthritis from DBA/1 to SCID mice can be prevented by blockade of MAC-1, (A1) 74
- TAYLOR PC see also Williams DG
- ^{99m}Tc human immunoglobulin (IgG), detection of active rheumatoid arthritis, (A2) 15
- tch gene
- expression in mouse embryos, (A2) 8
 - a mouse dwarfing gene, mapping, (A2) 9
- TCR restricted expression on synovial fluid CD4 and CD8 T cells in psoriatic arthritis, (A1) 160
- TEBIB J see Combe B
- TEBIB JG see Maillefert JF
- TEH LS, JOHNS CW, SHAFFER JL, RIELEY F, BAKER E, AARONS L, BENNETT RJM, HERRICK AL, JAYSON MIV. Ascorbic acid absorption in patients with systemic sclerosis, (A1) 43
- TEH LS see also Coombes GM; Sanders PA
- TEIXIDÓ M see Navarro M
- Telephone follow-up service, survey of ambulatory patients' attitudes towards, (A1) 130
- Telomere reduction in scleroderma patients: a possible cause for chromosomal instability, 732
- TEMPL E, KOELLER M, RIEDLE M, WAGNER O, GRANINGER W, LUGER A. Anterior pituitary function in patients with newly diagnosed rheumatoid arthritis, 350
- Temporal arteritis and polymyalgia rheumatica, prospective study of 287 patients, 1161
- Temporal artery biopsy results review in a large DGH—ophthalmic signs increase the positive biopsy rate, (A1) 56
- TEN BERGE M see van der Net J
- TENA X see Acebes JC; Olivé A
- Tenascin, serum levels in patients with rheumatoid and psoriatic arthritis, (A1) 176
- Tenidap
- decreases synovial growth (hyperplasia and increased extra-cellular matrix production) and cell infiltration in rabbits with chronic arthritis, (A1) 72
 - reduces cell infiltration and MCP-1 mRNA expression in a rabbit model of acute arthritis, (A1) 73
 - versus diclofenac in rheumatoid arthritis, (L) 602

- TENNANT A, HILLMAN M, FEAR J, PICKERING A, CHAMBERLAIN MA. Are we making the most of the Stanford Health Assessment Questionnaire, 574
- Tenosynovitis
of the hand: major determinant of functional ability in rheumatoid arthritis, (A1) 202
of the wrist, *Mycobacterium xenopi*—an unusual presentation in an immunocompetent patient, 1008
- Tenoxicam and bromazepam comparison in the treatment of fibromyalgia. A randomized, double-blind, placebo-controlled trial, (A1) 138
- TER HAAR NT see Dolhain RJEM
- TERATO K, YE XJ, MIYAHARA H, CREMER MA, GRIFFITHS MM. Induction of chronic autoimmune arthritis in DBA/1 mice by oral administration of type II collagen and *Escherichia coli* lipopolysaccharide, 828
- TERVAHARTIALA P see Pettersson T
- Testosterone
randomized trial of therapy in males with rheumatoid arthritis, 568
salivary and serum prolactin in young and middle-aged men with ankylosing spondylitis, (A1) 152
- TGF β , IL-1 α and retinoic acid effects on homeostasis of aggrecan and biglycan from articular cartilage, (A1) 7
- Th1 and Th2 cytokines in rheumatoid arthritis, (A1) 179
- THE LS see Davies EJ
- THEODORIDIS E see Davies KA
- Therapeutic trials, approach in rheumatology: the foundation for applied rheumatology research, 1000
- Thermal stress testing and video nail-fold capillaroscopy in children with Raynaud's phenomenon, (A2) 26
- Thermographic changes in patients with work-related chronic upper limb pain, (A1) 139
- THIELE B see Riemekasten G
- Third hypervariable region (HVR3), joint involvement in connective tissue disease, (A1) 30
- THOMAS E see Macfarlane GJ
- THOMPSON AV see Masud T; Taylor JC
- THOMPSON EM see Davies KA
- THOMPSON PW see Carr AJ; Dunne CA; Harwood RH; Shearing-Lynne C
- THOMSON EA see Capell HA; Duncan MR; McEntegart A; Morrison E; Porter DR
- Thrombocytopenia, refractory, successful therapy with danazol associated with autoimmune diseases, (A1) 55
- Thrombosis in Churg-Strauss syndrome. Beyond vasculitis, 1181
- Thyroid autoimmunity in systemic lupus erythematosus: the clinical significance of a fluctuating course, (L) 195
- Thyrototoxicosis and renal tubular acidosis presenting as hypokalaemic paralysis, 289
- Tibial stress fractures, simultaneous bilateral in a 15-year-old milkman—a case report, 905
- TICKLY M see Rudwaleit M
- TILL SH, AMOS RS. Neurofibromatosis masquerading as pauciarticular juvenile chronic arthritis, (A2) 39
- TINAHONES F see Collantes E
- TISHLER M see Ehrenfeld M
- Tissue inhibitor of metalloproteinases (TIMP-1)
gene promoter mapping, (A1) 68
inactivation and fragmentation by peroxynitrite, (A1) 69
- TIWANA H, TANI Y, HUKUDA S, NISHIOKA J, FIELDER M, RASHID T, WILSON C, BANSAL S, EBRINGER A. Antibodies to Klebsiella, Proteus and the peptides HLA-B27 and HLA-DR4 (DW15) in Japanese patients with ankylosing spondylitis and rheumatoid arthritis, (A2) 12
- TIWANA H, WILSON C, CUNNINGHAM P, BINDER A, EBRINGER A. Antibodies to four Gram-negative bacteria in rheumatoid arthritis which share sequences with the rheumatoid arthritis susceptibility motif, 592
- TNF A2 B3 and D2 microsatellite allele frequencies are increased in SLE and correlate with clinical outcome, (A1) 25
- TNF receptor promoter region, murine p75: DNA sequence and characterization of a cis-acting silencer, (A1) 9
- TNF-HLA haplotypes, extended, association with rheumatoid arthritis in males and females, (A1) 189
- TOBIAS JH, DALZELL N, PAZIANIS M, CHAMBERS TJ. Effect of cyclical didronel on bone loss in early postmenopausal women, (A1) 87
- TOBIAS JH, LAVERSUCH CJ, CHAMBERS TJ, GALLAGHER AC. Aminohexane bisphosphonate suppresses bone turnover in postmenopausal women more rapidly than oestrogen-gestagen therapy, 636
- TOBIAS JH see also Gallagher AC
- TODD A see McGonagle D; Ormond P
- TODD C see O'Neill TW
- TODESCO S see Fiocco U
- TOIVANEN A, TOIVANEN P. Current trends in Finnish rheumatology, 489
- TOIVANEN A see also Roivainen A
- TOIVANEN P see Isomäki P; Roivainen A; Toivanen A
- TOJO T see Nishikai M
- Tolerability
local tissue of meloxicam, a new NSAID: indications for parenteral, dermal and mucosal administration, (A1) 44
of multiple administration of intramuscular meloxicam: comparison with intramuscular piroxicam in patients with rheumatoid arthritis or osteoarthritis, (A1) 51
and safety, open study to assess meloxicam 15 mg in subjects with rheumatic disease and mild renal impairment, (A1) 56
- TOMAZIC S see Demars F
- TOMSIC M see Jevtic V
- TORISU T see Fujikawa Y
- TORRE-CISNEROS J see Gonzalez-Dominguez J
- TORRESANO M see Perez A
- TOSI S see Marchesoni A
- TOURAY M, LEMARCHAND P, SO A. Expression of MT-MMP1 by human fibroblast-like synoviocytes and its regulation by inflammatory cytokines, (A1) 67
- TOVAR JV see Navarro FJ
- Trabecular and cortical bone loss in systemic lupus erythematosus, 244
- TRACH V see Stei P
- Training
in rheumatology, (L) 1190
specialist, into the new millennium: the price of conformity with Europe, (E) 607
- TRATTNIG S see Kainberger F
- Treadmill and walking speed tests, how well do they reflect 24 hour activity?, (A1) 194
- Treatment
advances—when and how to inform the public, 1175
of rheumatoid arthritis—does it affect society's cost for the disease?, 791
- TREBELLAHR G see Riemekasten G
- TREGNAGHI A see Fiocco U
- Triamcinolone hexacetonide, comparison of glenohumeral injections in the painful rheumatic shoulder, (A1) 132
- TRIANTAPHYLLOPOULOS K see Chernajovsky Y
- TRIBBICK G see Laversuch CJ
- TRON AM see Hellio le Graverand MP
- TROUGHTON PR, PLATT R, BIRD H, EL-MANZALAWI E, BASSIOUNI M, WRIGHT V. Synovial fluid interleukin-8 and neutrophil function in rheumatoid arthritis and seronegative polyarthritis, 1244
- TSG-6: a hyaluronan binding matrix protein upregulated by cytokines in joint disease, (A1) 236
- TSIKAS D see Stichtenoth DO
- TSUBATA R see Furukawa M
- TSUBATA T see Furukawa M
- TSUKAMOTO M see Matsuda J
- TSUKAZAKI T see Okano K
- Tuberculosis, osteoarticular in two different populations: the HIV infection impact, (A1) 166
- Tumoral calcinosis: a case report, (A2) 35
- Tumour necrosis factor, paradoxical effects of a synthetic metalloproteinase inhibitor which blocks both p55 and p75 TNF receptor shedding and TNF α processing in RA synovial membrane cell cultures, (A1) 181
- Tumour necrosis factor α
antibody, treatment of rheumatoid arthritis decreases expression and shedding of E-selectin, (A1) 218
blockade, changes in naturally-occurring cytokine inhibitor levels after in rheumatoid arthritis, (A1) 181
comparison of two bisphosphonates on production by PBMCs, (A1) 4
feritin H-chain mRNA is upregulated in human articular cartilage stimulated with, (A1) 4
and its soluble receptors parallel clinical disease and autoimmune activity in systemic lupus erythematosus, 1067
promoter polymorphisms show interethnic differences in association with SLE, (A1) 28
release from human monocytes: a common mechanism of lupus-inducing drug immunomodulation, (A1) 3
and ricin A-chain, synthesis of an active cytotoxin by cross-linking, (A1) 3
role as an inhibitor of erythropoiesis in rheumatoid arthritis: insights from TNF α blockade *in vivo*, (A1) 217
- Tumour necrosis factor α -induced resorption of mouse calvariae *in vitro* is suppressed by xanthine oxidase inhibitors, (A1) 241
- Tumour necrosis factor receptor, human p75, mutational analysis of the membrane proximal cleavage site, (A1) 109
- TUNN EJ, ALCOCK J, HUSSEY R. Developing for low back pain and sciatica a care path from primary through secondary to tertiary care, (A1) 143
- TÜRK D, ROTH W, BUSCH U. Review of clinical pharmacokinetics of meloxicam, (S1) 13
- TURNER HE, MYSZOR MF, BRADLOW A, DAVID J. Lupus or lupoid hepatitis with mesenteric vasculitis, 1309
- TUSZEWSKI M see Braun J
- TUZÜN Y see Yazici H

- TYLER JA *see* Chevalier X
- Tyrosine nitration in the inflamed human synovium: evidence for the generation of peroxynitrite *in vivo*, (A1) 70
- Tyrosine phosphorylation in neutrophils from synovial fluid of patients with rheumatoid arthritis, 846
- TZIOUFAS AG *see* Vlachoyiannopoulos PG
- UEBELHART D *see* Hellio le Graverand MP
- UGARTEMENDIA C *see* Moragues-Pastor C
- ULRICHS T *see* Baerwald CGO
- Ultrasound measurements in glucocorticoid treated patients, (A2) 5
- Ultraviolet irradiation in systemic lupus erythematosus: friend or foe?, 1002
- UMAMAHESWARAN I *see* Malaviya AN
- Umbilical vein, human and dermal microvascular display heterogeneity in expression of surface antigens following activation of protein kinase, (A1) 11
- Undergraduate education in rheumatology, (E) 110, (E) 203
- UNSWORTH A *see* Speed CA
- Upper limb pain, chronic work-related, thermographic changes, (A1) 139
- Urate crystal-induced inflammation, characterization of E-selectin expression, leucocyte traffic and clinical sequelae, 323
- Urease, bacterial, increased activity in faeces in juvenile chronic arthritis: evidence of altered intestinal microflora?, 689
- USON J, Balsa A, Pascual-Salcedo D, Cabezas JA, Gijón-Baños J. Serum and synovial fluid soluble IL-6 receptor in patients with different arthropathies, (A1) 9
- UTHMAN I, Vazquez-Abad D, Senécal J-L. Distinctive features of idiopathic inflammatory myopathies in French Canadians, (A1) 49
- Uveitis
- chronic anterior, prevalence in New Zealand children with juvenile chronic arthritis, (A2) 25
 - juvenile chronic arthritis-associated, antibodies against retinal S-antigen, (L) 101
- VAISHNAV AK *see* Slingsby JH
- VAKALIS AN, Hull RJ, Hodgkins PR, Parnaby-Price A, Jaffrey MN, Evans AR. Review of temporal artery biopsy results in a large DGH—ophthalmic signs increase the positive biopsy rate, (A1) 56
- VAL-BERNAL F, González-Vela C, Garjón F, Mayorga M. Isolated polyarthritis nodosa of the prostate, (A1) 51
- VALENTINI G *see* Matucci Cericini M
- Valgus heel deformity and plantar pressure distribution of the forefoot in rheumatoid arthritis, (A1) 199
- Validity
- construct of EuroQol in 240 patients with rheumatoid arthritis stratified by functional class, (A2) 18
 - testing of the MOS SF36 generic health profile in rheumatoid arthritis, (A2) 17
- VALKENBERG HA *see* Odding E
- VALVERDE J, Clavaguera MT, Pérez A, Cabo X, Narváez FJ, Juanola X. Septic bursitis: predisposing factors, clinical and therapeutic features, (A1) 165
- VAN DE LAAR MAFJ *see* Brus HLM
- VAN DE PUTTE L *see* Stolk J
- VAN DE PUTTE LBA *see* Creemers MCW; Emery P; Prevoo MLL; van Riel PLCM
- VAN DEN HEUVEL WJA *see* Smedstad LM
- VAN DEN HOOGEN FHJ, Boerbooms AMT, Swaak AJG, Rasker JJ, van Lier HJJ, van de Putte LBA. Comparison of methotrexate with placebo in the treatment of systemic sclerosis: a 24 week randomized double-blind trial, followed by a 24 week observation trial, 364
- VAN DER LINDEN SJ *see* Boonen A
- VAN DER NET J, Prakken ABJ, Helders PJM, ten Berge M, van Herwaarden M, Sinnema G, de Wilde EJ, Kuis W. Correlates of disability in polyarticular juvenile chronic arthritis—a cross-sectional study, 91
- VAN DER PUTTE LBA *see* van den Hoogen FHJ
- VAN DER VEEN MJ, Scheven BAA, van Roy JLAM, Damen CA, Laféber FPJG, Bulsma JWJ. *In vitro* effects of methotrexate on human articular cartilage and bone-derived osteoblasts, 342
- VAN DER WARDT AM, Taal E, Rasker JJ, Wiegman O. General public's perceptions about rheumatic diseases, (A1) 126
- VAN DUKE CF *see* Demsar F
- VAN GESTEL AM *see* Prevoo MLL; van Riel PLCM
- VAN HERWAARDEN M *see* van der Net J
- VAN LANGELAAN EJ *see* Rondoy HK
- VAN LIER HJJ *see* van den Hoogen FHJ
- VAN MARLE J *see* Verschure PJ
- VAN MUIJEN GNP *see* Rondoy HK
- VAN PAASSEN HC *see* Riemsmma RP
- VAN RIEL PLCM, van Gestel AM, van de Putte LBA. Development and validation of response criteria in rheumatoid arthritis: steps towards an international consensus on prognostic markers, (S2) 4
- VAN RIEL PLCM *see also* Creemers MCW; Prevoo MLL
- VAN RIJSWIJK MH *see* Prevoo MLL
- VAN ROERMUND PM *see* Van Valburg AA
- VAN ROON JAG *see* Verhoef CM
- VAN ROY HJAM *see* Van Valburg AA
- VAN ROY JLAM *see* van der Veen MH
- VAN VALBURG AA, van Roermund PM, van Roy HJAM, Laféber FPJG, Bulsma JWJ. Actual repair of cartilage by Ilizarov joint distraction, tested in the Pond-Nuki dog model for osteoarthritis, (A1) 104
- VANDEN BERG WB *see* Verschure PJ
- VANDENBROUCKE JP *see* Vliet Vlieland TPM
- VANDENOUWELAND FA *see* Odding E
- VANDENROUCKE JP *see* Simon CH
- VANE JR. Mechanism of action of NSAIDs, (S1) 1
- VANT HOF MA *see* Creemers MCW; Prevoo MLL
- VARLOW J *see* O'Neill TW
- Vascular endothelial growth factor, angiogenesis: a critical process in the pathogenesis of RA, (E) 917
- Vasculitis
- damage index, (A1) 53
 - eruption during treatment with leflunomide for rheumatoid arthritis, (A1) 226
 - integrated total assessment log (vital): computerised system for assessment of vasculitis, (A1) 54
 - localised cutaneous, incidence and clinical associations of biopsy proven in the Norwich Health Authority, (A1) 54
 - mesenteric, lupoid hepatitis or lupus, 1309
 - reduced levels of soluble L-selectin, (A1) 42
 - relationship between soluble adhesion molecules and anti-endothelial cell antibodies, (A1) 52
- VAUGHAN R *see* Hameed K
- VAZQUEZ-ABAD D *see* Uthman I
- VEALE DJ *see* Reece RJ
- VEDHARA K *see* Lawson TM
- VEGAZO O *see* Carreño L
- VEIGA R *see* De La Mata J
- VÉLITAT P *see* Ghoslan PR
- VENABLES PJW. Polymyositis-associated overlap syndrome, (E) 305
- VENABLES PJW *see also* Cooke SP; Horsfall AC; Price EJ; Rigby S
- VENN G, Bayliss M, Hardingham T. Effects of TGF β , IL-1 α and retinoic acid on homeostasis of aggrecan and biglycan from articular cartilage, (A1) 7
- VENNART W *see* Gandy S; Gasson J
- Venous thrombosis, coagulation factor V gene mutation increases the risk in Behçet's disease, 1178
- VERA-LLAÓ R *see* Moragues-Pastor C
- VERHEIJEN JH *see* Rondoy HK
- VERHOEF CM, van Roon JAG, Bruinzeel CAFM, Laféber FPJG, Bulsma JWJ. Low prevalence of hay fever in patients with rheumatoid arthritis, (A1) 197
- VERRONEN P *see* Malin M
- VERSHURE PJ, van Marle J, Joosten LAB, Helsen MMA, Laféber FPJG, van den Berg WB. Localization of insulin-like growth factor-I receptor in human normal and osteoarthritic cartilage in relation to proteoglycan synthesis and content, 1044
- Vertebral deformity
- effect of physical activity on the risk, (A1) 89
 - hormonal and gynaecological factors and risk, (A1) 90
- VETTERLEIN O *see* Choy EHS
- V β gene, single anti-DNA-related encodes for multiple specificities: the influence of the light chain, (A1) 23
- Video image analysis of hands: development of an anatomic index as a potential outcome measure in rheumatoid arthritis, 1274
- VIGNON D *see* Lioté F
- VIGNON E *see* Conrozier T; Hellio le Graverand MP
- VILA L *see* Amat M; Camacho M
- VILJOEN D *see* Beighton PH
- VINCENT C *see* Houssien D
- VISCHER TL *see* Bas S
- VISIONI RA *see* Ciconelli RM
- Vitamin D receptor polymorphisms
- and bone turnover in female twins, (A1) 88
 - and osteoarthritis, (A1) 117
- VITTETA E *see* Gould DJ
- VIU P *see* Bologna C
- VIVANCOS J *see* García-Carrasco M
- Vlachoyiannopoulos PG, Gualis A, Tzioufas AG, Moutsopoulos HM. Predominance of IgM anti-U1RNP antibodies in patients with systemic lupus erythematosus, 534
- VLIET VLIELAND TPM, Zwiderman AH, Hazes JMW. Measurement of morning stiffness in rheumatoid arthritis clinical trials, (A1) 197
- VLIET VLIELAND TPM, Zwiderman AH, Vandenbroucke JP, Breedveld FC, Hazes JMW. Randomized clinical trial of in-patient

- multidisciplinary treatment versus routine out-patient care in active rheumatoid arthritis, 475
- VNTR loci, DNA allelic alterations within of scleroderma families, 1216
- VOCK P *see* Lehmann T
- VON KÄNEL S *see* Weber M
- WADA M, IMURA S, BABA H, SHIMADA S. Knee laxity in patients with osteoarthritis and rheumatoid arthritis, 560
- WADE AJ *see* Carbon RJ
- WADE M *see* Walsh DA
- WAGNER B *see* Stichenothe DO
- WAGNER O *see* Templ E
- WAHLE M *see* Baerwald CGO
- Waiting list
- orthopaedic, rheumatology initiative to reduce, (A1) 129
 - removal of patients from, (A1) 129
- WALDRON HA. Osteoarthritis of the hands in early populations, 1291
- WALKER DJ, HESLOP P, WRIGHT D, CLEARY R, ESSEX T, CHANDLER C. 24hr activity in rheumatoid arthritis: comparison with HAQ, (A1) 233
- WALKER DJ, HESLOP PS, CHANDLER C, MORANT SV, KIFF PS, STEAD H. Comparison of effect of Arthrotec 75 and Diclofenac 75 on patient activity using the Numact activity monitor, (A2) 33
- WALKER DJ *see also* Attea G; Chatterjee S; Davidson BKS; Kay LJ; McDonagh JE; Speed CA
- WALKER P *see* Maillefert JF
- Walking speed and treadmill tests, how well do they reflect 24 hour activity?, (A1) 194
- Walking time may be a very useful measure of clinical laboratory and radiological status in patients with rheumatoid arthritis, (A1) 195
- WALMSLEY M *see* Mauri C
- WALNE AJ, JAMES IT, COOKE N, PERRETT D. Relationships between clinical measures of disease activity and biochemical markers in rheumatoid arthritis, (A1) 230
- WALNE AJ *see also* James IT
- WALPORT M *see* Ocal L
- WALPORT MJ *see* Athanassiou P; Davies KA; Hogarth MB; Loizou S; Naughton M; Slingsby JH
- WALSH D *see* Gaffney K
- WALSH DA, CATRAVAS J, WHARTON J. Synovial angiotensin converting enzyme in human arthritis, (A1) 66
- WALSH DA, WADE M, HU DE, MAPP PI, BLAKE DR, FAN T-PD. Endothelial proliferation index in rat and human tissues, (A1) 10
- WALSH HPJ *see* Lees C
- WALSH S *see* Hardwick L; Hardwick LJ
- WANG CL, OOI L, WANG F. Prevalence and clinical significance of antibodies to ribonucleoproteins in systemic lupus erythematosus in Malaysia, 129
- WANG EY *see* Lawson TM
- WANG F *see* Wang CL
- WARD D *see* Ellis AJ
- WARD M *see* Wilkinson NZ
- WARD ME *see* Campbell F
- WARNOCK MG, LAYCOCK CA, DIAMOND AG, GOODACRE JA. Production of cartilage aggrecan domains as recombinant fusion proteins to analyse recognition by autoreactive T lymphocytes, (A1) 184
- WASSIF WS *see* Preedy VR
- Water, high mineral content, value in the short-term management of rheumatoid arthritis, (A1) 228
- WATERTON J *see* Creamer P
- WATKIN SW *see* Reid AJC
- WATT I. Role of MRI in the assessment of rheumatological disease, (S3) 1
- WATT I *see also* Creamer P; Jevtic V; Oliver C; Poleksic L; Sharma S
- WATTS RA *see* Carruthers DM; Jolliffe VA; Reid AJC
- WEBB GR, WESTACOTT CI, ELSON CJ. Osteoarthritic synovium induces an up-regulation of p55 TNF α receptor expression on non-arthritis chondrocytes, (A1) 237
- WEBB LMC *see* Price EJ
- WEBER H *see* Mau W
- WEBER M, CEDRASCHI C, ROUX E, KISSLING RO, VON KÄNEL S, DALVIT G. Prospective controlled study of low back school in the general population, 178
- WEBLEY M *see* Avernus HL
- WECHSLER B *see* Le Thi Huong D
- WEDEKIND F, SCHNEIDER M. Atypical synovial cells are involved in collagen type 1 mRNA-expression in rheumatoid arthritis, (A1) 178
- WEDEKIND F, SCHNEIDER M, PAULY T, FRIEMANN J, BURMEISTER G. Distribution of 27E10 macrophages in synovium of inflammatory and degenerative joint disease, (A1) 178
- WEDLEY JR *see* Dolan AL
- WEEKS D *see* Hall FC
- WEERASINGHE D *see* Rigby S
- Wegener's granulomatosis
- increasing incidence or increased recognition?, 142
 - silent myocardial infarction, 188
- WEIDEMANN HF *see* Mau W
- Weight loss and body composition in rheumatoid arthritis, (A1) 200
- WEINBLATT ME. Methotrexate in rheumatoid arthritis: toxicity issues, (E) 403
- WEINERTH J *see* Stoffel MP
- WEINERTH JD. Are antineutrophil cytoplasmic antibodies associated with spondylarthropathies, (L) 1032
- WELLS TM *see* Dudhia J
- WELSH K *see* McHugh NJ
- WELSH KI *see* Artlett CM
- WENDLING D. Combination therapy of anti-CD4 and anti-IL6 monoclonal antibodies in a case of severe spondylarthropathy, (L) 1330
- WEST J *see* Grahame R
- WESTACOTT CI *see* Webb GR
- WESTHOVEN R *see* Dequeker J
- WESTWOOD OMR *see* Laversuch CJ
- WEYL-CLERC D *see* Lioté F
- WHALE M *see* Baerwald CGO
- WHALLEY GA *see* Armstrong GP
- WHARTON J *see* Walsh DA
- WHEELER SK *see* Perretti M
- WHITE PH. Future expectations: adolescents with rheumatic diseases and their transition into adulthood, (E) 80
- WHITELAW D *see* Mack CS
- WHYTE J *see* Patel YI
- WICHERT P *see* Baerwald CGO
- WICHERT P von *see* Baerwald CGO
- WIEGLEB J *see* Stei P
- WIEGMAN O *see* Brus HLM; Riemsma RP; van der Wardt AM
- WILKINSON M *see* Carr AJ
- WILKINSON NZ, KINGSLEY GH, SIEPER J, BRAUN J, WARD M. Detection of *Chlamydia pneumoniae* and *Chlamydia trachomatis* in the synovium of patients with a range of rheumatic diseases, (A1) 162
- WILKINSON P *see* Talamo J
- WILKINSON PC *see* McInnes IB
- WILLIAMS AS, CAMILLERI JP, GOODFELLOW RM, WILLIAMS BD. Single intra-articular injection of liposomally conjugated methotrexate suppresses joint inflammation in rat antigen-induced arthritis, 719
- WILLIAMS AS *see also* Goodfellow RM
- WILLIAMS B *see* Barlow JH
- WILLIAMS BD *see* Byrne PAC; Goodfellow RM; Lawson TM; Lloyds D; Plant MJ; Richards PJ; Williams AS
- WILLIAMS C *see* Hardwick L
- WILLIAMS DG, TAYLOR PC. Clonally expanded plasma cells in rheumatoid synovium, (A1) 172
- WILLIAMS GP *see* Plant MJ
- WILLIAMS H *see* Martin JC
- WILLIAMS L *see* Brennan FM
- WILLIAMS MH *see* Carr AJ
- WILLIAMS P *see* Prouse P; Talamo J
- WILLIAMS PL *see* Bamji AN
- WILLIAMS R *see* Hawkins PN
- WILLIAMS RO *see* Marinova L; Mauri C; Ross S
- WILLIAMS SML *see* Evans MD
- WILLIAMSON L, JONES M, MOWAT AG. Use of steroids in patients undergoing lower limb arthroplasty, (A1) 209
- WILLOUGHBY M *see* Mills SY
- WILSON C *see* Tiwana H
- WILTSHIRE P *see* Taylor JC
- WIND K *see* Dasgupta B
- WINEARLS CG *see* Hawkins PN
- WINFIELD J *see* Prouse P; Talamo J
- WYNARD P *see* Nazhat NB
- WYNARD PG *see* Zhang Z
- WISNOM MR *see* Adams MA
- WITTE K *see* Hoffmeyer F
- WOJTULEWSKI JA, SCHATTEKIRCHNER M, BARCELÓ P, LE LOET X, BEVIS PJR, BLUMKI E, DISTEL M. Six-month double-blind trial to compare the efficacy and safety of meloxicam 7.5 mg and naproxen 750 mg in patients with rheumatoid arthritis, (S1) 22
- WOJTULEWSKI JA, SCHATTEKIRCHNER M, BARCELÓ P, LE LOET X, NORTH N, BLUMKI E. Lower gastrointestinal and renal side effects of the COX-2 inhibitor meloxicam, compared to naproxen, in rheumatoid arthritis patients, (A1) 223
- WOLFE F *see* Jones M; Jones MA
- WOLMAN R. Orthopaedic medicine, the remit of the rheumatologist, (E) 1
- WONG WM, ELLWOOD CN, HOWELL WM, SMITH JL, CAWLEY MID. Rheumatoid T cell functional deficiency can be reversed with trispecific antibody stimulation, (A1) 13
- Woo P
- British Paediatric Rheumatology Group news, 686, 1015
 - Paediatric rheumatology course, 687
- Woo P *see also* Fishman D

- WOODBURN J, HELLIWELL PS. Valgus heel deformity and plantar pressure distribution of the forefoot in rheumatoid arthritis, (A1) 199
- WOODCOCK JP *see* Plant MJ
- WOODFORD-RICHENS KL *see* Keen RW
- WOODHOUSE S *see* Sharif M
- WOODHOUSE SM *see* Sharif M
- WOODROW DF *see* Allard SA
- WOODS RJ, DAVID J, BAIGENT S, GIBBINS J, LOWRY PJ. Elevated levels of corticotrophin-releasing factor binding protein in the blood of patients suffering from arthritis and septicaemia and the presence of novel ligands in synovial fluid, 120
- WOODS Z *see* Leak AM
- WOODY J *see* Paleolog E
- WOLF AD *see* Deodhar AA; O'Neill TW
- WOOLHEAD GM *see* Carr AJ
- WORDSWORTH BP *see* Allen RL; Brown MA; Hall FC
- WORDSWORTH BPW *see* Hardwick LJ
- WORDSWORTH P *see* Hardwick L; Rudwaleit M
- Workload, Rheumatology outpatient increases inexorably, (A2) 29
- WORKALL JG *see* Dasgupta B
- WORTHINGTON J *see* Hajeer A; Hajeer AH; John S; John SL; Marlow A
- WRIGGLESWORTH JM *see* Mahmud T
- WRIGHT C *see* Barlow JH
- WRIGHT D *see* Kent RM; Walker DJ
- WRIGHT J, SAMANTA A. Survey of general practice requirements from the rheumatology services in Leicestershire, (A2) 29
- WRIGHT J *see also* Sheldon P
- WRIGHT MO *see* Bavington C
- WRIGHT P *see* Grabowski PS
- WRIGHT V *see* Troughton PR
- Wrist and hip fracture incidence reduction of an Asian population in Leicester, (A2) 3
- WULFFRAAT N. Familial presence of primary cryofibrinogenemia, report of three cases, (L) 102
- Xanthine oxidase inhibitors, rhTNF α -induced resorption of mouse calvariae *in vitro* is suppressed by, (A1) 241
- Y⁹⁰ synoviorthesis, effectiveness and response predictive factors, (A1) 208
- YACOUB M *see* Hawkins PN
- YAMANISHI Y. Atypical Cogan's syndrome associated with antineutrophil cytoplasmic autoantibodies, (L) 601
- YAMASHITA S *see* Okano K
- YANNI G, CAULI A, PITZALIS C, PANAYI G. CD1 expression in skin and synovial membrane of rheumatoid arthritis and psoriatic arthritis patients, (A1) 179
- YANNI G, RAMAGE W, PITZALIS C, HALL M, LANCHBURY J, PANAYI G. Th1 and Th2 cytokines in rheumatoid arthritis, (A1) 179
- YANNI G *see also* Iannone F; Johnson J
- YAQOOB M *see* Over KE
- YARWOOD H *see* Chapman PT; Mason JC
- YAZICI H, BAŞARAN G, HAMURYUDAN V, HIZLI N, YURDAKUL S, MAT C, TÜZÜN Y, ÖZYAZGAN Y, DIMITRIYADIS I. Ten-year mortality in Behçet's syndrome, 139
- YAZICI H *see also* Şahin (Özgün) Ş
- YE XJ *see* Terato K
- YEAP SS, AKIL M, HASSAN WU, ROY S, LIDDICOAT AJ, MOODY AR. Hand MRI changes in early rheumatoid arthritis, (A1) 118
- Pulmonary abnormalities in patients with early rheumatoid arthritis, (A2) 16
- YEAP SS, AKIL M, JONES AC, FLETCHER A, ROY S. Audit of muscle biopsies, (A1) 128
- YEAP SS, PEARSON D, CAWTE SA, HOSKING DJ. Ultrasound measurements in glucocorticoid treated patients, (A2) 5
- YEUNG VTF *see* Szeto CC
- YKL-40 serum levels in healthy children and adults. Comparison with serum and synovial fluid levels of YKL-40 in patients with osteoarthritis or trauma of the knee joint, 553
- YLI-JAMA T *see* Roivainen A
- YOCUM DE. Combination therapy with cyclosporin in rheumatoid arthritis, (S2) 19
- YOUNIOU P *see* Saraux A
- YOUNG A *see* Bodman-Smith K; Brown MA; Casey ATH; Dyce C; Pattison D; Prouse P; Talamo J
- YOUNG KA *see* Cox NL
- YOUNG SP *see* Carruthers DM
- YURDAKUL S *see* Yazici H
- ZANANIRI F *see* Creamer P
- ZDRAVKOVIC D *see* Poleksic L
- ZEIDLER H *see* Dreses-Werringloer U; Mau W
- ZHANG Z, FREARS ER, WINYARD PG, MURPHY G, O'CONNELL JP, BLAKE DR. Inactivation and fragmentation of tissue inhibitor of metalloproteinases (TIMP-1) by peroxynitrite, (A1) 69
- ZHANG Z *see also* Jawed S
- ZIAD F *see* Pina Salvador JM
- Zieve's syndrome case presenting with myalgia: not to be confused with polymyalgia rheumatica, (L) 495
- ZIGNEGO AL *see* Ferri C
- ZWINDERMAN AH *see* Vliet Vlieland TPM





